

## Predation on *Dorylus* army ants (Hymenoptera: Formicidae: Dorylinae) by *Agama agama* (Squamata: Agamidae) in Gabon

Thibaut DELSINNE<sup>1</sup>, Olivier S.G. PAUWELS<sup>2</sup>, Jean Pierre VANDE WEGHE<sup>3</sup> & Yves BRAET<sup>4</sup>

<sup>1</sup> Universidad Técnica Particular de Loja (UTPL), Departamento de Ciencias Naturales, Museo de Colecciones Biológicas (MUTPL), Loja, Ecuador (corresponding author: delsinthibaut@yahoo.fr).

<sup>2</sup> Vertébrés Récents, Institut royal des Sciences naturelles de Belgique, Rue Vautier 29, B-1000 Brussels, Belgium.

<sup>3</sup> BP 7847, Libreville, Gabon.

<sup>4</sup> Entomologie, Institut royal des Sciences naturelles de Belgique, Rue Vautier 29, B-1000 Bruxelles, Belgium; Unité d'Entomologie fonctionnelle et évolutive, Gembloux Agro-Bio Tech, Université de Liège, Passage des Déportés 2, B-5030 Gembloux, Belgium.

### Abstract

We report the first documented observations of predation on *Dorylus* (*Anomma*) *congolensis* Santschi, 1910 by the agamid lizard *Agama agama* (Linnaeus, 1758), the only reptile known to predate on this aggressive army ant. We also report two other observations of predation by Gabonese *Agama agama* on *Dorylus* sp. These observations confirm the extreme dietary adaptability of *Agama agama*.

**Keywords:** Africa, ants, ant-reptile interactions, natural history, predation.

*Dorylus* army ants are a noteworthy component of the Old World (Africa, Mediterranean Basin, Middle East and tropical Asia) entomofauna, especially in Africa where their species diversity is the highest (GOTWALD, 1995; KRONAUER, 2009). Species of the subgenus *Anomma* Shuckard, 1840 are particularly remarkable as their colonies can contain over ten million of adult individuals which predate a large array of prey<sup>1</sup>, mostly invertebrates, during massive swarm raiding on the soil surface and up into the vegetation (LEROUX, 1979; GOTWALD, 1995; KRONAUER, 2009). Although the sting of *Dorylus* is reduced and not functional (HERMANN, 1969; BOLTON, 1990), these ants possess a caste of large and aggressive workers with powerful disproportionate mandibles, which effectively defend the colony by the infliction of painful bites. Nonetheless, because colonies are conspicuous and composed of millions of nutrient- and protein-rich individuals (DEBLAUWE & JANSSENS, 2008; O'MALLEY & POWER, 2012), they represent an attractive food source for a wide diversity of animals.

The most studied predator of *Dorylus* is the chimpanzee, *Pan troglodytes* (Primates: Hominidae), which is renowned for using culturally transmitted harvesting techniques and tools to avoid being bitten (e.g. MÖBIUS *et al.*, 2008; RAUBENHEIMER & ROTHMAN, 2013; KOOPS *et al.*, 2015; and references therein). *Dorylus* also enters in the diet of various other African mammals (e.g. *Gorilla gorilla* (Primates: Hominidae) (DEBLAUWE & JANSSENS, 2008), *Manis* spp. (Pholidota: Manidae), *Bdeogale nigripes* (Carnivora: Herpestidae) (BEQUAERT, 1922), birds (e.g. *Alethe castanea* (Passeriformes: Muscicapidae), *Neocossyphus poensis* (Passeriformes: Turdidae)), skinks (*Trachylepis albilabris* (Squamata: Scincidae)), frogs (e.g.: *Amietophrynus* spp. (Anura: Bufonidae), *Hoplobatrachus occipitalis* (Anura: Dicroglossidae)) (BEQUAERT, 1922), and insects (e.g.: *Bengalia* spp. (Diptera: Calliphoridae), *Oecophylla longinoda* (Latreille, 1802) (Hymenoptera: Formicidae),

<sup>1</sup> We take this opportunity to mention here that the ants observed by PAUWELS *et al.* (2002) predated on the lamprophiid snake *Buroma depressiceps* belonged to the genus *Dorylus*.



Fig. 1. Predation by a subadult and a female *Agama agama* (lower and upper individual, respectively) on workers of *Dorylus (Anomma) congolensis* on 13 June 2006, at Pointe Wigombé, Pongara National Park, Gabon (photo Y. Braet).

other species of *Dorylus* (BEQUAERT, 1922; LEROUX, 1979)), sometimes as staple as indicated by the study of stomach contents of all the birds, frogs and mammals studied by BEQUAERT (1922). Here, we report a case of predation on *Dorylus (Anomma) congolensis* by *Agama agama* (Agamidae).

On 13 June 2006 at 10h40 at Pointe Wigombé, in Pongara National Park, Estuaire Province, northwestern Gabon, Yves Braet observed an adult female and a subadult agamas repeatedly catching workers from a large swarm of *Dorylus (Anomma) congolensis*, which was raiding along the bank of a lagoon (Fig. 1). The reptiles remained relatively still, even between prey catches, only a few centimeters away from the ant column. Vouchers of the ants collected at the Pointe Wigombé were deposited at the Oxford University Museum<sup>2</sup>. Subsequently, Jean-Pierre Vande weghe observed another *Agama agama* individual preying on *Dorylus* sp. in November 2012 in Minkébé National Park, Woleu-Ntem Province, northeastern Gabon. In the latter case, the agama was very cautious however, and after each catch it retreated about half a meter back. In both cases, the cost for the prey was certainly negligible as only a relatively tiny amount of workers was taken. The disappearance of the single queen of the *Dorylus* colony as a result of vertebrate predation is indeed exceptional (LEROUX, 1979).

*Agama agama* is widespread in West and Central Africa and feeds opportunistically on insects, small vertebrates (including cannibalism), flowers, fruits or human food reliefs (LOWES, 1954; HALSTEAD, 1970; BEIER, 1974; CLOUDSLEY-THOMPSON, 1981; PAUWELS *et al.*, 2004a, 2004b; PAUWELS & DAVID, 2008; LEACHÉ *et al.*, 2014). Faecal and stomach content analyses always confirmed that ants constitute an important part of its diet but, unfortunately, genera and species of the prey were in most cases not identified beyond the family level (SCHMIDT, 1919; HARRIS, 1964; CLOUDSLEY-THOMPSON, 1981; GUPTA, 1982; PAUWELS *et al.*, 2004a, 2004b; LUISELLI *et al.*, 2011; AKANI *et al.*, 2013). However, GRAMENTZ (1999) reported Gabonese *Agama agama* feeding on “Wanderameisen”, without specifying the generic identity of these ants; however he was actually referring to *Dorylus* (D. Gramentz, pers. comm.). These observations confirm *Agama agama* as an opportunistic, very adaptable predator, which explains its remarkable colonization abilities (PAUWELS *et al.*, 2004a, 2004b).

#### Acknowledgements

Thibaut Delsinne thanks the Secretary of Higher Education, Science, Technology and Innovation (SENESCYT) of the Government of Ecuador for financial support (Prometeo Grant) during the writing of this article. We thank Brian Taylor for his help in the identification of the ants, Dieter Gramentz for useful information, Roberto A.

<sup>2</sup> [http://antsafrica.org/ant\\_species\\_2012/dorylus/dorylus\\_congolensis/dorylus\\_congolensis.htm](http://antsafrica.org/ant_species_2012/dorylus/dorylus_congolensis/dorylus_congolensis.htm)

Pantaleoni for literature sending, and the ResearchGate platform for facilitating communication and knowledge sharing among scientists.

## References

- AKANI G.C., PETROZZI F., RUGIERO L., SEGNIAGBETO G.H. & LUISELLI L., 2013. - Effects of rainfall and geography on the comparative diets of eight rainbow lizard populations across Togo, Benin and Nigeria (West Africa). *Amphibia-Reptilia*, 34: 185-192.
- BEIER U., 1974. - Observations on the rainbow lizard. *Nigerian Field Journal*, 39: 137-140.
- BEQUAERT J., 1922. - The predaceous enemies of ants. *Bulletin of the American Museum of Natural History*, 45: 271-331.
- BOLTON B., 1990. - Army ants reassessed: the phylogeny and classification of the doryline section (Hymenoptera, Formicidae). *Journal of Natural History*, 24: 1339-1364.
- CLOUDSLEY-THOMPSON J.L., 1981. - Bionomics of the rainbow lizard *Agama agama* (L.) in eastern Nigeria during the dry season. *Journal of Arid Environments*, 4: 235-245.
- DEBLAUWE I. & JANSSENS G.P.J., 2008. - New insights in insect prey choice by chimpanzees and gorillas in southeast Cameroon: the role of nutritional value. *American Journal of Physical Anthropology*, 135: 42-55.
- GOTWALD W.H.Jr., 1995. - Army ants: The biology of social predation. Cornell University Press, Ithaca, 302 pp.
- GRAMENTZ D., 1999. - Zur Ökologie und Ethologie von *Agama agama* (Linnaeus, 1758) in Lambaréné, Gabun. *Salamandra*, 35(4): 193-208.
- GUPTA J.C., 1982. - Evaluation of *Agama agama* (Squamata, Agamidae) as a natural control agent of insect pests. *Zeitschrift für Angewandte Entomologie*, 93: 397-402.
- HALSTEAD L.B., 1970. - Some observations of the rainbow lizard, *Agama agama* (L.), at the University of Ife. *Nigerian Field Journal*, 35: 86-89.
- HARRIS V.A., 1964. - *The Life of the Rainbow Lizard*. Hutchison, London.
- HERMANN H.R., 1969. - The hymenopterous poison apparatus: evolutionary trends in three closely related subfamilies of ants (Hymenoptera: Formicidae). *Journal of the Georgia Entomological Society*, 4: 123-141.
- KOOPS K., SCHÖNING C., MCGREW W.C. & MATSUZAWA T., 2015. - Chimpanzees prey on army ants at Seringbara, Nimba Mountains, Guinea: predation patterns and tool use characteristics. *American Journal of Primatology*, 77: 319-329.
- KRONAUER D.J.C., 2009. - Recent advances in army ant biology (Hymenoptera: Formicidae). *Myrmecological News*, 12: 51-65.
- LEACHÉ A.D., WAGNER P., LINKEM C.W., BÖHME W., PAPPENFUSS T.J., CHONG R.A., LAVIN B.R., BAUER A.M., NIELSEN S.V., GREENBAUM E., RÖDEL M.-O., SCHMITZ A., LEBRETON M., INEICH I., CHIRIO L., OFORI-BOATENG C., ENIANG E.A., BAH EL DIN S., LEMMON A.R. & BURBRINK F.T., 2014. - A hybrid phylogenetic-phylogenomic approach for species tree estimation in African *Agama* lizards with applications to biogeography, character evolution, and diversification. *Molecular Phylogenetics and Evolution*, 79: 215-230.
- LEROUX J.M., 1979. - Sur quelques modalités de disparition des colonies d'*Anomma nigricans illiger* (Formicidae Dorylinae) dans la région de Lamto (Côte d'Ivoire). *Insectes Sociaux*, 26: 93-100.
- LUISELLI L., AKANI G.C., EBERE N. & PÉREZ-MELLADO V., 2011. - Stomach flushing affects survival/emigration in wild lizards: a study case with rainbow lizards (*Agama agama*) in Nigeria. *Amphibia-Reptilia*, 32: 253-260.
- LOWES R.H.G., 1954. - The rainbow lizard of West Africa. *Zoo Life*, 9: 27-28.
- MÖBIUS Y., BOESCH C., KOOPS K., MATSUZAWA T. & HUMLE T., 2008. - Cultural differences in army ant predation by West African chimpanzees? A comparative study of microecological variables. *Animal Behaviour*, 76: 37-45.
- O'MALLEY R.C. & POWER M.L., 2012. - Nutritional composition of actual and potential insect prey for the Kasekela chimpanzees of Gombe National Park, Tanzania. *American Journal of Physical Anthropology*, 149: 493-503.
- PAUWELS O.S.G & DAVID P., 2008 - Miscellanea Herpetologica Gabonica I. *Hamadryad*, 32: 13-18.
- PAUWELS O.S.G., GRAMENTZ D. & KAMDEM TOHAM A., 2004a - *Agama agama* (Red-headed Rock Agama). Nocturnal activity. *Herpetological Review*, 35: 164-165.
- PAUWELS O.S.G., BURGER M., GUIMONDOU S. & BRANCH W.R., 2004b. - *Agama agama* (Linnaeus, 1758), Red-headed Rock Agama. Nocturnal activity. *African Herp News*, 37: 20-21.
- PAUWELS O.S.G., KAMDEM TOHAM A. & CHIMSUNCHART C., 2002. - Recherches sur l'herpétofaune du Massif du Chaillu, Gabon. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 72: 47-57.
- RAUBENHEIMER D. & ROTHMAN J.M., 2013. - Nutritional ecology of entomophagy in humans and other primates. *Annual Review of Entomology*, 58: 141-160.
- SCHMIDT K.P., 1919. - Contributions to the herpetology of the Belgian Congo based on the collection of the American Museum Congo expedition, 1909-1915. Part I. Turtles, crocodiles, lizards, and chameleons. *Bulletin of the American Museum of Natural History*, 39: 385-624.