

A late Maastrichtian “out-of-the-ordinary” ophiuroid from The Netherlands

John W.M. JAGT & Mart J.M. DECKERS

Abstract

A tetramerous specimen of a new late Maastrichtian ophiuroid species from the Meerssen Member (Maastricht Formation) as exposed at the ENCI NV quarry, south of Maastricht, is described and briefly discussed. Non-pentamerous specimens of this species which dominates ophiuroid assemblages known from this member at this locality, are extremely rare.

Key-words: Late Maastrichtian, ophiuroids, pathology.

Résumé

Un exemplaire tétramérique d'une espèce nouvelle d'ophiure du Maastrichtien supérieur (Assise de Meerssen, Formation de Maastricht)

affleurant dans la carrière ENCI NV au sud de Maastricht, est décrite et brièvement discutée. Des exemplaires non-pentamériques de cette espèce, qui domine les assemblages d'ophiures dans cette assise à cet endroit, sont fort rares.

Mots-clés: Maastrichtien supérieur, ophiurides, pathologie.

Introduction

Recent field work at the Eerste Nederlandsche Cement Industrie (ENCI) NV quarry, south of Maastricht (The Netherlands), has yielded rich ophiuroid faunules from the upper part of the Meerssen Member (late Late Maastrichtian). Many well-preserved specimens of at least three species have been collected.




Fig. 1. — Ophiuridae (n. gen. ?) n. sp., late Late Maastrichtian, Meerssen Member (Maastricht Formation), ENCI NV quarry, Maastricht (The Netherlands), A - adoral aspect, B - oral aspect. Natuurhistorisch Museum Maastricht Collections, no. NHMM 1994675 (*ex* DECKERS Coll., no. 3239), x 12.5.

One of these, *Felderophiura vanderhami* JAGT, 1991, is comparatively rare, the dominant taxon being a superficially similar, undescribed species of ophiurid. Associated is an ?amphiurid species, which appears to be new as well. Taxonomical, palaeoecological and taphonomical aspects of these faunules will be published elsewhere (JAGT, in prep.). The fact that one of the ophiurid specimens before us is tetramerous and does not show any signs of predation warrants description.

Description and discussion

The present specimen (Fig. 1) is part of an ophiuroid faunule of some 40 specimens, recently collected from the upper part of the Meerssen Member (Maastricht Formation, late Late Maastrichtian, *Belemnella casimirovensis* Taxon Range Zone; see JAGT, in press), as exposed at the ENCI NV quarry, south of Maastricht.

Plotting the disc diameter of twenty-five specimens in the DECKERS Collection on a graph revealed that there are two size classes, one with diameters ranging between 5 and 6.1 mm, the other between 7.2 and 8.6 mm (a juvenile specimen in this lot measures 3.4 mm). The present tetramerous specimen belongs to the first size class, being 6 mm across.

This new species of ophiurid is easily distinguished from *Felderophiura vanderhami*, on details of adoral and oral disc plating. Typical features include a prominent primary circlet, well-developed but partially concealed radial shields, large interradial plates on disc margins, large adoral shields, single apical papillae, fairly many oral papillae, tentacle scales in proximal portions of arms and confinement of tentacle pores to proximal portions of arms.

Although the specimen before us (Fig. 1) lacks the centrodorsal plate, the primary circlet and some interradial plates, it is clear that it is perfectly tetramerous, and does not show any sign of accidental damage and regeneration. It can thus be assigned to Type I as distinguished by BOMWER & MEYER (1987). These authors described tetramerous and hexamerous specimens of the non-fissiparous Recent amphiurid *Amphiura filiformis* (O.F.

MÜLLER, 1776) and the fossil ophiurid *Ophiomusium gagnebini* (THURMANN, 1851) (see also MEYER, 1984). In discussing the occurrence of non-pentamerous, non-fissiparous fossil and modern ophiuroids, BOMWER & MEYER (1987, p. 899) distinguished two principal types of deviation; Type I, comprising "genuine alterations in total body symmetry" and Type 2, comprising "partial changes due to accidental causes and influenced possibly by regeneration."

BOMWER & MEYER (1987, p. 904) pointed out that symmetry deviation is common in fissiparous species (see also LAWRENCE, 1987), that their Recent and fossil tetramerous ophiuroids did not show any sign of recent regeneration of the disc and that it was possible that their type I symmetry deviation resulted from genetic factors which affected the early stages of development.

The fact that the present specimen (Fig. 1) does not display any trace of predation (see ARONSON, 1987 for a discussion of ophiuroid predation), shows that it is pathological and assignable to the above Type I.

Records in the literature of fossil and Recent non-pentamerous ophiuroids appear to be rare; MUNAR (1984, fig. 8) illustrated from off Mallorca a specimen of the non-fissiparous ophiomyxid species *Ophiomyxa pentagona* (LAMARCK, 1816), while KOEHLER (1922, pl. 84, fig. 1) figured a Recent tetramerous specimen of the ophiurid *Stegophiura sladeni* (DUNCAN, 1879) from the Pacific. The only other Cretaceous specimen we know of is that of SPENCER (1907, p. 105, pl. 28, fig. 4, 4a), who illustrated a small, tetramerous specimen (Natural History Museum London Collections, no. E.5058) which he assigned with a query to his new species *Ophiotitanos laevis* from the Cenomanian Lower Chalk of Dover, and about which he remarked that it was "peculiar in having only four arms, probably an abnormality."

Acknowledgements

We extend our best thanks to members of the "Vijlen Groep" for assistance during field work and, in particular, to Jef Reinders for preparing the photomicrographs, and to René Fraaye for an item of literature.

References

- ARONSON, R.B., 1987. Predation on fossil and Recent ophiuroids. *Paleobiology*, **13**: 187-192.
- BOMWER, T. & MEYER, C.A., 1987. Ophiurous oddities - deviation from pentamerous symmetry in non-fissiparous Ophiuroids: a comparison of recent and fossil species. *Eclogae geologicae Helvetiae*, **80**: 897-905.
- DUNCAN, P.M., 1879. On some Ophiuroidea from the Korean Seas. *Journal of the Linnean Society of London*, **14**: 445-482.
- JAGT, J.W.M., 1991. A new ophiuroid from the Maastrichtian type area (Late Cretaceous, SE Netherlands, NE Belgium). *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, **61**: 197-209.
- JAGT, J.W.M., in press. Late Maastrichtian and early Palaeocene key index macrofossils from the Maastrichtian type area (SE Netherlands, NE Belgium). *Geologie en Mijnbouw*.
- JAGT, J.W.M., in prep. Echinoderm extinction, survival and evolution across the K/T boundary in the type area of the Maastrichtian Stage (Late Cretaceous, The Netherlands).
- KOEHLER, R., 1922. Contributions to the biology of the Philippine Archipelago and adjacent regions. Ophiurans of the Philippine seas and adjacent waters. *Bulletin of the United States National Museum*, **100**: x + 1-486.
- LAMARCK, J.B.A.P.M. DE, 1816. *Histoire naturelle des animaux sans vertèbres*, 2. Verdière, Paris, 568 pp.
- LAWRENCE, J., 1987. A Functional Biology of Echinoderms. Croom Helm, London/Sydney, 340 pp.
- MEYER, C.A., 1984. Palökologie und Sedimentologie der Echinodermenlagerstätte Schafgraben (mittleres Oxfordian, Weissenstein, Kt. Solothurn). *Eclogae geologicae Helvetiae*, **77**: 649-673.
- MÜLLER, O.F., 1776. *Zoologicae Danicae Prodromus, seu Animalium Daniae et Norvegiae indigenarum Characteres, Nomena, et Synonyma imprimis popularium. Typis Hallageriis, Havniae*, xxxii + 282 pp.
- MUNAR, J., 1984. Anomalias en la simetria de los Asteroidea (Echinodermata). Casos observados en aguas de Mallorca.

Bulletin de Societat d'història Natural de les Balears, **28**: 59-66.

SPENCER, W.K., 1907. A monograph on the British fossil Echinodermata from the Cretaceous formations, II. The Asteroidea and Ophiuroidea. *Monograph of the Palaeontographical Society London*, **1907**: 91-132.

THURMANN, J., 1851. Abraham Gagnebin de la Ferrière. Fragment pour servir à l'histoire scientifique du Jura bernois et neuchâtelois pendant le siècle dernier. *Archives de la Société jurassique d'Emulation* (not seen).

Typescript received August 31, 1994

Corrected typescript received October 21, 1994

John W.M. JAGT
Dienst KCO
Natuurhistorisch Museum
Maastricht
Postbus 882
NL-6200 AW Maastricht
The Netherlands

Mart J.M. DECKERS
Boterbloemstraat 26
NL-5953 GJ Reuver
The Netherlands

