

Deep-water decapod crustacean fauna of the Eastern Ionian Sea

Chrissi-Yianna Politou¹, Porzia Maiorano², Gianfranco D'Onghia² and Chryssi Mytilineou¹

¹ Hellenic Centre for Marine Research, Aghios Kosmas, 16604 Helliniko, Greece

² Department of Zoology, University of Bari, Via Orabona 4, 70125 Bari, Italy

Corresponding author : C.-Y. Politou, e-mail : c-y@ncmr.gr

ABSTRACT. Knowledge on the decapod crustacean fauna of the E. Ionian Sea was enriched by a recent research program, carried out in deep waters (300-1200 m) of its northern part. The data were collected from a total of 148 hauls towed during four experimental trawl surveys from September 1999 to September 2000.

Thirty nine decapod species were identified, of which eight were Dendrobranchiata and 31 Pleocyemata (17 Caridea, 9 Brachyura, 3 Anomura, 1 Astacideum and 1 Palinurum). Concerning their depth distribution, 30 species were found in the depth zone 300-500 m, with *Parapenaeus longirostris* being the most abundant species. *Plesionika heterocarpus* and *P. antigai* followed in terms of abundance. Of the 27 species caught in the zone 500-700 m, *Aristaeomorpha foliacea* and *Plesionika martia* were the most abundant. In the zone 700-900 m, 19 species were found and *Aristaeomorpha foliacea* with *Aristeus antennatus* were the most numerous. Finally, the 18 decapod species encountered in the zone 900-1200 m showed low abundance, and *Sergia robusta* with *Polycheles typhlops* predominated in numbers.

From the identified decapods, *Acantheephyra eximia*, *Philoceras echinulatus* and *Pontophilus norvegicus* were mentioned for the first time in the E. Ionian Sea. Some other species, such as *Acantheephyra pelagica*, *Geryon longipes*, *Munida tenuimana*, *Paromola cuvieri*, *Parthenope macrochelos*, *Pasiphaea multidentata*, *Plesionika narval*, *Polycheles typhlops*, *Sergestes arachnoides* and *Sergestes arcticus* have been reported for the area only in the gray literature. Additionally, new depth distribution records for the Mediterranean were obtained for some species.

KEY WORDS : Decapoda, deep-water, Ionian Sea, Mediterranean.

INTRODUCTION

The literature on the decapod crustacean fauna of the Eastern Ionian Sea (Greece) is limited comparing to that referring to other Greek seas and more specifically to the Aegean Sea (e.g. THESSALOU-LEGAKIS & ZENETOS, 1985 ; THESSALOU-LEGAKIS, 1986 ; D'UDEKEM D'ACQZ, 1995). The systematic investigation of the decapod fauna of the Greek waters in the Ionian Sea has started quite recently in the framework of larger projects (MEDITS projects 1994-2001) or in restricted areas of the Ionian Sea, always at depths not exceeding 800 m ("Deep Water Fisheries" and RESHIO projects). Some information, obtained during the MEDITS project, is given by POLITOU et al. (1998, 2000).

The aim of the present work is to enrich the knowledge on the decapod crustacean fauna of the Eastern Ionian Sea with information obtained in the framework of the project INTERREG Italy-Greece, which was carried out in deep waters (300-1200 m) of its northern part. This information concerns the decapod faunistic composition, the species depth distribution, their frequency of occurrence and abundance.

MATERIALS AND METHODS

Sampling took place in depths from 300 to 1200 m of the Greek Ionian Sea during four experimental trawl surveys carried out from September 1999 to September 2000

(Fig. 1) using two chartered commercial trawlers. The gear used was a commercial bottom trawl with a cod end mesh size of 20 mm (side). The vertical and horizontal opening of the trawl were estimated, using a remote acoustic sensing system attached to the gear, as 1.5 m and 17 m respectively at a towing speed of 2.4 knots. The random stratified sampling design was applied using depth for the stratification of the study area. Three depth zones were defined, 300-500 m, 500-700 m and 700-900 m. Some additional experimental hauls were carried out in the depth zone 900-1200 m. A total of 148 hauls were carried out. The tow duration was 30 minutes for depths <500 m and one hour for depths >500 m. After each haul, catches were identified to species level. Species abundance was recorded on board. The mean abundance in number of individuals per fishing hour (CPUE) was estimated for each depth zone as :

$$CPUE = \frac{\sum N_n}{\sum t_n}$$

where $\sum N_n$ = sum of individuals of a species in the n hauls carried out in the depth zone and $\sum t_n$ = sum of fishing time of the n hauls in the depth zone.

RESULTS

Thirty nine decapod species were identified in total in the study area. Of them eight were Dendrobranchiata and 31 Pleocyemata (17 Caridea, 9 Brachyura, 3 Anomura, 1 Astacideum and 1 Palinurum) (Table 1).




Fig. 1. – Map of the study area showing the sampling stations.

Thirty species were found in the depth zone 300-500 m (6 Dendrobranchiata and 24 Pleocyemata : 14 Caridea, 6 Brachyura, 2 Anomura, 1 Astacideum and 1 Palinurum). *Parapenaeus longirostris* was the most important species, since it presented a remarkably high frequency of occurrence and abundance (Fig. 2). Although different species, such as *Plesionika antigai*, *P. gigliolii*, *P. heterocarpus*, *Nephrops norvegicus*, *Munida rullanti*, *P. edwardsii* and *Chlorotocus crassicornis*, were also found frequently, only *P. heterocarpus*, *P. antigai* and *P. edwardsii* were worth a mention in terms of their abundance.

Of the 27 species caught in the zone 500-700 m, six were Dendrobranchiata and 21 Pleocyemata : 9 Caridea, 7 Brachyura, 3 Anomura, 1 Astacideum and 1 Palinurum. *P. martia* was present in all stations, whereas other highly occurring species were, in order of importance, *Polycheles typhlops*, *Aristaeomorpha foliacea*, *P. acanthonotus*, *P. gigliolii*, *P. longirostris*, *N. norvegicus* and *P. antigai*. *Aristaeomorpha foliacea* and *Plesionika martia* were dominant in terms of abundance.

In the zone 700-900 m, 19 species were found (7 Dendrobranchiata and 12 Pleocyemata : 7 Caridea, 3 Brachyura, 1 Astacideum and 1 Palinurum). Many species, such as *A. foliacea*, *Polycheles typhlops*, *Aristeus antennatus*, *P. martia*, *Sergia robusta* and *P. acanthonotus*, showed high frequency of occurrence. However, their abundance was generally low with *A. foliacea* followed by *Aristeus antennatus* being the most numerous.

Finally, of the 18 decapod species encountered in the zone 900-1200 m, six were Dendrobranchiata and 12 Pleocyemata : 7 Caridea, 4 Brachyura and 1 Palinurum. *S. robusta* was the most frequently occurring species followed by *A. antennatus*, *P. typhlops*, *Acantheephyra pelagica*, *Pasiphaea multidentata*, *Sergestes arachnipodus*, *Acantheephyra eximia* and *P. acanthonotus*. Although, *Sergia robusta* and *Polycheles typhlops* predominated in numbers, their abundance was remarkably low.

DISCUSSION

Including results from the present study, the number of known decapods from the Ionian Sea has now reached 82 species. Of the identified decapods, *Acantheephyra eximia*, *Philoceras echinulatus* and *Pontophilus norvegicus* are reported for the first time in the E. Ionian Sea. *Acantheephyra eximia* was found in waters deeper than 800 m, which were not investigated earlier. Only one specimen of *Philoceras echinulatus* was caught in a single station. Finally, only one specimen of *Pontophilus norvegicus* was found in a deep station (965 m). Some other species, such as *Acantheephyra pelagica*, *Geryon longipes*, *Munida tenuimana*, *Paromola cuvieri*, *Parthenope macrochelos*, *Pasiphaea multidentata*, *Plesionika narval*, *Polycheles typhlops*, *Sergestes arachnipodus* and *Sergestes arcticus*, have been reported for the area only in the gray literature (Deep Water Fisheries technical report, unpublished data ; D'ONGHIA *et al.*, 2001 ; POLITOU *et al.*, 2001). Most of them are also uncommon species or mainly found in the two deepest strata. The two commercial deep-water shrimps *Aristaeomorpha foliacea* and *Aristeus antennatus* that were absent from the list of POLITOU *et al.* (1998) were found during the surveys of the Deep Water Fisheries project and later in the framework of the MEDITS project (KAPIRIS *et al.*, 1999 ; KAPIRIS & THESSALOU-LEGAKI, 2001 ; PAPA-KONSTANTINOY & KAPIRIS, 2001 ; CAU *et al.*, 2002). Most of the decapod species found in the present study, with the exception of *Acantheephyra eximia*, *Aegaeon lacazei*, *Munida rullanti*, *Paromola cuvieri*, *Philoceras echinulatus*, *Plesionika edwardsii*, *Plesionika narval* and *Pontophilus norvegicus*, were also found in the SE Adriatic Sea, which is adjacent to the E. Ionian (VASO & GJIKNURI, 1993 ; UNGARO *et al.*, 1999 ; MARSAN *et al.*, 2000). Comparing data from the rest of the Greek waters (Aegean and Cretan Sea) reveals that all the species found in the present study, with the exception of *Pontophilus norvegicus* and *Sergestes arachnipodus*, have been reported in the existing literature for these areas (KOUKOURAS *et al.*, 1992, 1997, 1998, 2000 ; KALLIANIOTIS *et al.*, 2000).

TABLE 1

List of decapod species collected in the E. Ionian Sea with species depth range and frequency of occurrence (%) per depth stratum.

No. of hauls	Depth range (m)	Depth strata (m)			
		300-500	500-700	700-900	900-1200
		45	52	38	13
Dendrobranchiata					
<i>Aristaeomorpha foliacea</i> (Risso, 1827)	388-1047	2.2	88.5	94.7	15.4
<i>Aristeus antennatus</i> (Risso, 1816)	480-1171	2.2	53.8	89.5	76.9
<i>Gennadas elegans</i> (Smith, 1882)	1082-1192				15.4
<i>Parapenaeus longirostris</i> (Lucas, 1846)	288-840	93.3	57.7	10.5	
<i>Sergestes arcticus</i> Krøyer, 1855	700-1192			2.6	23.1
<i>Sergestes arachnipodus</i> (Cocco, 1832)	318-1171	2.2	5.8	10.5	46.2
<i>Sergia robusta</i> (S.I. Smith, 1882)	480-1192	2.2	23.1	63.2	84.6
<i>Solenocera membranacea</i> (Risso, 1816)	322-823	6.7	1.9	2.6	
Pleocyemata					
Anomura					
<i>Munida intermedia</i> A. Milne Edwards & Bouvier, 1899	328-503	11.1	1.9		
<i>Munida rullanti</i> Zariquiey Alvarez, 1952	300-533	37.8	3.8		
<i>Munida tenuimana</i> G.O. Sars, 1872	518-518		1.9		
Astacidea					
<i>Nephrops norvegicus</i> (Linnaeus, 1758)	317-700	42.2	57.7	2.6	
Brachyura					
<i>Bathynectes longipes</i> (Risso, 1816)	620		1.9		
<i>Bathynectes maravigna</i> (Prestandrea, 1839)	322-1003	6.7	23.1	13.2	15.4
<i>Calappa granulata</i> (Linnaeus, 1758)	302-553	8.9	1.9		
<i>Geryon longipes</i> A. Milne Edwards, 1881	644-965		1.9	7.9	7.7
<i>Macropipus tuberculatus</i> (Roux, 1830)	343-462	11.1			
<i>Macropodia longipes</i> A. Milne Edwards & Bouvier, 1899	340-1003	2.2			7.7
<i>Monodaeus couchii</i> (Couch, 1851)	460-965	2.2	1.9		7.7
<i>Paromola cuvieri</i> (Risso, 1816)	597-742		7.7	2.6	
<i>Parthenope macrochelos</i> (Herbst, 1790)	302-582	11.1	1.9		
Caridea					
<i>AcanthePHYRA eximia</i> S.I. Smith, 1886	897-1047			2.6	38.5
<i>AcanthePHYRA pelagica</i> (Risso, 1816)	480-1192	2.2		15.8	61.5
<i>Aegaeon lacazei</i> (Gourret, 1887)	340-464	17.8			
<i>Alpheus glaber</i> (Olivi, 1792)	373	2.2			
<i>Chlorotocus crassicornis</i> (Costa, 1871)	300-614	35.6	3.8		
<i>Pasiphaea multidentata</i> Esmark, 1866	518-1171		9.6	28.9	53.8
<i>Pasiphaea sivado</i> (Risso, 1816)	377-1082	6.7	23.1	10.5	7.7
<i>Philoceras echinulatus</i> (M. Sars, 1861)	407	2.2			
<i>Plesionika acanthonotus</i> (Smith, 1882)	317-1047	13.3	86.5	44.7	30.8
<i>Plesionika antigai</i> Zariquiey Alvarez, 1955	288-700	77.8	34.6		
<i>Plesionika edwardsii</i> (Brandt, 1851)	305-700	35.6	23.1	2.6	
<i>Plesionika gigliolii</i> (Senna, 1903)	300-700	75.6	61.5		
<i>Plesionika heterocarpus</i> (Costa, 1871)	300-676	64.4	23.1		
<i>Plesionika martia</i> (A. Milne Edwards, 1883)	317-1085	15.6	100	65.8	15.4
<i>Plesionika narval</i> (Fabricius, 1787)	356	2.2			
<i>Pontophilus norvegicus</i> (M. Sars, 1861)	965				7.7
<i>Pontophilus spinosus</i> (Leach, 1815)	322-460	8.9			
Palinura					
<i>Polycheles typhlops</i> Heller, 1862	328-1171	17.8	98.1	92.1	76.9




Fig. 2. – Relative abundance of decapod species per depth stratum in the E. Ionian Sea. In parenthesis the mean number of specimens caught per hour (CPUE) is given.

Although comparison of abundance with other studies is difficult, because of the different types of gear and methods used, the general fauna distribution is quite similar to that found in other Mediterranean areas (ABELLO et al., 1988; CARTES & SARDA, 1992, 1993; CARTES et al., 1994; UNGARO et al., 1999; ABELLO et al., 2002). A dominance of shrimps in all depth strata observed in the present study was mentioned also by POLITOU *et al.* (1998) for the slope of the region. In relation to the western Mediterranean, our results are more comparable with those of ABELLO et al. (1988), whose samples were collected with a commercial trawl. A high presence of small species, such as *Calocaris macandreae* Bell, 1864 and *Processa noveli* Al-Adhub & Williamson, 1975, was mentioned by some authors (CARTES & SARDA, 1992; CARTES et al., 1994) for these depths in the Catalan Sea. These species were not found in the present study. This discrepancy can be explained by the smaller cod-end mesh size of the experimental trawl used in the former studies. Furthermore, although the species depth distribution observed in the present study was generally within the ranges given in the literature for the Mediterranean, in some cases new depth records were obtained (Table 2). More specifically, the distribution of the species *Aristaeomorpha foliacea*, *Bathynectes maravigna*, *Monodaeus couchii*, *Parapenaeus longirostris*, *Pasiphaea sivado*, *Plesionika antigai*, *Plesionika edwardsii* and *Plesionika martia* was extended into waters deeper than those

reported for the Mediterranean. Concerning *Bathynectes longipes*, it is considered a scarce species (ABELLO et al., 2001) and no depth distribution ranges are given for it.

A remarkable difference between the Greek Ionian Sea and the westernmost areas of the Mediterranean is the high abundance of *A. foliacea* in the former area and its absence or scarcity in the latter ones. This difference may be explained by the vulnerability of this species to overfishing (ORSI RELINI & RELINI, 1985; MATARRESE et al., 1997) in combination with the fishing pressure exercised in the deep waters of the westernmost areas. Such pressure is negligible in the Greek Seas. Furthermore, different hydrological conditions (i.e. salinity and temperature) between areas are reported as factors affecting the distribution of the species along the Mediterranean (RELINI & ORSI RELINI, 1987; MURENU et al., 1994). *A. foliacea* is considered to prefer waters of relatively high salinity and temperature such as those of the E. Ionian Sea in comparison to the westernmost Mediterranean areas (THEOCHARIS et al., 1993; HOPKINS, 1985).

A decline in the number of decapod species and of their abundance with depth was evident in the Ionian Sea. This pattern was observed also in the western Mediterranean (CARTES & SARDA, 1992, 1993; CARTES, 1993). The upper part of the slope (300-700 m) was characterized by a relatively high number of species found in abundance. *Parapenaeus longirostris*, *Aristaeomorpha foliacea*, *Ple-*

sionika martia, *P. heterocarpus*, *P. antigai* and *P. edwardsii* were the species that displayed the highest abundance values, and most of them are of important commercial value. Below 700 m of depth, the number of species and mainly their abundance decreased sharply, although red shrimps (mainly *A. foliacea* and secondarily *A. antenna-*

tus) were found in non negligible quantities in the zone 700-900 m. Given the relatively constant environmental conditions in these depths and the high oligotrophy of the area (STERGIOU et al., 1997), the main factor determining this reduction with depth seems to be the low trophic resource availability.

TABLE 2

Depth range of occurrence for decapod species in the Mediterranean updated with the results of the present study.

Species	Depth range (m)	
	Min	Max
<i>AcanthePHYra eximia</i>	421 (ABELLO et al., 2002)	2261 (CARTES, 1993)
<i>AcanthePHYra pelagica</i>	176 (ABELLO et al., 1988)	2261 (CARTES, 1993)
<i>Aegaeon lacazei</i>	13 (ABELLO et al., 1988)	1041 (CARTES, 1993)
<i>Alpheus glaber</i>	3 (ABELLO et al., 1988)	871 (ABELLO et al., 1988)
<i>Aristaeomorpha foliacea</i>	150 (MATARRESE et al., 1995)	1047 (present study)
<i>Aristeus antennatus</i>	150 (MATARRESE et al., 1995)	2200 (SARDÀ et al., 1994)
<i>Bathynectes longipes</i>		620 (present study)
<i>Bathynectes maravigna</i>	245 (ABELLO et al., 2001)	1003 (present study)
<i>Calappa granulata</i>	25 (ABELLO et al., 2002)	712 (ABELLO et al., 2002)
<i>Chlorotocus crassicornis</i>	55 (ABELLO et al., 1988)	742 (PIPITONE & TUMBILOLO, 1993)
<i>Gennadas elegans</i>	250 (KOUKOURAS et al., 1997)	2261 (CARTES, 1993)
<i>Geryon longipes</i>	439 (ABELLO et al., 1988)	1895 (CARTES, 1993)
<i>Macropipus tuberculatus</i>	48 (ABELLO et al., 1988)	748 (ABELLO et al., 1988)
<i>Macropodia longipes</i>	18 (ABELLO et al., 1988)	748 (ABELLO et al., 1988)
<i>Monodaeus couchii</i>	44 (ABELLO et al., 2002)	965 (present study)
<i>Munida intermedia</i>	35 (ABELLO et al., 1988)	871 (ABELLO et al., 1988)
<i>Munida rullanti</i>	40 (ABELLO et al., 2002)	587 (ABELLO et al., 2002)
<i>Munida tenuimana</i>	286 (ABELLO et al., 2002)	1899 (CARTES, 1993)
<i>Nephrops norvegicus</i>	58 (ABELLO et al., 2002)	871 (ABELLO et al., 1988)
<i>Parapenaeus longirostris</i>	26 (PIPITONE & TUMBILOLO, 1993)	840 (present study)
<i>Paromola cuvieri</i>	267 (PIPITONE & TUMBILOLO, 1993)	795 (PIPITONE & TUMBILOLO, 1993)
<i>Parthenope macrochelos</i>	20 (ABELLO et al., 1988)	655 (PIPITONE & TUMBILOLO, 1993)
<i>Pasiphaea multidentata</i>	128 (ABELLO et al., 1988)	2261 (CARTES, 1993)
<i>Pasiphaea sivado</i>	33 (ABELLO et al., 1988)	1082 (present study)
<i>Philoceras echinulatus</i>	55 (ABELLO et al., 1988)	871 (ABELLO et al., 1988)
<i>Plesionika acanthonotus</i>	141 (PIPITONE & TUMBILOLO, 1993)	1680 (CARTES, 1993)
<i>Plesionika antigai</i>	98 (ABELLO et al., 2002)	700 (present study)
<i>Plesionika edwardsii</i>	250 (GARCIA-RODRIGUEZ et al, 2000)	700 (present study)
<i>Plesionika gigliolii</i>	101 (ABELLO et al., 1988)	748 (ABELLO et al., 1988)
<i>Plesionika heterocarpus</i>	45 (ABELLO et al., 2002)	699 (ABELLO et al., 1988)
<i>Plesionika martia</i>	165 (ABELLO et al., 1998)	1085 (present study)
<i>Plesionika narval</i>	5 (THESSALOU-LEGAKI et al., 1989)	510 (POLITOU et al., 2000)
<i>Polycheles typhlops</i>	241 (ABELLO et al., 2002)	1927 (CARTES, 1993)
<i>Pontophilus norvegicus</i>	366 (ABELLO et al., 1998)	2261 (CARTES, 1993)
<i>Pontophilus spinosus</i>	69 (ABELLO et al., 2002)	871 (ABELLO et al., 1998)
<i>Sergestes arachnipedus</i>	279 (ABELLO et al., 2002)	1224 (CARTES, 1993)
<i>Sergestes arcticus</i>	160 (ABELLO et al., 2002)	2188 (CARTES, 1993)
<i>Sergia robusta</i>	220 (ABELLO et al., 1998)	2261 (CARTES, 1993)
<i>Solenocera membranacea</i>	3 (ABELLO et al., 1998)	871 (ABELLO et al., 1998)

ACKNOWLEDGEMENTS

The authors are grateful to their colleagues who participated in the sampling cruises and in the laboratory work. Special thanks to Stefanos Kavadas and John Dokos for their help in the data base management and the map drawing.

The project "INTERREG II Greece-Italy" was financially supported by the Greek Ministry of Economy and the EC.

REFERENCES

ABELLO, P., A. CARBONELL & P. TORRES (2002). Biogeography of epibenthic crustaceans on the shelf and upper slope off the

- Iberian Peninsula Mediterranean coasts : implications for the establishment of natural management areas. *Sci. Mar.*, 66 (Suppl. 2) : 183-198.
- ABELLO, P., N. UNGARO, C.-Y. POLITOU, P. TORRES, E. ROMAN, P. RINELLI, P. MAIORANO & G. NORRITO (2001). Notes on the distribution and biology of the deep-sea crab *Bathynectes maravigna* (Brachyura : Portunidae) in the Mediterranean Sea. *Hydrobiologia*, 449 : 187-192.
- ABELLO, P., F.J. VALLADARES & A. CASTELLON (1988). Analysis of the structure of decapod crustacean assemblages off the Catalan coast (North-West Mediterranean). *Mar. Biol.*, 98 : 39-49.
- CARTES, J.E. (1993). Deep-sea decapod fauna of the Western Mediterranean : bathymetric distribution and biogeographic aspects. *Crustaceana*, 65(1) : 29-40.
- CARTES, J.E., J.B. COMPANY & F. MAYNOU (1994). Deep-water decapod crustacean communities in the Northwestern Mediterranean : influence of submarine canyons and season. *Mar. Biol.*, 120 : 221-229.
- CARTES, J.E. & F. SARDA (1992). Abundance and diversity of decapod crustaceans in the deep-Catalan Sea (Western Mediterranean). *J. nat. Hist.*, 26 : 1305-1323.
- CARTES, J.E. & F. SARDA (1993). Zonation of deep-sea decapod fauna in the Catalan Sea (Western Mediterranean). *Mar. Ecol. Prog. Ser.*, 94 : 27-34.
- CAU A., A. CARBONELL, M.C. FOLLESA, A. MANNINI, G. NORRITO, L. ORSI-RELINI, C.-Y. POLITOU, S. RAGONESE & P. RINELLI (2002). MEDITS-based information on the deep-water red shrimps *Aristaeomorpha foliacea* and *Aristeus antennatus* (Crustacea : Decapoda : Aristeidae). *Sci. Mar.*, 66 (Suppl.2) : 103-124.
- D'ONGHIA, G., C.-Y. POLITOU, F. MASTROTOTARO, CH. MYTILINEOU & A. MATARRESE (2001). Biodiversity from the upper slope demersal community of eastern Mediterranean : preliminary comparison between two areas with and without fishing impact. *NAFO Scientific Council Meeting-September 2001*, SCR Doc. 01/135 : 11p.
- GARCIA-RODRIGUEZ, M., A. ESTEBAN & J.L. PEREZ GIL (2000). Considerations on the biology of *Plesionika edwardsi* (Brandt, 1851) (Decapoda, Caridea, Pandalidae) from experimental trap catches in the Spanish western Mediterranean Sea. *Sci. Mar.*, 64(4) : 369-379.
- HOPKINS, T.S. (1985). Physics of the Sea. In : MARGALEF, R. (ed.), *Key environments : Western Mediterranean*, Pergamon Press, New York : 100-125.
- KALLIANIOTIS, A., K. SOPHRONIDIS, P. VIDORIS, A. TSELEPIDES (2000). Demersal fish and megafaunal assemblages on the Cretan continental shelf and slope (NE Mediterranean) : seasonal variation in species density, biomass and diversity. *Progress in Oceanography*, 46 : 429-455.
- KAPIRIS, K. & M. THESSALOU-LEGAKI (2001). Sex-related variability of rostrum morphometry of *Aristeus antennatus* (Decapoda : Aristeidae) from the Ionian Sea (Eastern Mediterranean, Greece). *Hydrobiologia*, 449 : 123-130.
- KAPIRIS, K., M. THESSALOU-LEGAKI, M. MORAITOU-APOSTOLOPOULOU, G. PETRAKIS & C. PAPACONSTANTINOU (1999). Population characteristics and feeding parameters of *Aristaeomorpha foliacea* and *Aristeus antennatus* (Decapoda : Aristeidae) from the Ionian Sea (Eastern Mediterranean). In : VON VAUPEL KLEIN, J.C. & F.R. SCHRAM (eds), *The biodiversity crisis and Crustacea : Proceedings of the fourth international crustacean congress, Amsterdam, July 20-24, 1998*, A.A. Balkema, Rotterdam : 177-191.
- KOUKOURAS, A., S. DOULGERAKI & M.-S. KITSOS (2000). Notes on the vertical distribution of pelagic shrimps (Decapoda, Natantia) in the Aegean Sea. *Crustaceana*, 73(8) : 979-993.
- KOUKOURAS, A., S. DOULGERAKI, M.-S. KITSOS & C. PAPACONSTANTINOU (1997). The vertical distribution of the pelagic shrimps (Crustacea, Decapoda, Natantia) of the North Aegean Sea. In : POLITOU, C.-Y., M. STOUMBOUDI & D. SAKELLARIOU (eds), *Proc. 5th Hel. Symp. Oceanogr. & Fish.*, NCMR, Athens, vol. II : 91-94.
- KOUKOURAS, A., C. DOUNAS, M. TÜRKAY & E. VOULTSIADOU-KOUKOURA (1992). Decapod crustacean fauna of the Aegean Sea : new information, check list, affinities. *Senckenbergiana marit.*, 22(3/6) : 217-244.
- KOUKOURAS, A., A. KALLIANIOTIS & D. VAFIDIS (1998). The decapod crustacean genera *Plesionika* Bate (Natantia) and *Munida* Leach (Anomura) in the Aegean Sea. *Crustaceana*, 71(6) : 714-720.
- MARSAN, R., N. UNGARO, C.A. MARANO & M.C. MARZANO (2000). Remarks on distribution and fishery biology of some *Plesionika* species (Decapoda, Pandalidae) in the southern Adriatic basin (Mediterranean Sea). In : VON VAUPEL KLEIN, J.C. & F.R. SCHRAM (eds), *The biodiversity crisis and Crustacea : Proceedings of the fourth international crustacean congress, Amsterdam, July 20-24, 1998, volume 2*, A.A. Balkema, Rotterdam : 763-769.
- MATARRESE, A., G. D'ONGHIA, M. DEFLORIO, M. PANZA & G. COSTANTINO (1995). Recenti acquisizioni sulla distribuzione batimetrica di *Aristaeomorpha foliacea* e *Aristeus antennatus* (Crustacea, Decapoda) nel Mar Ionio. *Biol. Mar. Medit.*, 2(2) : 299-300.
- MATARRESE, A., G. D'ONGHIA, A. TURSI & P. MAIORANO (1997). Vulnerabilità e resilienza in *Aristaeomorpha foliacea* (Risso, 1827) e *Aristeus antennatus* (Risso, 1816) (Crustacei, Decapodi) nel Mar Ionio. *S.It.E. Atti*, 18 : 535-538.
- MURENU, A., D. CUCCU, C. FOLLESA, A. SABATINI & A. CAU (1994). The occurrence of *Aristaeomorpha foliacea* in Sardinia waters. In : BIANCHINI, M. & S. RAGONESE (eds), *Life cycles and fisheries of the deep-water red shrimps Aristaeomorpha foliacea and Aristeus antennatus*. N.T.R.-I.T.P.P. Special publication, 3 : 49.
- ORSI RELINI, L. & G. RELINI (1985). The red shrimps fishery in the Ligurian Sea : Mismanagement or not ? *FAO Fish. Rep.*, 336 : 99-106.
- PAPACONSTANTINOU, C. & K. KAPIRIS (2001). Distribution and population structure of the red shrimp (*Aristeus antennatus*) on an unexploited fishing ground in the Greek Ionian Sea. *Aquat. Living Resour.*, 14 : 303-312.
- PIPITONE, C. & M.L. TUMBILOLO (1993). Decapod and stomatopod crustaceans from the trawlable bottoms of the Sicilian Channel (Central Mediterranean Sea). *Crustaceana*, 65(3) : 358-364.
- POLITOU, C.-Y., M. KARKANI & J. DOKOS (1998). Distribution of decapods caught during MEDITS surveys in Greek waters. In : BERTRAND, J.A. & G. RELINI (eds), *Demersal resources in the Mediterranean*, Ifremer-Actes de Colloques, 26 : 196-207.
- POLITOU, C.-Y., M. KARKANI & J. DOKOS (2000). Distribution of pandalid shrimps in Greek waters (Ionian Sea and Argosaronikos). In : LABROPOULOU, M. (ed.), *Proc. 6th Hel. Symp. Oceanogr. & Fish.*, NCMR, Athens, vol. II : 61-66.
- POLITOU, C.-Y., S. KAVADAS, CH. MYTILINEOU, A. TURSI, G. LEMBO & R. CARLUCCI (2001). Fisheries resources in the deep waters of eastern Mediterranean (Greek Ionian Sea). *NAFO Scientific Council Meeting-September 2001*, SCR Doc. 01/104, 12p.
- RELINI, G. & L. ORSI RELINI (1987). The decline of red shrimps stocks in the Gulf of Genoa. *Inv. Pesq.*, 51(suppl. 1) : 245-360.
- SARDA, F., J.E. CARTES & W. NORBIS (1994). Spatio-temporal structure of the deep-water shrimp *Aristeus antennatus* population in the Western Mediterranean. *Fish. Bull.*, 92 : 599-607.
- STERGIOU, K.I., E.D. CHRISTOU, D. GEORGOPOULOS, A. ZENETOS & C. SOUVERMEZOGLU (1997). The Hellenic Seas : Physics, Chemistry, Biology and Fisheries. In : ANSELL, A.D., R.N.

- GIBSON & M. BARNES (eds), *Oceanography and Marine Biology : an Annual Review*, UCL Press, 35, 415-538.
- THEOCHARIS, A., D. GEORGOPOULOS, A. LASKARATOS & K. NITIS (1993). Water masses and circulation in the central region of the Eastern Mediterranean : Eastern Ionian, South Aegean and Northwest Levantine, 1986-1987. In : ROBINSON, A.R. & P. MALANOTTE-RIZZOLI (eds), *Topical studies in Oceanography, Part. II. Deep-Sea Res.*, 40(6) : 1121-1142.
- THESSALOU-LEGAKIS, M. (1986). Preliminary data on the occurrence of Thalassinidea (Crustacea, Decapoda) in the Greek seas. *Biologia Gallo-hellenika*, 12 : 181-187.
- THESSALOU-LEGAKI, M., A. FRANTZIS, K. NASSIOKAS & S. HATZINIKOLAOU (1989). Depth zonation in a *Parapandalus narval* (Crustacea, Decapoda, Pandalidae) population from Rhodes Island, Greece. *Estuar. coast. shelf Sci.*, 29 : 273-284.
- THESSALOU-LEGAKIS, M. & A. ZENETOS (1985). Autoecological studies on the Thalassinidea (Crustacea, Decapoda) of the Patras Gulf and Ionian Sea (Greece). *Rapp. Comm. int. Mer Médit.*, 29(5) : 309-312.
- UDEKEM D' ACOZ, C. D' (1995). Contribution à la connaissance des crustacés décapodes helléniques II : Penaeidea, Stenopodidea, Palinuridea, Homaridea, Thalassinidea, Anomura, et note sur les stomatopodes. *Bios (Macedonia, Greece)*, 3 : 51-77.
- UNGARO, N., C.A. MARANO, R. MARSAN, M. MARTINO, M.C. MARZANO, G. STRIPPOLI & A. VLORA (1999). Analysis of demersal species assemblages from trawl surveys in the South Adriatic Sea. *Aquat. Living Resour.*, 12(3) : 177-185.
- VASO, A. & L. GJIKNURI (1993). Decapod crustaceans of the Albanian coast. *Crustaceana*, 65(3) : 390-407.