

Taphrocampa Gosse, 1951

Taphrocampa annulosa Gosse, 1851

- Palaearctic region: Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960); Romania (Rudescu, 1960); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Impoundment Neunzehnhain, Germany (Althaus, 1976); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Coast of S Finland (Eriksen, 1969); Croatia, ex-Yugoslavia (Erben, 1987); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); District Ostrava, CSR (Dvorakova, 1960b); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Achmer Grasmoor, NW Germany (Koste, 1968b); All over Europe. Cosmopolitan (Bērziņš, 1978); Lake Balaton, Hungary (Zankai, 1968); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Peat bog Chambedaze, Massif Central, France (Francez, 1982); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Water courses in the Bucegi mts., Romania (Godeanu, 1963b); Small permanent ponds in the Bucegi mts., Rom. (Godeanu, 1969a); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Dübener heide, N Germany (Wulfert, 1960a, 1960b & 1961); Nat. Res. "Het Hol" near Kortenhoef, NL (Dresscher, 1976); In reed borders of the Danube delta, Romania (Rudescu, 1968); Lao-Shan mountains, China (Wulfert, 1968); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)
- Nearctic region: Lac Echo, Québec, Canada (Nogrady, 1976); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980); Kingston, Ontario, Canada (Nogrady, 1989); Canada (Chengalath, 1977); Wisconsin, USA (Bērziņš, 1984); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Laurentian Great Lakes (Stemberger, 1979)
- Oriental region: Singapore (Karunakaran & Johnson, 1978); Malaysia & Singapore (Suzuki, 1989)
- Australian region: NW Aust.: Magela Creek area (Koste, 1981b); Alligator river area, NT, Aust. (Tait, Shiel & Koste, 1984); Creek & river in Aust. (Bērziņš, 1982); NSW & Qld, Aust. (Shiel & Koste, 1979); NSW, NT, Qld & Vic, Aust. (Koste & Shiel, 1991); West Canterbury, New Zealand (Russell, 1960)
- Neotropical region: Brazil (Koste & José de Paggi, 1982); Varzea near Santarem, Brazil (Koste, 1974b); Mato Grosso, Brazil (Turner & Da Silva, 1992)

Taphrocampa clavigera Stokes, 1896

- Palaearctic region: Romania (Rudescu, 1960); Peat bogs in the Massif Central, France (Batut, 1965); C Eur. mts., Carpathes, C Eur. plains, Baltic province, UK (Bērziņš, 1978); Peat bogs in the Auvergne, France (Francez, 1981); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966)
- Nearctic region: N America (Bērziņš, 1978)
- Neotropical region: Brokopondo Lake, Surinam (Leentvaar, 1979); Surinam (Koste & José de Paggi, 1982)

Taphrocampa selenura Gosse, 1887

- Palaearctic region: Garden ponds in Tokyo, Japan (Suzuki, 1975); Lago Trasimeno, Italy (Taticchi, 1968); River Ybbarspå, Sweden (Bērziņš, 1979); Romania (Rudescu, 1960); Coast of S Finland (Eriksen, 1969); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); ? District Ostrava, CSR (Dvorakova, 1960b); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, Germany (Koste, 1965); Achmer Grasmoor, NW Germany (Koste, 1968b); All over Europe (Bērziņš, 1978); Southwest islands of Japan (Suzuki, 1990; Suzuki *et al.*, 1992); River Elbe estuary, Germany (Schulz, 1961); Anatolia, Turkey (Segers *et al.*, 1992); Dübener Heide, N Germany (Wulfert, 1960a & 1960b); Bog pond near Budapest, Hungary (Nogrady, 1962); Luga district of Leningrad area, R.F. (Kutikova, 1962b); In reed border of the Danube delta, Romania (Rudescu, 1968); Ponds in the Paris suburb, France (Pourriot, 1965)
- Nearctic region: Canada (Chengalath, 1977); Wisconsin, USA (Bērziņš, 1984); NW Canada (Chengalath & Koste, 1987); Arctic N America (Chengalath & Koste, 1989); Mt Tremblant Prov. park, Québec, Canada (Nogrady, 1980); Laurentian Great Lakes (Stemberger, 1979); Mirror Lake, New Hampshire, USA (Strayer, 1985)
- Oriental region: Impoundment Kinda, C Burma (Koste & Tobias, 1990)
- Australian region: Tasmania (Koste & Shiel, 1987); NSW, Aust. (Shiel & Koste, 1979); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, N Aust. (Koste & Shiel, 1983a; Tait, Shiel & Koste, 1984); Rare. NSW & NT (Koste & Shiel, 1991); Canterbury, New Zealand (Russell, 1960)
- Neotropical region: Ecuador: Oriente & Costa (Koste & Böttger, 1989); Ecuador: Oriente, Sierra & Costa (Koste & Böttger, 1992); Chile (Löffler, 1962); Chilean lakes (Schmid-Araya, 1991); Brazil, Argentina & Chile (Koste & José de Paggi, 1982); Amazonia, Brazil (Koste, 1972f); Varzea near Santarem, Brazil (Koste, 1974b); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Concepcion, Paraguay (Koste, 1986b); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Prov. Santa Fé, Argentina

(José de Paggi, 1990); Mato Grosso, C Brazil (Turner & Da Silva, 1992)

Taphrocampus spec.

- Palaearctic region: Periphytic biota in sewages & park waters, Jap. (Suzuki, 1981)
- Nearctic region: Bayou de Siard, Monroe, Louisiana, USA (Gallagher, 1965); Mirror Lake, New Hampshire, USA (Strayer, 1985)
- Oriental region: Singapore (Suzuki, 1991)

***Testudinella* Bory de St. Vincent, 1826**

***Testudinella ahlstromi* Hauer, 1956**

- Palaearctic region: Southwest islands of Japan (Suzuki *et al.*, 1992)
- Nearctic region: N Carolina, USA (Koste, 1972f)
- Australian region: S Tasmania (Koste, Shiel & Tan, 1988); Tasmania (Shiel, Koste & Tan, 1989)
- Neotropical region: Amazonia, Brazil (Thomasson, 1971); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Lago Paroni, Brazil (Koste, 1974a); Amazonia, Brazil & Venezuela (Koste, 1972f); Varzea near Santarem, Brazil (Koste, 1974b); Venezuela & Brazil (Koste & José de Paggi, 1982); Near Concepcion, Paraguay (Koste, 1986b); Ecuador: Oriente & Costa (Koste & Böttger, 1989); Ecuador: Oriente (Koste & Böttger, 1992); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Province Santa Fé, Argentina (José de Paggi, 1990); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Laguna Cocha Cashu, Amazonia, Brazil (Schaden, 1977c)

***Testudinella amphora* Hauer, 1938**

- Palaearctic region: Southwest islands of Japan (Suzuki *et al.*, 1992)
- Oriental region: Indonesia (Suzuki, 1989)
- Australian region: Tasmania (Koste & Shiel, 1987); ? Farm dams near Gloucester, NSW, Aust. (Suzuki & Timms, 1980); NT, Aust. (Koste & Shiel, 1990c); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984)
- Neotropical region: Amazonia, Brazil (Thomasson, 1971); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Varzea near Santarem, Brazil (Koste, 1974b); Brazil (Koste & José de Paggi, 1982); NW Brazil: Magela Creek area (Koste, 1981b)

***Testudinella angulata* Myers, 1934**

- Nearctic region: Northern Michigan inland lakes (Stemberger, 1990)

***Testudinella bidentata* (Ternetz, 1892)**

- See *T. parva* f. *bidentata* (Ternetz, 1892)

***Testudinella brevicaudata* Yamamoto, 1951**

- Palaearctic region: Japanese inland waters (Yamamoto, 1960a); Japan (Sharma, 1990)
- Oriental region: Kalimantan (Borneo), Indonesia (Koste, 1988a); Indonesia (Suzuki, 1989); New for India: Ganhati city, Assam State. (Sharma, 1990); Lake Manguao, Palawan, The Philippines (Davies & Green, 1990)

***Testudinella brycei* Hauer, 1938**

- Palaearctic region: Japanese inland waters (Yamamoto, 1960a); C Eur. plain (Bērziņš, 1978)
- Oriental region: Kalimantan (Borneo), Indonesia (Koste, 1988a); Indonesia (Suzuki, 1989)
- Neotropical region: Surinam (Leentvaar, 1975); Brokopondo Lake, Surinam (Leentvaar, 1979); Argentina & Surinam (Koste & José de Paggi, 1982); Province Santa Fé, Argentina (José de Paggi, 1990)

***Testudinella caeca* (Parsons, 1892)**

- Palaearctic region: Hortobagy Nat. Park, Hungary (Forró & Ronkay, 1983); Nat. Res. Jursky Sur, CSR (Hudec, 1978); Darnsee near Epe, NW Germany (Koste, 1962); River Hase, NW Germany (Koste, 1976b); W, C & E Eur. plains, Baltic province (Bērziņš, 1978); Peat bogs Chambéthane & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Luga district in the Leningrad area, R.F. (Kutikova, 1962b); Serbian section of river Danube (Zivkovic, 1987)

- Nearctic region: Ross Lake, Florida, USA (Bienert, 1986)

- Neotropical region: Argentina (Koste & José de Paggi, 1982)

***Testudinella caeca* (Parsons), f. *haueri* Wiszniewski, 1954**

- Palaearctic region: E Eur. plain (Bērziņš, 1978)

***Testudinella caeca* (Parsons), f. *lermaensis* Ahlstrom, 1932**

- Neotropical region: Province Rio Negro, Argentina (José de Paggi, 1990); Araucarian lakes, E of the Andes, Chile (Thomasson, 1963b)

***Testudinella carlini* Bartos, 1951**

- Palaearctic region: Rivers in Soviet Karelia, R.F. (Filimonova, 1976); N, W & C Iceland (De Ridder, 1966a, 1967a, 1969b, 1970, 1972); C Eur. plain, Iceland (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1960, 1968 & 1970); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); ? Lakes in the Leczna-Włodawa area, & "Plaine de la Grande Pologne" (Radwan, 1967)

***Testudinella clypeata* (O.F. Müller, 1786)**

- Palaearctic region: Romania (Rudescu, 1960); Sea & brackish water pools, S Finland (Björklund, 1972a); Brackish water pools in Norway (Eriksen, 1968); Coast of S Finland (Eriksen, 1969); Hortobagy Nat. Park, Hungary (Forro & Ronkay, 1983); All over Europe; Caspian area (Bērziņš, 1978); Rügener, Hiddenseer & Darszer Boddengewässer, Germany (Schwarz, 1963); Lake Balaton, Hungary (Zankai, 1968); Gulf of Riga, Baltic Sea, Belt Sea, Kattegat, Skagerak, N & S North Sea, E & W English Channel, Bristol Channel & Irish Sea, S & W Ireland, Norwegian Sea (Bērziņš, 1960); Camargue, S France (De Ridder, 1960a & 1961); Marismas, Spain (De Ridder, 1963b); W Iceland (De Ridder, 1969b, 1970 & 1972); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Oostduinkerke & Oostende, Belgium (De Ridder, 1986); Channel Nieuwpoort-Duinkerke in Oostduinkerke, Belgium (De Maeseneer, 1979); Roscoff, Atlantic coast of Brittany, France (D'Hondt, 1970); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Zuiderzee; r. Tjunmer near Tjun; channel Harlingervaat; near Midlum, NL (Dresscher, 1976); Salt marsh near Puerta da Pollense, Mallorca, Sp. (De Ridder, 1967c); Black Sea (Rudescu, 1961); Ponds in Oltina, Dobrogea, Romania (Popescu-Gorj & Costea, 1961); One brackish & 2 freshwater localities in Poland (Radwan, 1967); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Scandinavian coastal waters (Thane-Fenchel, 1968); Spain: Barcelona & Huelva; Balearics: Mallorca (Velasco, 1990); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)

- Nearctic region: Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 8)

***Testudinella clypeata* (O.F. Müller), f. *crassa* (Levander, 1892)**

- Palaearctic region: Gulf of Finland & N North Sea (Bērziņš, 1960); Baltic prov., Boreal highl. & NW Russian taiga (Bērziņš, 1978)

***Testudinella discoidea* Ahlstrom, 1938**

- Palaearctic region: River Loire, France (Lair, 1980); Middle course of river Loire, France (Lair & Sargas, 1981)

***Testudinella elliptica* (Ehrenberg, 1834)**

- Palaearctic region: Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); River Danube (Enaceanu & Brezeanu, 1970); Boom, Belgium (De Ridder, 1986); Romania (Rudescu, 1960); Coast of S Finland (Eriksen, 1969); Hortobagy Nat. Park, Hungary (Forro & Ronkay, 1983); Nat. Res. Jursky Sur, CSR (Hudec, 1978); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Darnsee near Epe, NW Germany (Koste, 1962); River Hase, NW Germany (Koste, 1976b); Lake Alfsee, NW Germany (Koste & Poltz, 1987); Fishponds near Warsaw, Poland (Hillbricht-Illkowska, 1964); Almost all over Europe (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1970); Rügener, Hiddenseer & Darszer Boddengewässer, Germany (Schwarz, 1963); Lake Balaton, Hungary (Zankai, 1968); Latorica, E Slovakia lowland, CSR (Terek, 1990a); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Japanese inland waters (Yamamoto, 1960a); Small water bodies in the Mikolajkie area, Poland (Klimowicz, 1967); Lowland brook-pond system in Boom, Belgium (Maes, 1985); Shores of embanked and free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); Prov. Noord-Holland; channel Noordzeekanaal near the Western Docks, A'dam; De Linde; Channel Zuidwillemsevaart between 's Hertogenbosch & Veghel, NL (Dresscher, 1976); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); In reed border of the Danube delta, Romania (Rudescu, 1968); One brackish & several freshw. areas in Poland (Radwan, 1967); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Ponds in Botanical Garden, Gent, Belgium (Segers *et al.*, 1991b); Serbian section of river Danube (Zivkovic, 1987); Spain: Tarragona (Velasco, 1990); Wiszni III Pond, Poland (Lewkowicz, 1972)

- Nearctic region: Canada (Chengalath, 1977); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988)

- Oriental region: Sri Lanka (Mendis & Fernando, 1962; Fernando, 1969)

- Neotropical region: Ecuador: Oriente (Koste & Böttger, 1989)

***Testudinella emarginula* (Stenroos, 1898)**

- Palaearctic region: L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960); Id. + Nat. Res. "Het Hol" near Kortenhoef, NL (Dresscher, 1976); N & W Iceland (De Ridder, 1969b, 1970 & 1972); Duszatynskie lakes, Poland (Kowalczyk & Radwan, 1962); Parczew area peat bogs, Poland (Radwan, 1974); Bieszcza, Poland (Radwan, 1967); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Impoundment in Spain (Velasco, 1990); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Romania (Rudescu, 1960); Small lakes in Estonia (Haberman & Mäemets, 1987); River

Ybbarspå, Sweden (Bērziņš, 1979); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Coast of S Finland (Eriksen, 1969); Asiatic Turkey (Dumont & De Ridder, 1987); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); N Swedish Lapland (Pejler, 1962a); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Achmer Grasmoor, NW Germany (Koste, 1968b); All over Europe (Bērziņš, 1978); Camargue, S France (De Ridder, 1960a & 1961); N & W Iceland (De Ridder, 1970 & 1972); Lake Neusiedlersee, Austria (Donner, 1979)

- Nearctic region: Wisconsin, USA (Bērziņš, 1984); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Lac Echo, Québec, Canada (Nogrady, 1976); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980); Jenny Lake, W North America (Thomasson, 1962, 1982a)

- Oriental region: Delhi, India (Sarma, 1988); Danau Teleko, near Palembang, Sumatra (Bērziņš); India: Delhi, Meghalaya State, Assam State & Dharamnayar in N tripura. Also in Orissa & W Bengal (Sharma, 1990); Impoundment Kinda, C Burma (Koste & Tobias, 1990)

- Australian region: Vic, Aust. (Koste & Shiel, 1980c); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984)

- Neotropical region: NW Brazil: Magela Creek area (Koste, 1981b); Brazil (Koste & José de Paggi, 1982); Rio Nhamunda, Amazonia, Bazil (Brandorff, Koste & Smirnov, 1982)

Testudinella epicopta Myers, 1934

- Nearctic region: Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988)

Testudinella gillardi De Ridder, 1966

- Palaearctic region: Pontic province (Bērziņš, 1978); W & SW Iceland (De Ridder, 1966a, 1967a, 1969b & 1972)

Testudinella greeni Koste, 1981

- Oriental region: In tectonic lake (Sone beel) in Cachar Valley of Assam State. (Sharma, 1990); Malaysia & Indonesia (Sudzuki, 1989)

- Australian region: NW Aust.: Magela Creek area (Koste, 1981b); NT, Australia (Shiel & Koste, 1985); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984)

- Neotropical region: Argentina (Koste & José de Paggi, 1982); River Parana, Argentina (José de Paggi, 1985); Parana medio river environments, Argentina (José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983); Province Santa Fé, Argentina (José de Paggi, 1990)

Testudinella hysseyi Shiel & Koste, 1985

- Australian region: N Queensland, Aust. (Shiel & Koste, 1985); Tasmania (Koste & Shiel, 1986b & 1987)

Testudinella incisa (Ternetz, 1892)

- Palaearctic region: 3 lakes of Sosnowickie, Poland (Radwan, 1966 & 1969); Lakes in the Leczna-Włodawa area & Silesia, Pol. (Radwan, 1967); Parczew area peat bogs, Poland (Radwan, 1974); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Fens in Genk, Belgium (Schepens, 1960); Pod padurka pond & Bajinice II pond, Poland (Lewkowicz, 1972); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Anatolia, Turkey (Segers *et al.*, 1992); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Fresh & brackish waters in the Netherlands (Dresscher, 1976); Romania (Rudescu, 1960); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); All over Europe; Caspian area (Bērziņš, 1978); Lake Balaton, Hungary (Zankai, 1968); N Iceland (De Ridder, 1972); Karelian inland waters (Kutikova, 1965)

- Nearctic region: N America (Bērziņš, 1978); Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); NW Canada (Chengalath & Koste, 1987); Wisconsin, USA (Bērziņš, 1984); Northern Michigan inland lakes (Stemberger, 1990)

- Oriental region: India (Sharma & Michael, 1980); Sri Lanka (Fernando, 1980a); Sri Lanka & India (Sudzuki, 1989); Sri Lanka: Gem Pit Ratnapura (Chengalath, Fernando & Koste, 1974)

- Australian region: L. Euramo, tropical Queensland crater lake, Aust. (Green, 1981); Qld, Aust. (Shiel & Koste, 1979); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); S Tasmania & Qld, Aust. (Koste, Shiel & Tan, 1988); Notornis Valley, New Zealand (Russell, 1960)

- Neotropical region: Panama (Koste & José de Paggi, 1982; Collado *et al.*, 1984); Near Panguana, Peru (Koste, 1988b); Pond in tropical rain forest, Peru (Schlüter, 1984)

Testudinella mucronata mucronata (Gosse, 1886)

- Palaearctic region: Middle course of river Loire, France (Lair & Sargas, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); River Sambre in Landrecies, France (Waeyaert, 1972); Ponds in the Paris suburb, France (Pourriot, 1965); Serbian section of river Danube (Zivkovic, 1987); Spain: Madrid & Valencia (Velasco, 1990); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Romania (Rudescu, 1960); River Klyas'ma, R.F. (Bogoslawski, 1961); Heiluntsian province, China (Bogoslawski, 1962); Lake Balaton, Hungary (Zankai & Kertesz, 1967; Zankai, 1968; Zankai & Ponzi, 1971a); River Loire, France (Lair, 1980); Hortobagy Nat. Park, Hungary (Forro & Ronkay, 1983); N Swedish Lapland

(Pejler, 1962a); Darnsee near Epe, NW Germany (Koste, 1962); River Hase, NW Germany (Koste, 1976b); Lake Alfsee, NW Germany (Koste & Poltz, 1987); Saratov reservoir, R.F. (Popchenko *et al.*, 1981); All over Europe (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1968 & 1970); Reservoir Sedlice, near Zeliv, Bohemia, CSR (Sladeckova, 1960); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Lake Donk, Overmere, Belgium (Dumont, 1988b); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Eeklo, Belgium (De Maeseneer, 1979; De Ridder, 1986); River Danube (Enaceanu & Brezeanu, 1970); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Anatolia, Turkey (Segers *et al.*, 1992); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Ponds in Oltina, SE Dobrogea, Romania (Popescu-Gorj & Costea, 1961); River Loire in St. Laurent-les-Eaux, France (Pourriot *et al.*, 1982); Lakes in the Leczna-Włodawa area & in several other parts of Poland (Radwan, 1967); River Rijn, Waal & Yssel; Pond in Enschede; Mastenbroeker kolk; Petgaten in NW Overijssel; De Linde, Friesland; River Maas; Kil van Hurwenen, NL (Dresscher, 1976)

- Nearctic region: Arctic N America (Chengalath & Koste, 1989)
- Oriental region: India (Sharma & Michael, 1980; Sudzuki, 1989)
- Australian region: Swamp near Lake Hayes, S Island, New Zealand (Bērziņš, 1982a); Chatham Islands, New Zealand (Russell, 1960)
- Neotropical region: Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Surinam (Leentvaar, 1975); Ecuador: Oriente (Koste & Böttger, 1989 & 1992); Argentina (Modenutti & Claps, 1988); Lago Açu, Maranhão, Brazil (Reid & Turner, 1988); Amazonia, Brazil (Hauer, 1965; Thomasson, 1971; Koste, 1972f); Paraguay, Brazil, Argentina & Surinam (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Prov. Corriente, Argentina (José de Paggi, 1990)

Testudinella mucronata tasmaniensis Koste & Shiel, 1986

- Australian region: Stock dam Triabunna, Tasmania (Koste & Shiel, 1986b); Tasmania (Shiel, Koste & Tan, 1989)

Testudinella mucronata (Gosse), f. *hauerensis* Gillard, 1967

- Neotropical region: Orinoco river basin, Venezuela (Vasquez, 1984a); Lower Orinoco and its delta, Venezuela (Vasquez & Rey, 1989); Lago Açu, Maranhão, Brazil (Reid & Turner, 1988); Lago do Macaco, Brazil (Koste, 1989a); Amazonia, Brazil (Gillard, 1967; Thomasson, 1971; Koste, 1972f); Brazil & Venezuela (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); River Saladillo basin, Argentina (José de Paggi & Koste, 1988); Ecuador: Oriente & Costa (Koste & Böttger, 1989); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Concepcion, Paraguay (Koste, 1986b); Near Panguana, Peru (Koste, 1988b); Prov. Chaco & Santa Fé, Argentina (José de Paggi, 1990); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980); Lago Jurucui, Amazonia, Brazil (Schaden, 1977c); L. Redondo & L. Jacaretinga, Amazonia, Brazil (Serruya & Pollingher, 1984); Ecuador: Costa (Koste & Böttger, 1992)

Testudinella munda Bērziņš, 1982

- Australian region: Creek in Australia (Bērziņš, 1982); Creek in Vic, Aust. (Shiel & Koste, 1985)

Testudinella naumanni Carlin, 1939

- Palaearctic region: N Swedish Lapland (Pejler, 1962a); C Eur. plain (Bērziņš, 1978)
- Nearctic region: NE Canada (Chengalath & Koste, 1983); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989)

Testudinella neboisi Bērziņš, 1982

- Australian region: Creek in Australia (Bērziņš, 1982); Creek in Vic, Aust. "Must be considered a doubtful species" (Shiel & Koste, 1985)

Testudinella obscura Althaus, 1957

- Palaearctic region: East Balkan (Bērziņš, 1978); Roscoff, Atlantic coast of Brittany, France (D'Hondt, 1970); Shores of the White Sea & the Barents Sea, R.F. (Ghilarov, 1967); Black Sea (Rudescu, 1961); Spain: Tarragona (Velasco, 1990)

Testudinella ohlei Koste, 1972

- Neotropical region: Orinoco river basin, Venezuela (Vasquez, 1984a); Caroni river basin, Venezuela (Vasquez, 1984b); Lower Orinoco and its delta, Venezuela (Vasquez & Rey, 1989); Lago Paroni, Brazil (Koste, 1974a); Amazonia, Brazil (Koste, 1972f); Varzea near Santarem, Brazil (Koste, 1974b); Amazonia, Brazil; Varzea near Santarem; Lago Paroni on the lower Rio Tapajos; in acid waters near Rio Nhamunda, N of Paratins, on the Amazonas (Koste, 1978b); Brazil (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste, Robertson & Hardy, 1984); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Ecuador: Oriente (Koste & Böttger, 1992)

***Testudinella ohlei* Koste, f. *pustulata* Koste, 1978**

- Neotropical region: In acid waters near Rio Nhamunda, Amaz., Braz. (Koste, 1978b); Brazil (Koste & José de Paggi, 1982); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982)

***Testudinella panonica* Zivkovic, 1987**

- Palaearctic region: Serbian section of river Danube (Zivkovic, 1987)

***Testudinella parva* (Ternetz, 1892)**

- Palaearctic region: L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Dübener Heide, N Germany (Wulfert, 1961); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); 3 lakes of Sosnowickie, Poland (Radwan, 1966 & 1969); Parczew area peat bogs, Poland (Radwan, 1974); Lakes in the Leczna-Włodawa area & other regions in Poland (Radwan, 1967); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Fens in Genk, Belgium (Schepens, 1960); Serbian section of river Danube (Zivkovic, 1987); Spain: Pontevedra (Velasco, 1990); Pond Chalinski Maly II & Lipowy pond, Poland (Lewkowicz, 1972); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Romania (Rudescu, 1960); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Coast of S Finland (Eriksen, 1969); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989c, 1989d); N Swedish Lapland (Pejler, 1962a); Darnsee near Epe, NW Germany (Koste, 1962); All over Europe. Cosmopolitan (Bērziņš, 1978); Lake Balaton, Hungary (Zankai & Kertesz, 1967); Lake Donk, Overmere, Belgium (Dumont, 1968b); Japanese inland waters (Yamamoto, 1960a); Southwest islands of Japan (Suzuki *et al.*, 1992); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Eeklo, Belgium (De Maeseneer, 1979; De Ridder, 1986); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960; Dresscher, 1976)

- Nearctic region: Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Great Dismal Swamp, N Carolina-Virginia border, USA (Anderson, Benfield & Buikema, 1977); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980); Laurentian Great Lakes (Stemberger, 1979)

- Oriental region: Sri Lanka (Fernando, 1980a); Malaysia, Sri Lanka & Indonesia (Suzuki, 1989); New record for India: in paddy-fields in Jaintia hills district, Meghalaya State; paddy-fields in Kokrajhar & Nalbari districts, Assam State (Sharma, 1990); Kalimantan (Borneo), Indonesia (Koste, 1988a); Sri Lanka. Rare; found in ponds (Chengalath, Fernando & Koste, 1973); W & E Malaysia & Singapore (Fernando & Zankai, 1981); 2 Kumaun Himalaya lakes, UP, India (Sharma & Pant, 1984 & 1985)

- Australian region: Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); NW Aust.: Magela Creek area (Koste, 1981b); Beauffort Dam, Aust. (Bērziņš, 1982); Qld, Aust. (as *T. insinuata* Hauer, 1938) (Shiel & Koste, 1979); NT, Aust. (Koste & Shiel, 1980c)

- Neotropical region: Venezuela & Brazil (Koste & José de Paggi, 1982); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Lago Paroni, Brazil (Koste, 1974a); Amazonia, Brazil (Koste, 1972f); Várzea near Santarem, Brazil (Koste, 1974b)

***Testudinella parva* (Ternetz), f. *bidentata* (Ternetz, 1892)**

- Palaearctic region: Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Romania (Rudescu, 1960); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Darnsee near Epe, NW Germany (Koste, 1962); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); Alps, C Eur. mts., Carpathes, Pontic province, W, C & E Eur. plains, UK, Ireland, Boreal highland, NW Russian taiga (Bērziņš, 1978); Japanese inland waters (Yamamoto, 1960a); 3 lakes of Sosnowickie, Poland (Radwan, 1968 & 1969); Eeklo & Lacuisine, Belgium (De Maeseneer, 1979; De Ridder, 1986); Knowsley Park, Liverpool, UK (Galliford, 1964); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960; Dresscher, 1976); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Parczew area peat bogs, Poland (Radwan, 1974); Lakes in the Leczna-Włodawa area & in several other regions in Poland (Radwan, 1967); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Fens in Genk, Belgium (Schepens, 1960); Serbian section of river Danube (Zivkovic, 1987); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)

- Nearctic region: Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980)

- Oriental region: New to India; in East Khasi & Jaintia Hills districts of Meghalaya State (Sharma, 1990)

***Testudinella parva* (Ternetz), f. *semiparva* Hauer, 1938**

- Palaearctic region: 2 regions in Poland (Radwan, 1967); Serbian section of river Danube (Zivkovic, 1987); C Eur. plain (Bērziņš, 1978)

- Nearctic region: Canada (Chengalath, 1977)

- Oriental region: Lake Ranu Lamongau, Indonesia (Green *et al.*, 1976); Indonesia (Suzuki, 1989); New to India: in pond in Lakhimpur, Assam State. Up to now only recorded from Java and Poland (Sharma,

1990)

- Australian region: Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); NT, Aust. (Shiel & Koste, 1985)
- Neotropical region: Amazonia, Brazil (Thomasson, 1971; Koste, 1972f); Brazil (Koste & José de Paggi, 1982); ? Crescent Lake, Rio Sniu Missu, Mato Grosso, C Brazil (Green, 1972); Lago Paroni, Brazil (Koste, 1974a); Varzea near Santarem, Brazil (Koste, 1974b)

Testudinella patina (Hermann, 1783)

- Palaearctic region: Romania (Rudescu, 1960); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Po river, Italy (Ferrari *et al.*, 1989); Volga river & Volgograd reservoir, R.F. (V'yushkova & Lakhnova, 1974); Peridunar pond Dulce, Doñana Nat. Park, Spain (Lopez *et al.*, 1991); Fishfattening ponds in the Ukraina (Krazhan *et al.*, 1976); Vilnius, Lithuania (Rachynnas, 1971); Aral Sea spawning grounds, R.F. (Yerekeyev & Kazakhbayev, 1972); All over Europe; Caucasus, caspian area, Asia Minor (Bērziņš, 1978); "Grote vijver" fort 8, Hoboken (Antwerp, Belg.) (De Smet *et al.*, 1986); "Kleine vijver" fort 8, Hoboken (Antwerp, Belg.) (De Smet *et al.*, 1989); Moat of fort Merksem (Antwerp, Belgium) (De Smet *et al.*, 1987); Soviet section of river Danube, R.F. (Parchuk, 1985); Saratov reservoir, R.F. (Popchenko *et al.*, 1981); Minorca, Spain (de Manuel, 1990); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); River Hase, NW Germany (Koste, 1976b); Lake Dümmer, NW Germany (Koste & Poltz, 1983 & 1984); Lake Alfsee, NW Germany (Koste & Poltz, 1987); Delta of river Danube, Romania (Prunescu-Arion, 1989; Godeanu & Zinevici, 1983; Zinevici & Teodorescu, 1990); Fishponds near Warsaw, Poland (Hillbricht-Ilkowska, 1964); Lac Léman, Fr./CH (Martin, 1961; Balvay & Laurent, 1981 & 1989; Balvay *et al.*, 1991); Lac d' Aydat, Massif Central, France (Oulad Ali *et al.*, 1990); River Elbe near Hamburg, Germany (Caspers & Schulz, 1964); Freshwater pools in S Finland (Björklund, 1972a); Coast of S Finland (Eriksen, 1969); River Loire, France (Lair *et al.*, 1978; Lair, 1980); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Asiatic Turkey (Dumont & De Ridder, 1987); Croatia, ex-Yugoslavia (Erben, 1987); District Ostrava, CSR (Dvorakova, 1960b); Knowsley Park, Liverpool, UK (Galliford, 1964); Boatracing canal Gent, Belgium (De Maeseneer, 1972; Vanderbeken, 1977; Coussement, 1982); Lake Donk, Overmere, Belgium (Dumont, 1968b; Coussement, 1976); Pond in Botanical Garden, Gent, Belgium (Verheyen, 1980); Karagöl, near Izmir, Turkey (Ustaoglu, 1986); Akgöl (Selçuk-Izmir), Turkey (Ustaoglu & Balik, 1987); Lake Marmara, Turkey (Ustaoglu, in press); Camargue, S France (De Ridder, 1960a & 1961); Marismas, S Spain (De Ridder, 1962b); N & W Iceland (De Ridder, 1969b, 1970 & 1972); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Leczna-Włodawa lake district, E Poland (Radwan, 1984); Backwater Labicko, Middle Elbe, CSR (Straskraba, 1965); Korea (Turner, 1986); Japanese inland waters (Yamamoto, 1960a); Canal Isebek, Hamburg, Germany (Caspers & Schulz, 1960); Inundation area of Danube delta, Romania (Dorobantu, 1964); Lower Silesia, Poland (Pawlowski, 1960); River Grabia basin, Poland (Pawlowski, 1968, 1970 & 1973); Reservoir Sedlice, near Zeliv, Bohemia, CSR (Sladeckova, 1960); Lake Balaton, Hungary (Zankai & Kertesz, 1967; Zankai, 1968); Latorica, E Slowakia lowland, CSR (Terek, 1990a); Canals in Hran area, CSR (Terek, 1990c); Mountain Fuji, Japan (Suzuki, 1971); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Brackish waters in the Lorraine orientale, France (De Ridder, 1969a); All over Belgium (De Ridder, 1986); Lake Neusiedlersee, Austria (Donner, 1979); Clay pits in Stuivekenskerke, Belgium (Espeel, 1979); Eeklo, Belgium (De Maeseneer, 1979); Lake Constance, Germany (Walz *et al.*, 1987); River Danube (Enaceanu & Brezeanu, 1970); River Elbe estuary, Germany (Schulz, 1961); Peat bog Chambedaze, Puy-de-Dôme, France (Francez, 1982); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Canal Leeds-Liverpool, UK (Galliford, 1960b); Dozmary pool, Bodmin Moor, Cornwall, UK (Galliford, 1967); District Ardennen, Belgium (Gijsels *et al.*, 1976); Lakes N of Bucharest, Romania (Godeanu, 1969b); Lakes of the Black Sea shore, Romania (Godeanu, 1970b); Nat. Res. Wellmeersen, Belgium (Dumont *et al.*, 1975); River Vistula in the Warsaw area, Poland (Klimowicz, 1981); Near water works at Okocim, Poland (Klimowicz, 1975); Sewage ponds near Czestochowa, Poland (Klimowicz, 1968); Small water bodies in the Mikolajkie area, Poland (Klimowicz, 1967); Anatolia, Turkey (Segers *et al.*, 1992); River Bistritza & its tributaries, Romania (Miron, 1960); Lowland brook-pond system at Boom, Belgium (Maes, 1985); Bakkerskooi, Wanneperveen, NW Overijssel, NL (Leentvaar, 1967b); Shores of embanked or free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1976); Dübener Heide, N Germany (Wulfert, 1960a & 1960b); Brielse Maas & dune lakes on Voorne; all over the Netherlands (Dresscher, 1976); Impoundment Helme, near Kelbra, Kyffhäuser, Germany (Heynig, 1974); Etang de Soustons, SW France (De Ridder, 1967b); Bog pond near Budapest, Hungary (Nogradi, 1962); Black Sea (Rudescu, 1961); Ellesmere District, UK (Galliford, 1960a); Central Finland (Eloranta, 1981b); Lake Bolsena, C Italy (Ferrari, 1972); Hortobagy Nat. Park, Hungary (Forro & Ronkay, 1983); Nat. Res. Jursky Sur, CSR (Hudec, 1978); Near Asperg, Württemberg, Germany (Klement, 1960); "Altes Land" near

Hamburg, Germany (Caspers & Heckman, 1981 & 1982); Hamburg area, Germany (Heckman, 1982); Lake Bielersee, CH (Berner-Fankhauser, 1986); Middle course of river Po, Italy (Ferrari *et al.*, 1986); Lake Palaeostomi, Georgia, R.F. (Haberman & Laugaste, 1982); Afghanistan (Bērziņš, 1961); Monsa, Shetlands (De Smet *et al.*, 1988); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Alkal. Péteri-See, Hungary (Kertesz, 1960); Impoundment Kninicki, CSR (Kubicek, 1964); Lake Muy on Texel, NL (Leentvaar, 1967); L. Quackjeswater & L. Breedewater on Voorne; Lake Vogelmeer, Ymuiden, NL (Leentvaar, 1963); Broads & peat pits in NW Overijssel, NL (Schroevens, 1965); Karstic lake Percile, Latium, Italy (Margaritora, 1981); N Swedish Lapland (Pejler, 1962a); Baisse salée, Camargue, S France (Aguesse, 1960); Lago di Giulianello, Latium, Italy (Ferrara & Margaritora, 1977; Bazzanti & Stella, 1977); Lake Bodensee, near Konstanz, Germany (Althaus, 1962); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); River Klyas'ma, R.F. (Bogoslowski, 1961); Heiluntsian province, China (Bogoslowski, 1962); River Ybbarså, Sweden (Bērziņš, 1979); All over Europe (Bērziņš, 1978); Lake Monterosi, Latium, Italy (Stella & Margaritora, 1966); Karatas Lake, Turkey (Emir, 1991); Southwest islands of Japan (Sudzuki *et al.*, 1992); Waldviertel area, lower Austria (Naidenow & Wawrik, 1992); Mickletown Lagoons, lower Aire valley, UK (Ruttner-Kolisko, 1979); Ponds in the Paris suburb, France (Pourriot, 1965); Ponds in the Botanical Garden, Gent, Belgium (Segers *et al.*, 1991b); Garden ponds in Tokyo, Japan (Sudzuki, 1975); Lakes around mountain Fuji, Japan (Sudzuki, 1971); Mountain Fuji area, Japan (Sudzuki, 1978a); Periphytic biota in sewages & park waters, Jap. (Sudzuki, 1981); Pool, S of Trondheim, Norway (Thomasson, 1979); Scandinavian coastal waters (Thane-Fenchel, 1968); Lago Trasimeno, Italy (Taticchi, 1968); Serbian section of river Danube (Zivkovic, 1987); All over Spain. Baleares: Ibiza, Menorca, Mallorca (Velasco, 1990); Lake Biwa-ko, Japan (Yamamoto, 1960b); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Impoundment Twistetalsperrre, Nordhessen, G. (Meinel *et al.*, 1983); Danube delta, Romania (Popescu-Marinescu & Zinevici, 1967); 4 lakes in the Puy-de-Dôme, France (Persoone *et al.*, 1968); Luga district in the Leningrad area, R.F. (Kutikova, 1962b); In reed borders of the Danube delta, Romania (Rudescu, 1968); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); Ponds in Oltina, SE Dobrogea, Romania (Popescu-Gorj & Costea, 1961); Rés. biol. de Villars-la-Dombes, France (Rostan, 1976); River Loire, in St. Laurent-les-Eaux, France (Pourriot *et al.*, 1982); Lakes in the Leczna-Włodawa area & in many other regions in Poland (Radwan, 1967); 3 lakes of Sosnowickie, Poland (Radwan, 1969); Parczew area peat bogs, Poland (Radwan, 1974); Lake Jorzec, Mazurian lakeland, Poland (Radwan *et al.*, 1985); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Middle course of river Loire, France (Lair & Sargas, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979)

- Nearctic region: Canada (Chengalath, 1977); Ontario, Canada (Chengalath, Fernando & George, 1971; Chengalath & Fernando, 1973); NE Canada (Chengalath & Koste, 1983); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Gregg's Lake, Fentrus L., Highland L., Little Swede n° 1 L., Parvin L. & Hayden's L., Colorado, USA (Pennak, 1966); Wisconsin, Illinois & Kentucky, USA (Bērziņš, 1984); Ross Lake, Florida (Bienert, 1986); Illinois river, USA (Brown *et al.*, 1989); Lake O'Woods, West Virginia, USA (Havens & Da Costa, 1987); Colorado, USA (Turner, 1988); Kingston, Ontario, Canada (Nogrady, 1989); Northern Michigan inland lakes (Stemberger, 1990); Atchafalaya river basin, Louisiana, USA (Holland *et al.*, 1983); Lakes in Santee, California, USA (Nogrady, 1982); Lac Echo, Québec, Canada (Nogrady, 1976); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980); Desaulniers reservoir, Québec, Canada (Pinel-Alloul *et al.*, 1982); Greens Lake, Beaver Island, Michigan, USA (Foran & King, 1982); Near Monroe, Louisiana, USA (Gallagher, 1966); Crater Lake & Walker L., Arizona, USA (Green, 1986); Experimental ponds near Bakersfield, Kern County, California, USA

- Oriental region: 2 sewage channels of Gwalior, India (Saksena & Kulkarni, 1986); Taiwan (Sudzuki, 1988); India (Sharma & Michael, 1980); Kashmir, Gujarat, Rajasthan, Andhra Pradesh, W Bengal, Orissa, Assam & Meghalaya + different states of NE India (Sharma, 1990); Kalimantan (Borneo), Indonesia (Koste, 1988a); Philippines (Mamaril & Fernando, 1978); Sri Lanka (Chengalath, Fernando & Koste, 1973; Fernando, 1980a); Singapore & Malaysia (Karunakaran & Johnson, 1978); Fishponds in Thailand (Boonsom, 1984a); Thailand (Boonsom, 1984b); W & E Malaysia & Singapore (Fernando & Zankai, 1981); Maini Lake, Stocking tank & Nursery tank, Delhi, India (George, 1966); Fishponds at Barrackpore, W Bengal State, India (Michael, 1968); Punjab State, India (Sharma & Sharma, 1984); Impoundment Kinda, C Burma (Koste & Tobias, 1990, 1982); 2 Kumaun Himalaya lakes, UP, India (Sharma & Pant, 1984 & 1985); Lake Ranu Lamongau, Indonesia (Green *et al.*, 1976); Parakrama Samudra, Sri Lanka (Fernando & Rajapaksa, 1983); Ricefields in NE Thailand (Heckman, 1979); West Bengal, NE India (Sharma, 1979b); Orissa, E India (Sharma, 1980b); Sri Lanka (Mendis & Fernando, 1962; Fernando, 1969); South India (Naidu, 1967); Punjab State, NW India (Sharma, 1976); Storage reservoir Ajwa, Prata-pura reservoir & Nimeta water works, near Baroda, India (Wulfert, 1966); Lake Khan Payao (Serruya & Pollingher, 1984)

- Australian region: In many localities in Australia (Bērziņš); SW Aust. (Koste, Shiel & Brock, 1983); Lake

Purrumbete, freshwater Aust. crater lake (Green, 1981); NW Aust.: Magela Creek area (Koste, 1981b); Tasmania (Koste & Shiel, 1986b); Alligator rivers area, NT, Aust. (Koste & Shiel, 1983a; Tait, Shiel & Koste, 1984); Billabongs in the river Murray floodplain, NSW, Aust. (Hillman & Shiel, 1991); NSW, Qld & Vic, Aust. (Shiel & Koste, 1979); Lower river Murray, S Aust. (Shiel, Walker & Williams, 1982); Saddle Hill pond, New Zealand (Bryas, 1960); Waihola & Canterbury, New Zealand (Russell, 1960)

- Neotropical region: Antilles françaises (Pourriot, 1975); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Suape area, Pernambuco, Brazil (Neumann-Leitao *et al.*, 1992); Laguna Cocha Cashu, Amazonia, Brazil (Schaden, 1977c); Ecuador: Oriente & Costa (Koste & Böttger, 1989 & 1992); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980); Chilean lakes (Schmid-Araya, 1991); River Santa Fé, Argentina (José de Paggi, 1976b); River Saladillo basin, Argentina (José de Paggi & Koste, 1988); River Parana, Argentina (Paggi & José de Paggi, 1974; José de Paggi, 1978a & 1985); Argentina (Modenutti & Claps, 1988); Orinoco river basin, Venezuela (Vasquez, 1984a); Lower Orinoco river & its delta, Venezuela (Vasquez & Rey, 1989); Lago do Macaco, Brazil (Koste, 1989a); Provinces in NW Argentina (José de Paggi, 1989); Lago Paroni, Brazil (Koste, 1974a); Amazonia, Brazil (Gillard, 1967; Thomasson, 1971; Koste, 1972f); Varzea near Santarem, Brazil (Koste, 1974b); Argentina, Venezuela, Colombia, Brazil, Chile, Paraguay, Antilles, Peru, Surinam, Panama (Koste & José de Paggi, 1982); Rio Uatumá, Amazonia, Brazil (Koste & Hardy, 1984); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Concepcion, Paraguay (Koste, 1986b); Near Panguana, Peru (Koste, 1988b); Galapagos Islands (De Smet, 1987 & 1989); Island Maraca, Roraima, Brazil (Koste & Robertson, 1990); Surinam (Leentvaar, 1975); Argentine Patagonia (Kuczynski, 1987b); Amazonia, Brazil (Hauer, 1965b); Lake Escondido, Rio Negro prov., Argentina (Baleiro & Modenutti, 1990); Prov. Chubut & Rio Negro, Argentina (José de Paggi, 1990); Caribbean part of the Neotropical region (De Ridder, 1977); Aruba, Bonaire, Curaçao, Bahamas, Caribbean Islands, Cuba & Haiti (Collado *et al.*, 1984); Lago do Valencia, Venezuela (De Infante, 1980); Brokopondo Lake, Surinam (Leentvaar, 1979); Santa Cruz & Tierra del Fuego, Argentina (Olivier, 1965)

Testudinella patina (Hermann), f. *aspis* Carlin, 1939

- Palaearctic region: C Eur. mountains & plains, Baltic province (Bērziņš, 1978); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973)

Testudinella patina (Hermann), f. *dendradena* de Beauchamp, 1955

- Palaearctic region: Southwest islands of Japan (Suzuki, 1990; Suzuki *et al.*, 1992)
- Oriental region: Kalimantan (Borneo), Indonesia (Koste, 1988a); Indonesia (Suzuki, 1989a); Taiwan, Malaysia, Singapore, Indonesia, Sri Lanka, Philippines & India (Suzuki, 1989b)
- Neotropical region: Orinoco river basin, Venezuela (Vasquez, 1984a); Lower Orinoco & its delta, Venezuela (Vasquez & Rey, 1989); Venezuela (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil. First record (Koste & Robertson, 1983); for the Amazon region (Koste, Robertson & Hardy, 1984); Near Concepcion, Paraguay (Koste, 1986b); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980); Mato Grosso, C Brazil (Turner & Da Silva, 1992)

Testudinella patina (Hermann), f. *intermedia* (Anderson, 1889)

- Palaearctic region: Lowland brook-pond system in Boom, Belgium (Maes, 1985); Heiluntsian province, China (Bogoslawski, 1962); C Eur. mts., Hungarian plain, C & E Eur. plains, Baltic province, UK, Caucasus, Caspic area (Bērziņš, 1978); Alkal. Péteri-See, Hungary (Kertesz, 1960); Impoundment Kninicki, CSR (Kubicek, 1964); River Svatka & new impoundments in the CSR (Kubicek, 1968); Lakes of Sosnowickie, Poland (Radwan, 1968 & 1969); Parczew area peat bogs, Poland (Radwan, 1974); Lakes in the Leczna-Włodawa area & "Plaine de la Grande Pologne", Poland (Radwan, 1967); Serbian section of river Danube (Zivkovic, 1987); Spain: Madrid (Velasco, 1990); Southwest islands of Japan (Suzuki *et al.*, 1992); Boom, Belgium (De Ridder, 1986); Side-arm of river Danube near Bratislava, CSR (Lichardova, 1964)

- Nearctic region: Kingston, Ontario, Canada (Nogrady, 1989)

- Oriental region: Calcutta (Anderson, 1889). Paddy-field in Tihu (Sharma, 1990); (Malbari district, Assam State, India)

- Australian region: Lake Barracoota, Aust. (Bērziņš, 1982); Vic, Aust. (Shiel & Koste, 1979 & 1985)

- Neotropical region: Near Panguana, Peru (Koste, 1988b)

Testudinella patina (Hermann), f. *lindbergi* Bērziņš, 1961

- Palaearctic region: Afghanistan (Bērziņš, 1961)

Testudinella patina (Hermann), f. *trilobata* (Anderson & Shephard, 1892)

- Palaearctic region: Lakes in the Leczna-Włodawa area & Haute Silésie, Poland (Radwan, 1967); Lakes of Sosnowickie, Poland (Radwan, 1966 & 1969); Serbian section of river Danube (Zivkovic, 1987); Spain: Madrid (Velasco, 1990); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Heiluntsian province, China (Bogoslawski, 1962); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); District

Ostrava, CSR (Dvorakova, 1960b); Alkal. Péteri-See, Hungary (Kertesz, 1960); Lake Balaton, Hungary (Zankai & Kertesz, 1967); Darnsee near Epe, NW Germany (Koste, 1962); C Eur. mts., Carpathes, Hungarian plain, C & E Eur. plains, Caspian area (Bērziņš, 1978); Obretin lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Southwest islands of Japan (Sudzuki *et al.*, 1992); Romania (Godeanu, 1961); Lakes, N of Bucarest, Romania (Godeanu, 1969b)

- Nearctic region: Kentucky, USA (Bērziņš, 1984); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990)

- Oriental region: W & E Malaysia & Singapore (Fernando & Zankai, 1981); Malaysia (Sudzuki, 1989); Singapore & Taiwan (Sudzuki, 1991)

- Australian region: Vic, Aust. (Shiel & Koste, 1979)

- Neotropical region: Surinam (partim *sub nomine T. lobosa*) (Leentvaar, 1975); Brokopondo Lake, Surinam (Leentvaar, 1979); Amazonia, Brazil (Koste, 1972f); Brazil, Argentina, Surinam, Panama (Koste & José de Paggi, 1982); Orinoco river basin, Venezuela (Vasquez, 1984a); Chilean lakes (Schmid-Araya, 1991); Prov. Santa Fé, Argentina (José de Paggi, 1990); Panama (Collado *et al.*, 1984); Laguna Costa Cashu, Amazonia, Brazil (Schaden, 1977c)

***Testudinella pseudoclypeata* Bērziņš, 1943**

- Palaearctic region: W Eur. plain & Baltic province (Bērziņš, 1978)

***Testudinella reflexa* (Gosse, 1887)**

- Palaearctic region: Peats in the Caucasus (Tarnogradsky, 1961); Nat. Res. "Leiemeersen", Assebroek, Belgium (De Ridder, 1992); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Dübener Heide, N Germany (Wulfert, 1961); Romania (Rudescu, 1960); Prov. Zuid-Limburg, NL (Dresscher, 1976); Coast of S Finland (Eriksen, 1969); C & S Sweden (Bērziņš & Pejler, 1987); Hortobagy Nat. Park, Hungary (Forro & Ronkay, 1983); Near Asperg, Württemberg, Germany (Klement, 1960); N Swedish Lapland (Pejler, 1962); Darnsee near Epe, NW Germany (Koste, 1962); Alps, C Eur. mts., Carpathes, Hungarian plain, C & E Eur. plains, Baltic prov., Fennoscandia, Caspian area. Cosmopolitan (Bērziņš, 1978); Lake Balaton, Hungary (Zankai, 1968); Small water bodies in the Mikolajkie area, Poland (Klimowicz, 1967); Anatolia, Turkey (Segers *et al.*, 1992); Several regions in Poland (Radwan, 1967)

- Nearctic region: Canada (Chengalath, 1967); Laurentian Great Lakes (Stemberger, 1979)

- Australian region: Creek in Aust. (Bērziņš); NSW & Qld, Aust. (Shiel & Koste, 1979); Mt. Cook, New Zealand (Russell, 1960)

- Neotropical region: Parana medio river environments, Argentina (José de Paggi, 1982); Brokopondo Lake, Surinam (Leentvaar, 1979); Surinam & Argentina (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste, Robertson & Hardy, 1984); Province Santa Fé, Argentina (José de Paggi, 1990)

***Testudinella robertsoni* Koste & Robertson, 1990**

- Neotropical region: Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990)

***Testudinella sculpturata* Bartos, 1951**

- Palaearctic region: C Eur. mountains (Bērziņš, 1978); Southwest islands of Japan (Sudzuki, 1990; Sudzuki *et al.*, 1992)

***Testudinella sphagnicola* Rudescu, 1960**

- Palaearctic region: In peat bogs of S Bukowina, Romania (Rodewald-Rudescu & Godeanu, 1961); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Carpathes (Bērziņš, 1978)

***Testudinella striata* (Murray, 1913)**

- Australian region: Müller Glacier, New Zealand (Russell, 1960)

***Testudinella triangularis* Myers, 1934**

- Nearctic region: Lac Echo, Québec, Canada (Nogrady, 1976); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980)

***Testudinella tridentata* Smirnov, 1931**

- Palaearctic region: Heiluntsian province, China (Bogoslawski, 1962); Amur region, Ussuri & China (Koste, 1981b)

- Nearctic region: Mt. Temblant Prov. park, Québec, Canada (Nogrady, 1980)

- Oriental region: New record for India: in 2 ponds in Somithur district of Assam State and in a paddy-field in Kailashaker (N Trip.) (Sharma, 1990); Kalimantan (Borneo), Indonesia (Koste, 1988a)

- Australian region: Jubiluka billabong, NT, Aust. (Koste, 1981b); Alligator rivers area, NT, Aust. (Koste & Shiel, 1983; Tait, Shiel & Koste, 1984); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Concepcion, Paraguay (Koste, 1986b); E Aust. (Koste & Shiel, 1980b); Vic, Aust. (Koste & Shiel, 1980c)

- Neotropical region: Brazil, Argentina & Surinam (Koste & José de Paggi, 1982); Surinam (Leentvaar, 1975); River Parana, Argentina (José de Paggi, 1985 & 1987b); Ecuador: Oriente (Koste & Böttger, 1989 & 1992); Lago Paroni, Brazil (Koste, 1974a); Province Santa Fé, Argentina (José de Paggi, 1990)

Testudinella tridentata Smirnov, f. *amazonica* Thomasson, 1971

- Neotropical region: Orinoco river basin, Venezuela (Vasquez, 1984a); Caroni river basin, Venezuela (Vasquez, 1984b); Lower Orinoco and its delta, Venezuela (Vasquez & Rey, 1989); Varzea near Santarem, Brazil (Koste, 1974b); Lago do Macaco, Brazil (Koste, 1989a); Amazonia, Brazil (Thomasson, 1971; Koste, 1981b); Lago Paroni, Amazonia, Brazil (Koste, 1974a); Brazil & Venezuela (Koste & José de Paggi, 1982); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980)

Testudinella tridentata Smirnov, f. *braziliensis* Koste, 1972

- Australian region: NW Aust.: Magela Creek area (Koste, 1981b)
- Neotropical region: Amazonia, Brazil; Brazil; also R.F., USA & Africa (Koste, 1972f)

Testudinella tridentata Smirnov, f. *curvata* Wulfert, 1965

- Oriental region: W & E Malaysia & Singapore (Fernando & Zankai, 1981)

Testudinella tridentata Smirnov, f. *dicella* Myers, 1934

- Nearctic region: Canada (Chengalath, 1977); Wisconsin & New Jersey, N America (Koste, 1981b); Ross Lake, Florida, USA (Bienert, 1986)

Testudinella tridentata Smirnov, f. *insulana* Bērziņš, 1982

- Oriental region: Danau Teleko, near Palembang, Sumatra (Bērziņš, 1982a)

Testudinella trilobata haterunensis Sudzuki, 1992

- Palaearctic region: Southwest islands of Japan (Sudzuki *et al.*, 1992)

Testudinella trilobata triangularis Sudzuki, 1992

- Palaearctic region: Southwest islands of Japan (Sudzuki *et al.*, 1992)

Testudinella trilobata amphora Sudzuki, 1992

- Palaearctic region: Southwest islands of Japan (Sudzuki *et al.*, 1992)

Testudinella truncata (Gosse, 1886)

- Palaearctic region: River Grabia basin, Poland (Pawlowski, 1960, 1968 & 1970); Lake Balaton, Hungary (Zankai, 1968); Latorica, E Slovakia lowland, CSR (Terek, 1990a); Canals in Hran area, CSR (Terek, 1990c); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Shores of embanked or free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); Domein De Linde, Friesland, NL (Dresscher, 1976); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); 3 lakes of Sosnowickie, Poland (Radwan, 1969); Lakes in the Leczna-Wlodawa area & several other regions in Poland (Radwan, 1967); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Romania (Rudescu, 1960); Lake Stechlinsee, Brandenburg, Romania (Althaus, 1963; Ronneberger & Schönborn, 1985); Coast of S Finland (Eriksen, 1969); Pools of the Upper Elbe, CSR (Dvorakova, 1961); District Ostrava, CSR (Dvorakova, 1960b); Lake Nehmitzsee, Brandenburg, Germany (Kasprzak & Ronneberger, 1982); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Nat. Res. Neeruti lakes, NE Estonia (Mäemets A. & A., s.d.); Darnsee near Epe, NW Germany (Koste, 1962); River Hase, NW Germany (Koste, 1976b); C Eur. mountains, Carpathes, Pontic prov., C & E Eur. plains, Baltic province, Ireland, UK, NW Russian taiga, Caspian area (Bērziņš, 1978)

- Nearctic region: Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988)

Testudinella truncata (Gosse), f. *ecornis* Wiszniewski, 1953

- Palaearctic region: C Eur. mountains & plains (Bērziņš, 1978); Pelagic zone of the Leczna-Wlodawa lakes, Pol. (Radwan, 1975); Serbian section of river Danube (Zivkovic, 1987)

Testudinella truncata (Gosse), f. *usmaensis* Bērziņš, 1943

- Palaearctic region: Baltic province (Bērziņš, 1978)

Testudinella unicornuta Koste & Shiel, 1987

- Australian region: Tasmania (Koste & Shiel, 1986b & 1987; Shiel, Koste & Tan, 1989)

Testudinella vidzemensis Bērziņš, 1943

- Palaearctic region: River Ybbarspså, Sweden (Bērziņš, 1979); C & S Sweden (Bērziņš & Pejler, 1987, 1989a & 1989d); C Eur. plain & Baltic province (Bērziņš, 1978)
- Australian region: Creek in Australia (Bērziņš, 1982); Vic, Aust. (Shiel & Koste, 1985)

Testudinella walkeri Koste & Shiel, 1980

- Australian region: NT, Australia (Koste & Shiel, 1980c); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984)

Testudinella spec.

- Palaearctic region: Lake Oerd, Ameland, NL (Leentvaar, 1967); Norfolk Broads, UK (Irvine *et al.*, 1990); Delta of river Danube (Prunescu-Arion, 1989); Lake Balaton, Hungary (Zankai, 1968); Lowland brook-pond system in Boom, Belgium (Maes, 1985); Bakkerskooi, NW Overijssel; Loenerveense plas Wijde Blik; Petgaten in NW Overijssel; Lake Het Oerd, Ameland; Lake Veerse Meer, NL (Dresscher, 1976); Etang de

- Soustons, SW France (De Ridder, 1967b); Lac Bourdouze, Puy-de-Dôme, France (Persoone *et al.*, 1968); Bulgarian bank of river Danube (Naidenow, 1963)
- Nearctic region: Northern Michigan inland lakes (Stemberger, 1990); NW Colorado, USA (Cauton & Ward, 1981); Lake Wabamun, Alberta, Canada (Gallup & Hickman, 1975); Mirror Lake, New Hampshire, USA (Strayer, 1985)
 - Oriental region: Malaysia & Singapore (Sudzuki, 1989); Thailand (Boonsom, 1984b); Rothanara tank, Indranagar tank, Stocking tank & Nursery tank, Delhi, India (George, 1966); River Yamuna, India (Rai, 1974)
 - Australian region: Lake Catani, Mt. Buffalo, Australia (Bērziņš, 1982)
 - Neotropical region: S Brazil (Arcifa, 1984); Crescent Lake, Rio Snia Missu, Mato Grosso, C Brazil (Green, 1972); Near Panguana, Peru (Koste, 1988a); Lake Monte Alegre, SE Brazil (Arcifa *et al.*, 1992)

Tetrasiphon Ehrenberg, 1840

Tetrasiphon hydrocora Ehrenberg, 1840

- Palaearctic region: Coast of S Finland (Eriksen, 1969); Germany, France, UK, Sweden (Koste, 1968d); In peat bogs in the "Massif Central", France (Batut, 1965); Small water bodies in Soviet Karelia, R.F. (Filimonova, 1976); C Eur. mts., W & C Eur. plains, UK, NW Russian taiga (Bērziņš, 1978); Lakes of Sosnowickie, Poland (Radwan, 1968 & 1969); Pond near Besse-en-Chandesse, France (Omaly & Dévaux, 1971); Peat bogs in the Paris suburb, France (Pourriot, 1965)
- Nearctic region: USA (Koste, 1968d); N America (Bērziņš, 1978); Crater Lake, Arizona, USA (Green, 1986); Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Wisconsin & Kentucky, USA (Bērziņš, 1984); Mt. Tremblant Prov. park, Québec, Canada (Nograday, 1980); Mirror Lake, New Hampshire, USA (Strayer, 1985)
- Oriental region: Sri Lanka (Fernando, 1980a); Malaysia & Sri Lanka (Sudzuki, 1989); Sri Lanka: Gem pit, Ratnapura (Chengalath, Fernando & Koste, 1974)
- Australian region: Tasmania (Koste & Shiel, 1987); NSW, Aust. (Shiel & Koste, 1979); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Billabongs of upper Murray tributaries, NSW; Vic; dune lakes in Tasmania NSW Koste & Shiel, 1991)
- Neotropical region: Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Ecuador: Oriente & Sierra (Koste & Böttger, 1992)

Tetrasiphon spec.

- Nearctic region: The Sacramento-San Joaquin delta estuary, USA (Siegfried & Kopache, 1980)
- Oriental region: Ponds in Chinglepat, Madras, India

Trichocerca Lamarck, 1801

Trichocerca agnatha Wulfert, 1939

- Palaearctic region: C Eur. plain (Bērziņš, 1978); Southwest islands of Japan (Sudzuki *et al.*, 1992); Sinaia, S Carpathes, Romania (Rodewald-Rudescu & Godeanu, 1961); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Impoundment Twistetalsperre, Nordhessen, G. (Meinel *et al.*, 1983)
- Australian region: Vic, Aust. (Koste & Shiel, 1980c); Single record from the river Murray at Echuca, Vic. (Koste & Shiel, 1992)

Trichocerca arctica Voigt, 1957

- Palaearctic region: Pools with fresh and brackish water, S Finland (Björklund, 1972a); NW Russian tundra (Bērziņš, 1978)

Note : See also *Trichocerca minuta* (Olofsson)

Trichocerca barsica (Varga & Dudich, 1939)

- Palaearctic region: C Eur. mountains & Hungarian plain (Bērziņš, 1978)

Trichocerca bicristata (Gosse, 1887)

- Palaearctic region: Romania (Rudescu, 1960); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Afghanistan (Bērziņš, 1961); Lakes in Estonia (Haberman, 1978); Coast of S Finland (Eriksen, 1969); Delta of river Danube (Godeanu & Zinevici, 1983); Sierra Nevada, Spain (Morales-Baquero, 1987); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Lake Muy (Texel) & Lake Oerd (Ameland), NL (Leentvaar, 1967); Lac Léman, Fr./CH (Balvay & Laurent, 1990); (Balvay *et al.*, 1985); Fish-fattening ponds in the Ukraina (Krazhan *et al.*, 1976); All over Europe (Bērziņš, 1978); Fish ponds near Warsaw,

Poland (Hillbricht-Ilkowska, 1964); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); Alpine lakes, Sierra Nevada, Spain (Morales-Baquero *et al.*, 1989); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Japanese inland waters (Yamamoto, 1960a); Kalken, Belgium (De Ridder, 1986); River Grabia basin, Poland (Pawlowski, 1960, 1968 & 1970); Lake Balaton, Hungary (Zankai, 1968); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Water-courses in the Bucegi mts., Romania (Godeanu, 1963b); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); High mt. lakes & ponds of Sierra Nevada, Spain (Morales-Baquero *et al.*, 1988); Temporary waters in the Bucegi mts., Romania (Godeanu *et al.*, 1960); Small water bodies in the Mikolajki area, Poland (Klimowicz, 1967); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Prov. of Drente; Lake Muy on Texel; Lake Oerd on Ameland, NL (Dresscher, 1976); Luga district of Leningrad area, R.F. (Kutikova, 1962b); Karelian inland waters, R.F. (Kutikova, 1965); Ponds with dense vegetation in Fontainebleau, Ile-de-France, France (Pourriot, 1970); Pool in Fontainebleau Forest, Ile-de-France, Fr. (Pourriot, 1976); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Rés. biol. de Villars-la-Dombes, France (Rostan, 1976); Parczew area peat bog, Poland (Radwan, 1974); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Fens in Genk, Belgium (Schepens, 1960); Lago Trasimeno, Italy (Taticchi, 1968); Serbian section of river Danube (Zivkovic, 1987); Spain : Gerona & Granada (Velasco, 1990); Lake Biwa-ko, Japan (Yamamoto, 1960b)

- Nearctic region: Pinehurst lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Lac Echo, Québec, Canada (Nogrady, 1976); Mt. Tremblant Prov. Park, Québec, Canada (Nogrady, 1980); Bay of Quinte, Lake Ontario, Canada (Nogrady, 1988); Stoneman Lake, Arizona, USA (Green, 1986); Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); 3 lakes in N Carolina, USA (Lemly & Dimmick, 1982); Wisconsin & Illinois, USA (Bērziņš, 1984); Ross Lake, Florida, USA (Bienert, 1986); Kingston, Ontario, Canada (Nogrady, 1989); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990)

- Oriental region: Sri Lanka (Fernando, 1980a); Malaysia & Sri Lanka (Sudzuki, 1989); Kalimantan (Borneo), Indonesia (Koste, 1988a); W & E Malaysia & Singapore (Fernando & Zankai, 1981); Fishponds in Thailand (Boonsom, 1984a); Thailand (Boonsom, 1984b); Sri Lanka. Fairly common in lakes & ponds (Chengalath, Fernando & Koste, 1973)

- Australian region: River in Aust. (Bērziņš, 1982); Qld, Aust. (Shiel & Koste, 1979); Lower river Murray, S Aust. (Shiel, Walker & Williams, 1982); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Rare in Murray-Darling basin, also Cape Cork, Qld and Kakadu Nat. Park, NT, Aust. (Shiel & Koste, 1992)

- Neotropical region: Prov. San Luis & Chaco, Argentina (José de Paggi, 1990); Panama (Collado *et al.*, 1984); Ecuador : Oriente (Koste & Böttger, 1989 & 1992); Galapagos Islands (De Smet, 1987 & 1989); Surinam (Leentvaar, 1975); River Saladillo basin, Argentina (José de Paggi & Koste, 1988); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Lower Orinoco and its delta, Venezuela (Vasquez & Rey, 1989); Lago Paroni, Brazil (Koste, 1974a); Amazonia, Brazil (cosmopolitan?) (Koste, 1972f); Varzea near Santarem, Brazil (Koste, 1974b); Brazil, Argentina, Venezuela & Panama (Koste & José de Paggi, 1982); Rio Yatapu & Rio Uatumá, Amazonia, Brazil (Koste & Hardy, 1984); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Rio Negro, Amazonia, Brazil (Schaden, 1977c); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Concepcion, Paraguay (Koste, 1986b); Near Panguana, Peru (Koste, 1988b); Amazonia, Brazil (Hauer, 1965b; Thomasson, 1971); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980); Araucarian ponds & pools, Chile (Thomasson, 1963b)

Trichocerca bicristata (Gosse), f. *mucosa* (Stokes, 1896)

- Palaearctic region: Southwest islands of Japan (Sudzuki *et al.*, 1992)

- Nearctic region: Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Canada (Chengalath, 1977); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990)

- Oriental region: Kalimantan (Borneo), Indonesia (Koste, 1988a); Indonesia (Sudzuki, 1989)

- Australian region: Qld, Australia (Shiel & Koste, 1974a)

- Neotropical region: Lago Paroni, Brazil (Koste, 1974a); Varzea near Santarem, Brazil (Koste, 1974b); Amazonia, Brazil (Hauer, 1965b; Koste, 1972f); Brazil, Panama, Surinam, Venezuela (Koste & José de Paggi, 1982); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980)

Trichocerca bicuspis (Pell, 1890)

- Nearctic region: Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Wisconsin, USA (Bērziņš, 1984); Kingston, Ontario, Canada (Nogrady, 1989); Laurentian

Great Lakes (Stemberger, 1979)

Trichocerca bidens (Lucks, 1912)

- Palaearctic region: River Vistula in the Warsaw area, Poland (Klimowicz, 1981); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Lake Stråken, Aneboda, Sweden (Bérziņš, 1972); Afghanistan (Bérziņš, 1961); Romania (Rudescu, 1960); Coast of S Finland (Eriksen, 1969); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); C & S Sweden (Bérziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Edgeøya, Svalbard (De Smet *et al.*, 1988b); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); All over Europe (Bérziņš, 1978); Lake Balaton, Hungary (Zankai & Kertesz, 1967; Zankai, 1968); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Peat bog Chambedaze, Puy-de-Dôme, France (Francez, 1982, 1987 & 1988); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Dübener Heide, N Germany (Wulfert, 1961); Lake Nobel, Ukraina (Polishchuk, 1991); Nat. Res. Het Hol, near Kortenhoef, NL (Dresscher, 1976); Karelian inland waters, R.F. (Kutikova, 1965); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Peats in the Caucasus (Tarnogradsky, 1961); Serbian section of river Danube (Zivkovic, 1987)
- Nearctic region: NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Mt. Tremblant Prov. Park, Québec, Canada (Nogrady, 1980)
- Oriental region: Taiwan (Suzuki, 1988); Taiwan, Malaysia & Indonesia (Suzuki, 1989)
- Australian region: Creek, river and lake in Aust. (Bérziņš, 1982); Swamp near Dunedin, S Island, New Zealand (Bérziņš, 1982a); Vic, Aust. (Shiel & Koste, 1985); S Tasmania & Vic, Aust. (Koste, Shiel & Tan, 1988); Cosmopolitan; isolated finds in acid waters, particularly *Sphagnum*. Rare. Tas & Vic (Shiel & Koste, 1992); Westland, New Zealand & Campbell Islands (Russell, 1960)
- Neotropical region: Galapagos Islands (De Smet, 1987 & 1989); Varzea near Santarem, Brazil (Koste, 1974b); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Brazil (Koste & José de Paggi, 1982); Ecuador : Oriente & Sierra (Koste & Böttger, 1992)
- Antarctic region: Antarctica & Subantarctica (José de Paggi & Koste, 1984)

Trichocerca birostris (Minkiewicz, 1900)

- Palaearctic region: Lake Zürichsee, CH (Nipkow, 1961); Lago di Giulianello, Latium, Italy (Ferrara & Margaritora, 1972 & 1977); Romania (Rudescu, 1960); Lakes in S Sweden (Arnemo *et al.*, 1968); Impoundment "Eisernes Tor", Danube, Romania (Brezeanu & Petén, 1973); Goggausee, Karinthia, Austria (Herzig & Moog, 1976); River Ybbarspå, Sweden (Bérziņš, 1979); Afghanistan (Bérziņš, 1961); Coast of S Finland (Eriksen, 1979); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); C & S Sweden (Bérziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Buchseen, Bodensee (Einsle, 1966); Mindelsee, Bodensee (Einsle, 1967); Lake Bolsena & L. Bracchiano, C Italy (Ferrari, 1972); Lake Biwa, Japan (Yamamoto, 1967 & 1968; Hillbright-Ilkowska, 1983a); C Sweden (Pejler, 1972); Lakes in C Sweden (Pejler, 1964 & 1965); Lago Maggiore, Italy (Tonolli, 1962); Lake Sempachersee, CH (Zimmermann, 1974); Lake Mälaren, Sweden (Grönberg, 1975); Fishponds near Warsaw, Poland (Hillbright-Ilkowska, 1964); Lake Erken, Sweden (Nauwerck, 1963); (Blomquist & Pejler, 1980); All over Europe. Cosmopolitan (Bérziņš, 1978); Lake Monterosi, Latium, Italy (Stella & Margaritora, 1966); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Lake Constance, Germany (Kiefer, 1968; Walz *et al.*, 1987); Lakes N of Bucarest, Romania (Godeanu, 1969b); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960); Lake Lauerzersee, CH (Odermatt, 1970); Lake Barlosche Kolk; Nat. Res. Oisterwijkse vennen; Nat. Res. Het Hol, near Kortenhoef, NL (Dresscher, 1976); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Lake Ösbysjön, near Stockholm, Sweden (Pejler, 1961); Lac Léman, Fr./CH (Rougier, 1969); Aegean lakes, Macedonia, ex-Yugoslavia & Greece (Serafimova-Hadzisce, 1974); Periphytic biota in sewages & park waters, Japan (Suzuki, 1981); A tarn in Värmland, Sweden (Thomasson, 1970); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)
- Nearctic region: Wisconsin, Illinois & Kentucky, USA (Bérziņš, 1984); Northern Michigan inland lakes (Stemberger, 1990); Lac Echo, Québec, Canada (Nogrady, 1976); Carter Lake, N Colorado, USA (Thomasson, 1964b)
- Oriental region: Girikbe-Tank, Sri Lanka (Bérziņš, 1982a)
- Australian region: Creek & 2 reservoirs in Aust. (Bérziņš, 1982); Farm dams near Gloucester, NSW, Aust. (Suzuki & Timms, 1980); Qld, Aust. (Shiel & Koste, 1979); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Lake Sorell, Tasmania (Suzuki, 1967)
- Neotropical region: Chile (Löffler, 1962); Chilean lakes (Schmid-Araya, 1991); Brokopondo Lake, Surinam (Leentvaar, 1979); Chile & Surinam (Koste & José de Paggi, 1982); Araucarian lakes, W & E of the Andes, Chile (Thomasson, 1963b); Amazonia, Brazil (Thomasson, 1971); Prov. Tierra del Fuego, Chubut, Rio Negro & Neuquen, Argentina (José de Paggi, 1990)

***Trichocerca brachyura* (Gosse, 1851)**

- Palaearctic region: Lakes, N of Bucharest, Romania (Godeanu, 1969b); Romania (Rudescu, 1960); River Loire, France (Lair, 1980); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); N Swedish Lapland (Pejler, 1962a); Darnsee near Epe, NW Germany (Koste, 1962); ? Lake Orog-Nur, Gobi desert, Mongolia (Koste & Wulfert, 1964); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Achmer Grasmoor, NW Germany (Koste, 1968b); ? La Mancha, Castill., Spain (Armengol *et al.*, 1975); Impoundment on Rio Guardinia, C Spain (Armengol, 1982); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Impoundment Saidenbach, Germany (Althaus, 1976); River Klyas'ma, R.F. (Bogoslawski, 1961); River Ybbarspān, Sweden (Bērziņš, 1979); Afghanistan (Bērziņš, 1961); Pools with fresh & brackish water, S Finland (Björklund, 1972a); Coast of S Finland (Eriksen, 1969); River Hase, NW Germany (Koste, 1976b); Lake Dümmer, NW Germany (Koste & Poltz, 1983); Lake Alfsee, NW Germany (Koste & Poltz, 1987); Fishponds near Warsaw, Poland (Hillbricht-Ilkowska, 1964); All over Europe. Cosmopolitan (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1970); Lake Balaton, Hungary (Zankai, 1968); Camargue, S France (De Ridder, 1960a & 1961); W Iceland (De Ridder, 1972); Japanese inland waters (Yamamoto, 1960a); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Prov. Drente; Nat. Res. Oisterwijkse vennen; Nat. Res. Het Hol, near Kortenhoef; Maria- and Helenapeel, NL (Dresscher, 1976); Brackish waters in the Lorraine orientale, France (De Ridder, 1969a); Samrée, Belgium (De Ridder, 1980 & 1986); Lake Constance, Germany (Walz *et al.*, 1987); Watercourses in the Bucegi mts., Romania (Godeanu, 1963b); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1960); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960); Temporary waters, small permanent ponds & marshy zones in the Bucegi mts., Rom. (Godeanu, 1969a); In periphyton of mosses & *Myriophyllum* in the Danube, and in some Alpine brooks, Austria (Donner, 1964); Shores of embanked or free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); River Salzach and its tributaries, Austria (Donner, 1970); Dübener Heide, N Germany (Wulfert, 1960b); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Bog pond near Budapest, Hungary (Nogrady, 1962); Karelian inland waters, R.F. (Kutikova, 1965); In reed borders of the Danube delta, Rom. (Rudescu, 1968); Peat bogs Poiana Stampei, Bukowina, Rom. (Rudescu, 1966); Middle course of river Loire, France (Lair & Sargas, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990); Lago Trasimeno, Italy (Taticchi, 1968); Serbian section of river Danube (Zivkovic, 1987); Spain : Ciudad real (?), Madrid, Orense, Pontevedra; Baleares : Menorca; Andorra (Velasco, 1990); Lake Biwa-ko, Japan (Yamamoto, 1960b); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)
 - Nearctic region: Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Illinois & Kentucky (Bērziņš, 1984); Kingston, Ontario, Canada (Nogrady, 1989); Miller's Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Lac Echo, Québec, Canada (Nogrady, 1976); Laurentian Great Lakes (Stemberger, 1979); Bayou de Siard, Monroe, Louisiana, USA (Gallagher, 1965)
 - Oriental region: India (Sharma & Michael, 1980); Malaysia, Indonesia & India (Sudzuki, 1989); Kalimantan (Borneo), Indonesia (Koste, 1988a); Lake Lano, Island of Mindanao, the Philippines (Lewis, 1977); Lakelet Tan Daha, Kathmandu valley, Nepal (Daems & Dumont, 1974)
 - Australian region: NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Creek in Aust. (Bērziņš, 1982); SW Aust. (Koste, Shiel & Brock, 1983); Tasmania (Koste & Shiel, 1987); NSW & Qld, Aust. (Shiel & Koste, 1979); Cosmopolitan, generally solitary in psammon and littoral of most freshwaters; pH-tolerant. Uncommon; probably pancontinental, but not yet recorded from S Aust. (Shiel & Koste, 1992); Lincoln & Christchurch, New Zealand; Chatham Islands (Russell, 1960)
 - Neotropical region: Provinces in NW Argentina (José de Paggi, 1989); Lago Paroni, Brazil (Koste, 1974a); Varzea near Santarem, Brazil (Koste, 1974b); Argentina, Brazil, Surinam, Panama, Costa Rica (Koste & José de Paggi, 1982); Near Concepcion, Paraguay (Koste, 1986b); Prov. Rio Negro, Tucuman & Santa Fé, Arg. (José de Paggi, 1990); Curaçao (De Ridder, 1977); Curaçao & Panama (Collado *et al.*, 1984); Brokopondo Lake, Surinam (Leentvaar, 1979); Prov. Neuquen & Tierra del Fuego, Argentina (Olivier, 1965); Araucarian Lakes, E of the Andes, Chile (Thomasson, 1963b); Lago do Castanho, C Amazonia, Brazil (Schaden, 1977b); Lago Rotondo, Amazonia, Brazil (Serruya & Pollingher, 1984); Ecuador: Costa (Koste & Böttger, 1992)
 - Antarctic region: Antarctica (Dartnall & Hollowday, 1985)
- Trichocerca brasiliensis* (Murray, 1913)**
- Oriental region: India (Sharma & Michael, 1980); Sri Lanka (Fernando, 1980a); Indonesia, Sri Lanka & India (Sudzuki, 1989); Kalimantan (Borneo), Indonesia (Koste, 1988a); Sri Lanka. Rare, found only from one pond, near Tobbawa (Chengalath, Fernando & Koste, 1983); W Bengal, NE India (Sharma, 1979b)

- Australian region: S Tasmania (Koste, Shiel & Tan, 1988); Tasmania (Shiel, Koste & Tan, 1989); Single record, Tasmania, from a roadside marsh, Deloraine (Shiel & Koste, 1992)
- Neotropical region: Ecuador : Oriente, Sierra & Costa (Koste & Böttger, 1989); Ecuador : Oriente (Koste & Böttger, 1992); Lago do Macaco, Brazil (Koste, 1989a); Amazonia, NE Brazil & Rio de Janeiro, Brazil (Koste, 1972f); Varzea near Santarem, Brazil (Koste, 1974b); Brazil (Koste & José de Paggi, 1982); Rio Yatapu & r. Uatuma, Amazonia, Brazil (Koste & Hardy, 1984); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Near Concepcion, Paraguay (Koste, 1986b); Near Panguana, Peru (Koste, 1988b); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Prov. Tucuman,, Santiago del Estero & Santa Fé, Argentina (José de Paggi, 1990); Amazonia, Brazil (Hauer, 1965; Thomasson, 1971)

Trichocerca brevidactyla (Daday, 1890)

- Palaearctic region: Hungarian plain (Bērziņš, 1978)

Trichocerca brevistyla (Lucks, 1912)

- Palaearctic region: Coast of S Finland (Eriksen, 1969)

Trichocerca capucina (Wierzejski & Zacharias, 1893)

- Palaearctic region: Lakes in Estonia (Haberman, 1978); Peat pits in Finland (Selin & Koskinen, 1988); Coast of S Finland (Eriksen, 1969); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Delta of river Danube, Romania (Godeanu & Zinevici, 1983; Zinevici & Teodorescu, 1990); Romania (Rudescu, 1960); Slovenia, ex-Yugoslavia (Markic, 1983); Lakes in Estonia (Määmets, 1983); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Buchseen, Bodensee, CH (Einsle, 1966); Mindelsee, Bodensee, CH (Einsle, 1967); C Finland (Eloranta, 1982b); Lake Bolsena & L. Vico, C Italy (Ferrari, 1972); Hortobagy Nat. Park, Hungary (Forró & Ronkay, 1983); Lago Maggiore, Italy (Tonolli, 1962; Goldman *et al.*, 1968); Lake Biwa, Japan (Yamamoto, 1967 & 1968; Hillbricht-Ilkowska, 1983a); Lake Stechlinsee & L. Nehmitzsee, Brandenburg, Germany (Kasprzak & Ronneberger, 1982); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Lake Mikolajskie & L. Sniardny, ibid. (Kosicki, 1960); Lago di Giulianello, Latium, Italy (Stella & Margaritora, 1972; Ferrara & Margaritora, 1977); Lakes in S Sweden (Arnemo *et al.*, 1968); Groszer Müggelsee, Berlin, Germany (Barthelmes, 1960); Small lakes in Estonia (Haberman & Määmets, 1987); Lake Bielersee, CH (Berner-Frankhauser, 1986); Middle course of river Po, Italy (Ferrari *et al.*, 1986); Bannsee, near Seoon, Bavaria, Germany (Preiszler, 1977a & 1977b); Lake Balaton, Hungary (Zankai & Ponyi, 1971a; Zankai, 1968); Grasmere Lake, Lake district, UK (Elliott, 1977); Nat. Res. Neeruti Lakes, NE Estonia (Määmets A. & A., s.d.); Karstic Lake Percile, Latium, Italy (Margaritora *et al.*, 1981); Lakes C Sweden (Pejler, 1964 & 1965); Po river, Italy (Ferrari *et al.*, 1989); Lake Sempachersee, CH (Zimmermann, 1974); C Sweden (Pejler, 1972); Darnsee near Epe, NW Germany (Koste, 1962); Lake Mälaren, Sweden (Grönberg, 1975); Fish ponds near Warsaw, Poland (Hillbricht-Ilkowska, 1964); Lac Léman, Fr./CH (Martin, 1961; Rougier, 1969; Naef & Martin, 1985; Balvay, 1989; Balvay & Laurent, 1981, 1989, 1990a & 1990b); Rybnik dam reservoir, Poland (Krzanowski, 1979); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Volga river & Volgograd reservoir, R.F. (V'yushkova & Lakhnova, 1974); Lakes in Soviet Karelia, R.F. (Filimonova, 1976); Dniester Estuary, R.F. (Polishchuk, 1976); In large & medium-size rivers, tributary to Kiev reservoir (Polishchuk, 1987); Lake Ladoga, R.F. (Nikolayev & Petrova, 1978); All over Europe. Cosmopolitan (Bērziņš, 1978); Lake Volvi, Macedonia, Greece (Zarfdjian *et al.*, 1990); Rügener, Hiddensee & Darszer Boddengegewässer, NE Germany (Schwarz, 1963); Etang du Brochet, Rambouillet, France (Pourriot *et al.*, 1969); Small lakes SE of Gothenburg, W Sweden (Morling & Pejler, 1990); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b); Gulf of Finland (Bērziņš, 1960); Lake Kus, Turkey (Ustaglu & Balik, 1990a); Leżnica & Włodawa Lake district, E Poland (Kowalczyk & Radwan, 1982; Radwan, 1984); Japanese inland waters (Yamamoto, 1960a); Lakes Vechtplassen, NL (Leentvaar, 1969); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Lake Constance, Germany (Kiefer, 1968; Walz *et al.*, 1987); River Danube (Enaceanu & Brezeanu, 1970); River Elbe estuary, Germany (Schulz, 1961); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960); Impoundment Weidatalsperrre, E Thüringen, G. (Hoffmann, 1966); Bolshoe Lake, White Sea, R.F. (Ghilarov, 1965); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Lac de Nantua, Ain, France (Balvay, 1985); All over the Netherlands (Dresscher, 1976); Bog pond near Budapest, Hungary (Nogradi, 1962); Lake Zürichsee, CH (Nipkow, 1961); Lake Erken, Sweden (Nauwerck, 1963); Karelian inland waters, R.F. (Kutikova, 1965); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Lake Ösbysjön, near Stockholm, Sweden (Pejler, 1961); Lac d'Annecy, France (Balvay & Druart, 1992); Black Sea (Rudescu, 1961); In reed borders of the Danube delta, Romania (Rudescu, 1968); 3 lakes of Sosnowickie, Poland (Radwan, 1969); Gemünderer Maar, Weinfelder Maar & Pulvermaar, Eifel, Germany (Oehms & Seitz, 1992); Hamburg area, Germany (Rühmann, 1961); Hamburg harbour, Germany (Rühmann,

1962); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Etang de la Tour, Ile-de-France, France (Pourriot, 1965); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990); Kakhova reservoir, near Lysaya Gora, R.F. (Pashkova, 1992); Aegean lakes, Macedonia, ex-Yugoslavia & Greece (Serafimova-Hadzisce, 1974); Fens in Genk, Belgium (Schepens, 1960); Pool, S of Trondheim, Norway (Thomasson, 1979); Lake Ohrid, Balkan peninsula (Stankovic, 1960); Lago Trasimeno, Italy (Taticchi, 1968); Serbian section of river Danube (Zivkovic, 1987); Lake Stechlinsee, Brandenburg, Germany (Ronneberger & Schönborn, 1985); Waters of the Szczeczin Lagoon, Poland (Wiktor, 1960); Lake Biwa-ko, Japan (Yamamoto, 1960b)

- Nearctic region: Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1972); Canada (Chengalath, 1977); 3 lakes in N Carolina, USA (Lemly & Demmick, 1982); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Cape Herschel, Ellesmere Island, NWT, Canada (Nogrady & Smol, 1989); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); Newfoundland lakes, Canada (Chengalath *et al.*, 1984); Mt. Tremblant Prov. Park, Québec, Canada (Nogrady, 1980); Jenny Lake, W North America (Thomasson, 1962)

- Oriental region: Philippines (Mamaril & Fernando, 1989); Taiwan & Philippines (Sudzuki, 1989); Perennial pond at Chotanagpur, India (Srivastava, 1981)

- Australian region: Reservoir in Australia (Bērziņš, 1982); Farm dams near Gloucester, NSW, Aust. (Sudzuki & Timms, 1980); NSW & Qld, Aust. (Shiel & Koste, 1979); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Magela Creek billabongs, NW Aust. (Shiel & Koste, 1983); Tasmania (Koste & Shiel, 1987); Uncommon; NT, Qld, Tas & Vic (Shiel & Koste, 1992)

- Neotropical region: Orinoco river basin, Venezuela (Vasquez, 1984a); Lower Orinoco river and its delta, Venezuela (Vasquez & Rey, 1989); Lago do Macaco, Brazil (Koste, 1989a); Amazonia, Brazil (Hauer, 1965; Thomasson, 1971); Amazonia, Brazil. Cosmopolitan (Koste, 1972f); Brazil, Venezuela & Panama (Koste & José de Paggi, 1982); Rio Yatapu & r. Uatumá, Amazonia, Brazil (Koste & Hardy, 1984); Lago Camaleão, Amazonia, Brazil (Koste, Robertson & Hardy, 1984); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Lake Hemmen, Rio Snia Missu, Mato Grosso, C Brazil (Green, 1972); Lago do Castanho, C Amazonia, Brazil (Schaden, 1977b); Lago Yurucui & Rio Negro, Amazonia, Brazil (Schaden, 1977c); Lake Cocibolca & L. Masaya, Nicaragua (Moreno *et al.*, 1992)

Trichocerca cavia (Gosse, 1886)

- Palaeartic region: Romania (Rudescu, 1960); Lake Myvatn, Iceland (Adalsteinsson, 1979); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1989); River Klyas'ma, R.F. (Bogoslawski, 1961); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); Sierra Nevada, Spain (Morales-Baquero, 1987); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Lake Taltowisko, Mazurian lakeland, Poland (Klimowicz, 1982); N swedish Lapland (Pejler, 1962a); Alpine lakes, Sierra Nevada, Spain (Morales-Baquero *et al.*, 1989); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); All over Europe. Cosmopolitan (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1970); Lake Balaton, Hungary (Zankai, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b); SW, W & N Iceland (De Ridder, 1966a, 1967a, 1969b, 1970 & 1972); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Peat bog Chambedaze, Puy-de-Dôme, France (Francez, 1982); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Leżnica-Włodawa Lake District, E Poland (Radwan, 1984); Pelagic zone of the Leżnica-Włodawa lakes, Pol. (Radwan, 1975); Water-courses in the Bucegi mts., Romania (Godeanu, 1963b); Small permanent ponds & marshy zones, ibid. (Godeanu, 1969a); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Anatolia, Turkey (Segers *et al.*, 1992); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); River Salzach & its tributaries, Austria (Donner, 1970); Predeltaic area of the Black Sea, Romania (Popescu-Marinescu, 1978); Dübener Heide, Germany (Wulfert, 1960b); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Nat. Res. Meyendel; Nat. Res. Het Hol near Kortenhoef, NL (Dresscher, 1976); In reed borders of the Danube delta, Romania (Rudescu, 1968); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Ponds in Botanical Garden Gent, Belgium (Segers *et al.*, 1991b); Spain : Granada (Velasco, 1990); Baginiec II pond & Lipowy pond, Poland (Lewkowicz, 1972); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)

- Nearctic region: Crater Lake, Arizona, USA (Green, 1986); Kingston, Ontario, Canada (Nogrady, 1989); NE Canada (Chengalath & Koste, 1983); Cap Breton Ilsand, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Illinois, USA (Bērziņš, 1984); Laurel Creek reservoir, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980)

- Oriental region: India (Sharma & Michael, 1980; Sudzuki, 1989)

- Australian region: NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait,

Shiel & Koste, 1984); Creek and river in Aust. (Bērziņš, 1982); Swamp near Dunedin, S Island, New Zealand (Bērziņš, 1982a); Tasmania (Koste & Shiel, 1987); Vic, Aust. (Koste & Shiel, 1980c); Rare, one record each from NT, Tas & Vic (Shiel & Koste, 1992); Wellington, New Zealand (Russell, 1960)

- Neotropical region: Venezuela (Koste & José de Paggi, 1982); Prov. Rio Negro, Argentina (José de Paggi, 1990); Araucarian lakes, E of the Andes, Chile (Thomasson, 1963b)

Trichocerca chattoni (de Beauchamp, 1907)

- Palaearctic region: Impoundment on Rio Guadiana, C Spain (Armengol, 1982); Spanish reservoirs (Guiset, 1977); 3 reservoirs in Rio Guadiana, C Spain (Armengol *et al.*, 1987); Lake Biwa, Japan (Yamamoto, 1967 & 1968); Azibo reservoir, near Porto, Portugal (Vascoucelos, 1990); W & C Eur. plains (Bērziņš, 1978); Southwest islands of Japan (Sudzuki *et al.*, 1992); All over Spain (Velasco, 1990)

- Oriental region: Asia (Bērziņš, 1978); Lakes Sunda-expedition (Serruya & Pollingher, 1984); Sri Lanka (Fernando, 1980a); Taiwan (Sudzuki, 1988); Taiwan, Malaysia, Singapore, Indonesia & Sri Lanka (Sudzuki, 1989); Indonesia (Koste, 1972f); Sri Lanka. Fairly common in lakes & ponds (Chengalath, Fernando & Koste, 1973); Taiwan. Distribution : warm waters (Sudzuki, 1991); W & E Malaysia & Singapore (Fernando & Zankai, 1981)

- Australian region: Sidney reservoirs, NSW, Aust. (Thomasson, 1973); NSW, Aust. (Shiel & Koste, 1979); (Koste & Shiel, 1980c); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area (Koste & Shiel, 1983a; Tait, Shiel & Koste, 1984); Rare. Confirmed from ca 14 localities (Shiel & Koste, 1992)

- Neotropical region: Araucarian Lakes, E of the Andes, Chile (Thomasson, 1963b); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Rio Tocantins, Lago do Rei & Rio Negro, Amazonia, Brazil (Schaden, 1977c); Lago Taruma-Mirim, Amazonia, C Brazil (Hardy, 1980); Lago Castanho, L. Cristalino & L. Taruma-Mirim, Amazonia, Brazil (Serruya & Pollingher, 1984); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980); Prov. Rio Negro, Catamarca, Chaco & Santa Fé, Argentina (José de Paggi, 1990); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Orinoco river basin, Venezuela (Vasquez, 1984a); Caroni river basin, Venezuela (Vasquez, 1984b); Lower Orinoco r. and its delta, Venezuela (Vasquez & Rey, 1989); Provinces in NW Argentina (José de Paggi, 1989); Lago Açu, Maranhao, Brazil (Reid & Turner, 1988); Brazil, Panama & Venezuela (Koste & José de Paggi, 1982); Rio Yatapu & R. Uatumá, Amazonia, Brazil (Koste & Hardy, 1984); Amazonia, Brazil (Hauer, 1965b; Thomasson, 1971;; Koste, 1972f)

Trichocerca collaris (Rousselet, 1896)

- Palaearctic region: Romania (Rudescu, 1960); Coast of S Finland (Eriksen, 1969); Asiatic Turkey (Dumont & De Ridder, 1987); C & S Sweden (Bērziņš & Pejler, 1987 & 1989d); N Swedish Lapland (Pejler, 1962a); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); Europe, Japan, China, R.F. (Koste, 1972f); In peat bogs of the Massif Central, France (Batut, 1965); All over Europe (Bērziņš, 1978); Lake Balaton, Hungary (Zankai, 1968); N & W Iceland (De Ridder, 1969b & 1972); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Peat bog Chambedaze, Massif Central, France (Francez, 1982, 1987 & 1988); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Japanese inland waters (Yamamoto, 1960a); Predeltaic area of Black Sea, Romania (Popescu-Marinescu, 1974); River Maas, Rijn, Waal & Yssel; L. Gerritsfles, near Kootwijk, NL (Dresscher, 1976); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Peat bogs in the Paris suburb, France (Pourriot, 1965); Tarn in Värmland, Sweden (Thomasson, 1970)

- Nearctic region: Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); N America (Koste, 1972f); Northern Michigan inland lakes (Stemberger, 1990)

- Oriental region: Sumatra (Koste, 1972f); Indonesia (Sudzuki, 1989)

- Australian region: ? Tasmania (Koste & Shiel, 1987); NT, Aust. (Koste & Shiel, 1980c); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); New Zealand (Koste, 1972f); Cosmopolitan in acid waters with sphagnum; rare; several records from Tasmania, only one from mainland : Magela Creek, NT (Shiel & Koste, 1992); Takahe valley, New Zealand (Russell, 1960)

- Neotropical region: Argentina, Brazil & Venezuela (Koste & José de Paggi, 1982); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Concepcion, Paraguay (Koste, 1986b); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Amazonia, Brazil (Hauer, 1965b; Thomasson, 1971; Koste, 1972f); Lago Paroni, Brazil (Koste, 1974a); Varzea near Santarem, Brazil (Koste, 1974b); Prov. Corrientes, Argentina (José de Paggi, 1990); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980); Ecuador : Oriente & Costa (Koste & Böttger, 1992)

Trichocerca cryptopus Hauer, 1938

- Oriental region: Singapore & Indonesia (Sudzuki, 1989); Singapore & Java (Sudzuki, 1991)

Trichocerca cylindrica (Imhof, 1891)

- Palaearctic region: Lake Biwa-ko, Japan (Yamamoto, 1960b); Romania (Rudescu, 1960); Delta of river Danube, Romania (Prunescu-Arion, 1989; Zinevici & Teodorescu, 1990); Nat. Res. Neeruti Lakes, Estonia (Määmets A & A., s.d.); Lakes in C Sweden (Pejler, 1964 & 1965); Lago Maggiore, Italy (Tonolli, 1962); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); Eutrophic waters in E Poland (Radwan & Popiolek, 1989); Carp ponds in S Poland (Zurek, 1980); Lake Mälaren, Sweden (Grönberg, 1975); Fish ponds near Warsaw, Poland (Hillbright-Ilkowska, 1964); Lac Léman, Fr./CH (Balvay, 1989); (Balvay & Laurent, 1981, 1989 & 1990a); Lac d'Aydat, Massif Central, France (Oulad Ali *et al.*, 1990; Lair, 1991); Volga river & Volgograd reservoir, R.F. (V'yushkova & Lakhnova, 1974); Lakes in Soviet Karelia, R.F. (Filimonova, 1976); Lake Groszer Müggelsee, Berlin, Germany (Barthelmes, 1960); Heiluntsian province, China (Bogoslawski, 1962); 2 lakes in N Sweden (Axelson, 1961); Lakes in Estonia (Haberman, 1978; Määmets, 1983); Coast of S Finland (Eriksen, 1969); Lake Ormajärvi, S Finland (Cajander, 1983); Lake Biwa, Japan (Hillbright-Ilkowska, 1983a); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); District Ostrava, CSR (Dvorakova, 1960b); Lapland State Preserve, R.F. (Ghilarov, 1972); Oligotrophic neutral to alkaline lakes in the NW R.F. (Salazkin, 1971); All over Europe (Bērziņš, 1978); Experimental fishponds near Warsaw, Poland (Spodniewska *et al.*, 1966); Lake Balaton, Hungary (Zankai, 1968); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Etang du Brochet, Rambouillet, France (Pourriot *et al.*, 1969); Lake Kus, Turkey (Ustaoglu & Balik, 1990a); Leżna-Włodawa Lake District, E Poland (Radwan, 1984); Japanese inland waters (Yamamoto, 1960a); Serbian section of river Danube (Zivkovic, 1987); 3 lakes of Sosnowickie, Poland (Radwan, 1969); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); River Danube (Enaceanu & Brezeanu, 1970); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Artificial lakes near Köln, Germany (Trahms, 1972); River Salzach & its tributaries, Austria (Donner, 1970); Lac d'Aydat, Massif Central, France (Lair, 1990); Pond in Enschede; surroundings of Nijmegen; de Gerritsfles near Kootwijk; Kil van Hurwenen; area between Linge & Rhene; Susteren (Lb); pool Vechten near Utrecht; Loosdrecht Lakes, NL (Dresscher, 1976); Karelian inland waters, R.F. (Kutikova, 1965); Lakes Aydat, Chambon & Bourdouze, Puy-de-Dôme, France (Persoone *et al.*, 1968); Leżna-Włodawa Lake District, E Poland (Kowalczyk & Radwan, 1982); Rés. biol. de Villars-la-Dombes, France (Rostan, 1976); Lakes in the Leżna-Włodawa area of E Poland (Radwan, 1976); Hamburg area, Germany (Rühmann, 1961); Middle course of river Loire, France (Lair & Sargas, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Ponds in the Paris suburb, France (Pourriot, 1965); Tarn in Värmland, Sweden (Thomasson, 1970); Spain : Barcelona, Valencia & Zamore; Portugal : Beja & Portalegre (Velasco, 1990); Experimental ponds at Golysz, Poland (Lewkowicz, 1971; Zurek, 1974)
- Nearctic region: 2 lakes in Idaho, USA (Buscemi, 1964); Acid lakes in Maine, USA (Brett, 1989); 18 arctic & subarctic lakes, Canada (Moore, 1978); Ontario, Canada (Chengalath, Fernando & George, 1971; Chengalath & Fernando, 1973); NE Canada (Chengalath & Koste, 1983); 3 lakes in N Carolina, USA (Lemly & Dimmick, 1982); Wisconsin, USA (Bērziņš, 1984); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Illinois river, USA (Brown *et al.*, 1989); Lakes Adirondack, NY State, USA (Siegfried *et al.*, 1984 & 1989); Northern Michigan inland lakes (Stemberger, 1990); Clearwater L. & Swan L., near Sudbury, Ontario (Yan & Geiling, 1985); Columbia Lakes, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Newfoundland lakes, Canada (Chengalath *et al.*, 1984); Lake 122 & L. 132, NW Ontario, Canada (Schindler & Novén, 1971); Silver Lake, Adirondack mts., NY State, USA (Siegfried, 1991); Crecy Lake, New Brunswick, Canada (Smith, 1969); Experimental ponds, C Florida, USA (Fry & Osborne, 1980); Greens Lake, Beaver Island, Michigan, USA (Foran & King, 1982); Atchafalaya river basin, Louisiana, USA (Holland *et al.*, 1983); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980); Lake Oglethorpe, NE Georgia, USA (Orcutt & Pace, 1984); Laurentian Great Lakes (Stemberger, 1979); Desaulniers reservoir, Québec, Canada (Pinel-Alloul *et al.*, 1982); James Bay area, Canada (Pinel-Alloul *et al.*, 1979); R. Opinaca & Small Opinaca, James Bay area, N Québec, Canada (Armengol *et al.*, 1983); Bayou de Siard, Monroe, Louisiana, USA (Gallagher, 1965); Long Lake, N Carolina, USA (Moore, 1980); Lake Ontario, Canada (Nauwerck, 1978); Bay of Quinte, Lake Ontario, Canada (Roff & Kwiatkowski, 1977); Lakes in the Churchill Falls area of Labrador (Duthie & Ostrowsky, 1974)
- Oriental region: Dal Lake, Kashmir Himalayan lake, India (Vass & Zutshi, 1983); Fishpond in Burdwan, W Bengal, India (Mandal, 1976); India (Sharma, 1980); Sri Lanka (Fernando, 1980a); Taiwan, Sri Lanka & India (Suzuki, 1989); Phewa Tal, Pokhara valley, Nepal (Daems & Dumont, 1974); Sri Lanka. Rare; found in lakes (Chengalath & Koste, 1973); Fishponds in Thailand (Boonsom, 1984a); Thailand (Boonsom, 1984b); 2 sewage channels of Gwalior, India (Saksena & Kulkarni, 1986); 2 Kumaun Himalayan lakes, UP, India (Sharma & Pant, 1984 & 1985); Lakes of the Sunda Expedition (Serruya & Pollingher, 1984)
- Australian region: Lake Sorrell, Tasmania (Suzuki, 1967); Lake Sorrell in Tasmania (after Suzuki, 1967); L. Encumbene, Snowy mts. & L. Dartmouth (Shiel & Koste, 1992)
- Neotropical region: Ecuador : Sierra (Koste & Böttger, 1989); Orinoco river basin, Venezuela (Vasquez,

1984a); Argentina & Panama (Koste & José de Paggi, 1982); Prov. Rio Negro, Chaco & Corrientes, Argentina (José de Paggi, 1990); Araucarian lakes, E of the Andes, Chile (Thomasson, 1963b)

***Trichocerca dixonnuttalli* (Jennings, 1903)**

- Palaearctic region: Romania (Rudescu, 1960); Afghanistan (Bērziņš, 1961); Seat Ness, Shetland (De Smet, 1988); Koyama Ike, a coastal pond in Japan (Suzuki & Fukuta, 1989); Lac Léman, Fr./CH (Balvay *et al.*, 1985; Balvay & Laurent, 1989, 1990a & 1990b); C Eur. mountains, Carpathes, Pontic province, C & E Eur. plains, Baltic prov., UK, Caucasus (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1968 & 1970); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Southwest islands of Japan (Suzuki, 1990; Suzuki *et al.*, 1992); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Lakes, N of Bucharest, Romania (Godeanu, 1969b); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Lac de Nantua, Ain, France (Balvay, 1985); Province Zuid-Limburg, NL (Dresscher, 1976); Lac d'Annecy, France (Balvay & Druart, 1992); River Loire in St. Laurent-les-Eaux, France (Pourriot *et al.*, 1982); Pelagic zone of the Leżnica-Włodawa lakes, P. (Radwan, 1975); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Serbian section of river Danube (Zivkovic, 1987)

- Nearctic region: Wisconsin, Illinois & Kentucky, USA (Bērziņš, 1984); NW Canada (Chengalath & Koste, 1987); Altiplanic Central, Mexico (Vilaclara & Sladecik, 1989); Lake Wabamun, Alberta, Canada (Gallup & Hickman, 1975); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980)

- Oriental region: Sri Lanka (Fernando, 1980a); Kalimantan (Borneo), Indonesia (Koste, 1988a); Taiwan, Indonesia, Malaysia, Singapore & Sri Lanka (Suzuki, 1989); Sri Lanka. Only in ricefields, but in fairly large numbers (Chengalath, Fernando & Koste, 1973); W & E Malaysia & Singapore (Fernando & Zankai, 1981); Impoundment Kinda, C Burma (Koste & Tobias, 1990); Lake Pakis & L. Singkarak (Serruya & Pollingher, 1984)

- Australian region: Lake Barracoota, Aust. (Bērziņš, 1982); Qld, Aust. (Shiel & Koste, 1979); S Tasmania & Qld, Aust. (Koste, Shiel & Tan, 1988; Shiel & Koste, 1992)

- Neotropical region: Lago Açu, Maranhão, Brazil (Reid & Turner, 1988); Amazonia, Brazil (Koste, 1972f); Brazil, Panama & Venezuela (Koste & José de Paggi, 1982); Near Panguana, Peru (Koste, 1988b); Chilean lakes (Schmid-Araya, 1991); Costa Rica & Panama (Collado *et al.*, 1984); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980); Araucarian lakes, W of the Andes, Chile (Thomasson, 1963b); Lago do Passarinho & L. Buiuçu, Amazonia, Brazil (Schaden, 1977c)

***Trichocerca elongata elongata* (Gosse, 1886)**

- Palaearctic region: Peat bogs Chambédaize, Puy-de-Dôme, France (Francez, 1982 & 1987); River Danube (Enaceanu & Brezeanu, 1970); Romania (Rudescu, 1960); Asiatic Turkey (Dumont & De Ridder, 1987); District Ostrava, CSR (Dvorakova, 1960b); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Po river, Italy (Ferrari *et al.*, 1989); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Achmer Grasmoor, NW Germany (Koste, 1968b); Lake Myvatn, Iceland (Adalsteinsson, 1979); Heilunshan province, China (Bogoslawski, 1962); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Coast of S Finland (Eriksen, 1979); River Loire, France (Lair, 1980); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); Lake Alfsee, NW Germany (Koste & Poltz, 1987); Lac Léman, Fr./CH (Balvay *et al.*, 1985; Balvay & Laurent, 1989 & 1990a); Kiev reservoir, R.F. (Polishchuk, 1987); Soviet section of river Danube, R.F. (Parchuk, 1985); All over Europe (Bērziņš, 1978); "Altes Land" near Hamburg, Germany (Caspers & Heckman, 1981 & 1982); Middle course of river Po, Italy (Ferrari *et al.*, 1986); Japanese inland waters (Yamamoto, 1960a); River Grabia basin, Poland (Pawlowski, 1970); Lake Balaton, Hungary (Zankai, 1968); Mt. Fuji, Japan (as *Paratrichocerca elongata*) (Suzuki, 1971); Lake Donk, Overmere, Belgium (Dumont, 1968b); N & W Iceland (De Ridder, 1969b, 1970 & 1972); Peat bogs Chambédaize & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Southwest islands of Japan (Suzuki *et al.*, 1992); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Samrée, Belgium (De Ridder, 1980 & 1986); Pools with sphagnum in Fontainebleau forest & in the Sologne, Ile-de-France, France (Pourriot, 1970); Middle course of river Loire, France (Lair & Sargos, 1981); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990); Pool, S of Trondheim, Norway (Thomasson, 1979); Lago Trasimeno, Italy (Taticchi, 1968); Serbian section of river Danube (Zivkovic, 1987); Lake Biwa-ko, Japan (Yamamoto, 1960b); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Lakes of the Black Sea shore, Romania (Godeanu, 1970b); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); River Vistula in the Warsaw area, Poland (Klimowicz, 1981); Small water bodies in the Mikolajski area, Poland (Klimowicz, 1967); Düben Heide, N Germany (Wulfert, 1961); Lake Aydat, Massif Central, France (Lair & Ayadi, 1989); Peat bog Chambédaize, Massif Central, France (Francez, 1988); River Rijn, Waal & Yssel; Maria- & Helenapeel; Nat. Res. de Gerritsfles near Kootwijk; R. Maas; Broekhuizen &

Susteren, Limburg, NL (Dresscher, 1976); Bog pond near Budapest, Hungary (Nogrady, 1962); Karelian inland waters, R.F. (Kutikova, 1965); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); Parczew area peat bogs, Poland (Radwan, 1974); Peat bogs Poiana Stampei, Bukowina, Poland (Rudescu, 1966); Mountain Fuji area, Japan (Suzuki, 1978a); Lakes around mountain Fuji, Japan (Suzuki, 1971); Lake Biwa, Japan (Yamamoto, 1967)

- Nearctic region: Walker Lake, Arizona, USA (Green, 1986); Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Canada (Chengalath, 1977); Gregg's Lake & Parvin Lake, Colorado, USA (Pennak, 1966); Kentucky, USA (Bērziņš, 1984); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Colorado, USA (Turner, 1988); Acid lakes in Maine, USA (Brett, 1989); Kingston, Ontario, Canada (Nogrady, 1989); Laurentian Great Lakes (Stemberger, 1979); Pinehurst Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamootttil, 1975); Bearverskin Lake, Kejumkujik Nat. Park, Nova Scotia, Canada (Blouin *et al.*, 1984); Adirondack lakes, NY State, USA (Siegfried *et al.*, 1984)

- Oriental region: Kalimantan (Borneo), Indonesia (Koste, 1988a); Malaysia, Singapore & Indonesia (Suzuki, 1989); W & E Malaysia & Singapore (Fernando & Zankai, 1981); Ricefields in NE Thailand (Heckman, 1979); Sri Lanka (Mendis & Fernando 1962); (Fernando, 1969)

- Australian region: SW Aust. (Koste, Shiel & Brock, 1983); NSW & Qld, Aust. (Shiel & Koste, 1979); River in Australia (Bērziņš, 1982); River Murray, Aust. (Koste, 1979b); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Pancontinental, common in billabongs (Shiel & Koste, 1992); South Canterbury, New Zealand (Russell, 1960)

- Neotropical region: Ecuador : Oriente (Koste & Böttger, 1989 & 1992); Lake Cocibolca, Nicaragua (Moreno *et al.*, 1992); River Saladillo basin, Argentina (José de Paggi & Koste, 1988); Lower Orinoco river and its delta, Venezuela (Vasquez & Rey, 1989); Lago Paroni, Brazil (Koste, 1974a); Argentina, Brazil & Paraguay (Koste & José de Paggi, 1982); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Chilean lakes (Schmid-Araya, 1991); Prov. Santa Cruz, Chubut, Corriente, Santa Fé & Buenos Aires, Argentina (José de Paggi, 1990); Prov. Buenos Aires & Santa Cruz, Argentina; Paraguay (Olivier, 1965); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Lago do Castanho, C Amazonia, Brazil (Schaden, 1977b); Lago Tefé, Amazonia, Brazil (Schaden, 1977c)

Trichocerca elongata braziliensis (Murray, 1913)

- Oriental region: W & E Malaysia & Singapore (Fernando & Zankai, 1981)

Trichocerca euodonta Hauer, 1938

- Oriental region: Malaysia & Indonesia (Suzuki, 1989)

- Australian region: River in Aust. (Suzuki, 1989); Vic, Aust. Is a synonym of *Tr. myersi* Hauer, 1931. Known only from North Sumatra and one unconfirmed Australian record from the Moorabool river at Ballan, Vic (Bērziņš, 1982) (Shiel & Koste, 1992)

Trichocerca falx Edmondson, 1936

- Nearctic region: Canada (Chengalath, 1977)

Trichocerca flagellata Hauer, 1937

- Palaearctic region: Southwest islands of Japan (Suzuki *et al.*, 1992)

- Oriental region: Malaysia (Brandorff, Koste & Smirnov, 1982); Malaysia & India (Suzuki, 1989); India (Sharma & Michael, 1980); W & E Malaysia & Singapore (Fernando & Zankai, 1981)

- Australian region: NT, Aust. (Koste & Shiel, 1980c); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Lake Purumbete, freshwater Aust. crater lake (Green, 1981); 3 localities known : Magela Creek, NT; Cape Cork, Qld & Lake Purumbete, Vic (Shiel & Koste, 1992)

- Neotropical region: Orinoco river basin, Venezuela (Vasquez, 1984a); Lower Orinoco river & its delta, Venezuela (Vasquez & Rey, 1989); Brazil & Venezuela (Koste & José de Paggi, 1982); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Concepcion, Paraguay (Koste, 1986b); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980)

Trichocerca flava (Voronkov, 1907)

- Palaearctic region: NW Russian taiga (Bērziņš, 1978)

Trichocerca fusiformis (Gosse, 1886)

- Palaearctic region: Peat bog Poiana Stampei, Bukowina, Rom. (Rudescu, 1966); In reed borders of the Danube delta, Rom. (Rudescu, 1968)

Trichocerca gracilis (Tessin, 1890)

- Palaearctic region: Area between rivers Linge & Waal, NL (Dresscher, 1976); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Romania (Rudescu, 1960); Mt. Hoverla,

Maramarosch Alps, Hungary (Varga, 1962); Darnsee near Epe, NW Germany (Koste, 1962); Europe, R.F., Tibet (Koste, 1972f); Fish ponds near Warsaw, Poland (Hillbright-Ilkowska, 1964); Southwest islands of Japan (Sudzuki, 1990; Sudzuki *et al.*, 1992); Lakes, N of Bucarest, Romania (Godeanu, 1969b); Ponds in Oltina, SE Dobrogea, Romania (Popescu-Gorj & Costea, 1961); Pelagic zone of the Leżna-Włodawa lakes, P. (Radwan, 1975); Hamburg area, Germany (Rühmann, 1961); Lago Trasimeno, Italy (Taticchi, 1968); River Cerna basin, Romania (Brezeanu & Marinescu-Popescu, 1964)

- Nearctic region: Little Swede n° 1 Lake, Colorado (Pennak, 1966); Illinois & Missouri, USA (Bērziņš, 1984); Colorado, USA (Turner, 1988); N America (Koste, 1972f); Laurentian Great Lakes (Stemberger, 1979); San Francisco, USA (Nogrady, 1983)

- Oriental region: Java (Koste, 1972f); Singapore (Sudzuki, 1989)

- Australian region: Qld, Aust. (Shiel & Koste, 1985); Europe, N & S America. Rare in Aust. Single record, Salomon Dam, Palm Island, N Qld (Shiel & Koste, 1992)

- Neotropical region: Rio Tocantins, Lago Moratu, L. do Castanho & L. Tefé, Amazonia, Brazil (Schaden, 1977c); Lago do Castanho, Amazonia, C Brazil (Schaden, 1977b); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Lower Orinoco & its delta, Venezuela (Vasquez & Rey, 1989); Amazonia, Brazil (Koste, 1972f); Brazil, Argentina, Venezuela & Panama (Koste & José de Paggi, 1982); Prov. Corrientes, Argentina (José de Paggi, 1990); Costa Rica (Collado *et al.*, 1984); Flooded savanna at Mantecal, Venezuela (Michelangeli *et al.*, 1979/1980)

Trichocerca harveyensis Myers, 1942

- Palaearctic region: In peat bogs in the Ile-de-France, France (Pourriot, 1965 & 1970); In peat bogs in the Ile-de-France et en Sologne, France (Pourriot, 1966); W Eur. plain (Bērziņš, 1978)

Trichocerca helminthoides (Gosse, 1886)

- Palaearctic region: Almost all over Europe (Bērziņš, 1978)

- Australian region: Campbell Island (Russell, 1960)

Trichocerca heterodactyla (Tschugunoff, 1921)

- Palaearctic region: N Part of Caspian Sea & L. Aral (Mordukhai-Boltovskoi, 1979); NW Russian taiga (Bērziņš, 1978); Southwest islands of Japan (Sudzuki *et al.*, 1992)

Trichocerca iernis (Gosse, 1887)

- Palaearctic region: Small water bodies in the Mikolajki area, Poland (Klimowicz, 1967); Romania (Rudescu, 1960); Lake Myvatn, Iceland (Adalsteinsson, 1979); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); River Klyas'ma, R.F. (Bogoslawski, 1961); Coast of S Finland (Eriksen, 1969); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Delta of river Danube, Romania (Prunescu-Arion, 1989); N Swedish Lapland (Pejler, 1962a); Italy, Alps, all over Europe. Caspian area. Cosmopolitan (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1970 & 1973); Lake Balaton, Hungary (Zankai, 1968); N & W Iceland (De Ridder, 1969b & 1972); Japanese inland waters (Yamamoto, 1960a); Southwest islands of Japan (Sudzuki *et al.*, 1992); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Dübener Heide, N Germany (Wulfert, 1961); Nat. Res. "Het Hol", near Kortenhoef, NL (Dresscher, 1976); Karelian inland waters, R.F. (Kutikova, 1965); Parczew area peat bogs, Poland (Radwan, 1974); Pelagic zone of the Leżna-Włodawa lakes, Poland (Radwan, 1975); Fens in Genk, Belgium (Schepens, 1960); Serbian section of river Danube (Zivkovic, 1987); Spain : Lerida & Orense (Velasco, 1990); Lake Biwa-ko, Japan (Yamamoto, 1960b); River Cerna basin, Romania (Brezeanu & Marinescu-Popescu, 1964); Lipowy pond, Baginice pond & pond Chylinski Maly II, Poland (Lewkowicz, 1972); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)

- Nearctic region: Canada (Chengalath, 1977); Illinois & Kentucky, USA (Bērziņš, 1984); Kingston, Ontario, Canada (Nogrady, 1989); Lakes in Santee, California, USA (Nogrady, 1982); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); Calabogie Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975)

- Oriental region: Girikbe-Tank, Sri Lanka (Bērziņš, 1982a); India (Sharma & Michael, 1980); Indonesia & India (Sudzuki, 1989); Lake Manguao, Palawan, the Philippines (Davies & Green, 1990)

- Australian region: River Yarra, Australia (Bērziņš, 1982); Vic, Aust. (Shiel & Koste, 1985); Single unconfirmed record from Yarra river at Warburton, Vic (Shiel & Koste, 1992); West Coast, New Zealand (Russell, 1960)

- Neotropical region: Galapagos Islands (De Smet, 1987 & 1989); Argentina & Venezuela (Koste & José de Paggi, 1982); Near Concepcion, Paraguay (Koste, 1986b); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Prov. Chaco & Santa Fé, Argentina (José de Paggi, 1990); Santa Cruz, Galapagos Islands (Segers, 1991a)

Trichocerca inermis (Linder, 1904)

- Palaearctic region: Parczew area peat bogs, Poland (Radwan, 1974); Serbian section of river Danube

(Zivkovic, 1987); Romania (Rudescu, 1960); Coast of S Finland (Eriksen, 1969); All over Europe (Bērziņš, 1967); Upper part of Volga delta, R.F. (Chuykov, 1981); River Danube (Enaceanu & Brezeanu, 1970); Barlosche Kolk, near Wijlre, Drente, NL (Dresscher, 1976); W Iceland (De Ridder, 1969b & 1972)

- Nearctic region: N America (Bērziņš, 1978); Illinois & Kentucky, USA (Bērziņš, 1984); Northern Michigan inland lakes (Stemberger, 1990)

- Australian region: Vic, Aust. (Shiel & Koste, 1985); Single record : sheepwash billabongs, Yea, Vic (Shiel & Koste, 1992)

- Neotropical region: Araucarian lakes, W of the Andes, Chile (Thomasson, 1963b); Amazonia, Brazil (Thomasson, 1971); Chilean lakes (Schmid-Araya, 1991); Lower Orinoco & its delta, Venezuela (Vasquez & Rey, 1989)

Trichocerca insignis (Herrick, 1885)

- Palaearctic region: C Eur. mts., Carpathes, W, C & E Eur. plains, Baltic province, NW Russian taiga (Bērziņš, 1978); Coast of S Finland (Eriksen, 1969); N Swedish Lapland (Pejler, 1962a); Germany, CH, R.F., Sweden (Koste, 1972f); Pools with vegetation in the Fontainebleau forest, Ile-de-France, France (Pourriot, 1970); Karelian inland waters, R.F. (Kutikova, 1965); Romania (Rudescu, 1960); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); River Grabia basin, Poland (Pawlowski, 1970); Japanese inland waters (Yamamoto, 1960a); Water-courses in the Bucegi mts., Romania (Godeanu, 1963b); Temporary waters & marshy zones in the Bucegi mountains, Romania (Godeanu, 1969a); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Pool in Fontainebleau forest, Ile-de-France, Fr. (Pourriot, 1976); Pelagic zone of the Leżna-Włodawa lakes, P. (Radwan, 1975); Lake Chenbu, Changjiang basin, China (Weu, 1992); Peat bog Poiana Stampei, Bukowina, Rom. (Rudescu, 1966); Pool, S of Trondheim, Norway (Thomasson, 1979); Pond Chylinski Maly II & Lipowy pond, Poland (Lewkowicz, 1972)

- Nearctic region: Lake Tahoe, Western N America (Thomasson, 1962); Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); N America (Koste, 1972f; Bērziņš, 1978); Ontario, Canada & Wisconsin, USA (Bērziņš, 1984); Pinehurst Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Laurentian Great Lakes (Stemberger, 1979)

- Australian region: New Zealand (Koste, 1972f)

- Neotropical region: S America : Argentina, Venezuela, Brazil (Koste, 1972f); Ecuador : Oriente (Koste & Böttger, 1989); Orinoco river basin, Venezuela (Vasquez, 1984a); Caroni river basin, Venezuela (Vasquez, 1984b); Lago Paroni, Brazil (Koste, 1974a); Prov. Rio Negro & Corrientes, Argentina (José de Paggi, 1990); Araucarian lakes, E of the Andes, Chile (Thomasson, 1963b)

Trichocerca insolens Myers, 1936

- Nearctic region: Laurentian Great Lakes (Stemberger, 1979); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988)

Trichocerca insulana (Hauer, 1938)

- Palaearctic region: C Eur. plains & Asia (Bērziņš, 1978); Small water bodies in Soviet Karelia, R.F. (Filimonova, 1976); Afghanistan (Bērziņš, 1961)

- Nearctic region: Canada (Chengalath, 1977); Cranberry bog, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975)

- Oriental region: Swamp near Menggala, Sumatra (Bērziņš, 1982a); Singapore & Malaysia (Karunakaran & Johnson, 1978); Malaysia, Singapore & Indonesia (Suzuki, 1989)

- Australian region: River Yarra, Aust. (Bērziņš, 1982); NW Aust.: Magela Creek area (Koste, 1981b); ? Tasmania (Koste & Shiel, 1986b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); NT, Aust. (Shiel & Koste, 1985); Two records : Magela Creek, NT, & Lake Dulverton, Tasmania (Shiel & Koste, 1992)

- Neotropical region: Varzea near Santarem, Brazil (Koste, 1974b); Brazil (Koste & José de Paggi, 1982); Mato Grosso, C Brazil (Turner & Da Silva, 1992)

Trichocerca intermedia (Stenoos, 1898)

- Palaearctic region: Brook in Lao-Shan mts. & pool in Kaitao, China (Wulfert, 1968); Spain : Ibiza (Baleares) (Velasco, 1990); Bog pond near Budapest, Hungary (Nogradi, 1962); Romania (Rudescu, 1960); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Minorca, Spain (de Manuel, 1990); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Coast of S Finland (Eriksen, 1969); All over Europe (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1960, 1968 & 1970); Lake Balaton, Hungary (Zankai, 1968); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c)

- Nearctic region: Colorado, USA (Turner, 1988); N America (Bērziņš, 1978); NW Canada (Chengalath

& Koste, 1987); Arctic N America (Chengalath & Koste, 1989); Kingston, Ontario, Canada (Nogrady, 1989); Canada (Chengalath, 1977); Parvin Lake, Colorado, USA (Pennak, 1966); Illinois, USA (Bērziņš, 1984)

- Australian region: Tasmania (Koste & Shiel, 1987); Bromfield Swamp, Tropical Qld Aust. crater lake (Green, 1981); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Qld, Aust. (Shiel & Koste, 1985); River in Aust. (Bērziņš, 1982); 4 widely separated records, probably more widespread (Shiel & Koste, 1992); Westland, New Zealand (Russell, 1960)

- Neotropical region: Brazil, Surinam, Venezuela (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984)

Trichocerca intermedia (Stenroos), f. *arenosa* (Wiszniewski, 1953)

- Palaearctic region: E Eur. plain (Bērziņš, 1978)

Trichocerca jenningsi Voigt, 1957

- Palaearctic region: Lipowy pond, Poland (Lewkowicz, 1972); Romania (Rudescu, 1960); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Coast of S Finland (Eriksen, 1969); Almost all over Europe (Bērziņš, 1978); Inundation area of the Danube delta, Romania (Dorobantu, 1964); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Peat bog Chambedaze, Puy-de-Dôme, France (Francez, 1982); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); In reed borders of the Danube delta, Romania (Rudescu, 1968)

- Nearctic region: Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988)

- Oriental region: W & E Malaysia & Singapore (Fernando & Zankai, 1981); Malaysia (Sudzuki, 1989)

- Australian region: NT, Aust. (Koste & Shiel, 1980c); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); NT & Tas. Rare (Shiel & Koste, 1992)

- Neotropical region: Chile (Koste & José de Paggi, 1982); Chilean lakes (Schmid-Araya, 1991)

Trichocerca lata (Jennings, 1894)

- Palaeartic region: Coast of S Finland (Eriksen, 1969); ? NW Russian taiga (Bērziņš, 1978)

- Nearctic region: N America (Bérziņš, 1978); Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); Wisconsin, USA (Bérziņš, 1984); NW Canada (Chengalath & Koste, 1987); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); Columbia Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Adirondack Lakes, NY State, USA (Siegfried et al., 1984)

- Australian region: Westland, New Zealand (Russell, 1960)

Trichocerca longiseta (Schrank, 1802)

- Palaearctic region: Romania (Rudescu, 1960); La Mancha, Cast., Spain (Armengol *et al.*, 1975); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); N Swedish Lapland (Pejler, 1962a); C & S Sweden (Bérziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Lago Maggiore, Italy (Tonolli, 1962); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Achmer Grasmoor, NW Germany (Koste, 1968b); River Hase, NW Germany (Koste, 1976b); Fishponds near Warsaw, Poland (Hillbricht-Ilkowska, 1964); Peat bogs in the Massif Central, France (Batut, 1965); Lac Léman, Fr./CH (Balvay & Laurent, 1981 & 1990a); Volga river, R.F. (V'yushkova & Lakhnova, 1974; Mordukhai-Boltovskoi, 1979); Oligotrophic mesoacid lakes in the NW R.F. (Salazkin, 1971); Afghanistan (Bérziņš, 1961); Moor Skopun, Faröer (De Smet, 1988); Peat pits in Finland (Selin & Koskinen, 1988); Freshwater pools, S Finland (Björklund, 1972a); Coast of S Finland (Eriksen, 1969); Lakes in the Pyrenees, Spain (Miracle, 1981); River Loire, France (Lair, 1980); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Slovenia, ex-Yugoslavia (Markic, 1983); S Finland (Eloranta, 1982b); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Impoundment Kninicki, CSR (Kubicek, 1964); Lake Oerd, Ameland, & L. Van Hunenplak, Terschelling, NL (Leentvaar, 1967); Hilversumse wasmeer & lake on Leersumse veld, NL (Leentvaar, 1967b); Lake Quackjeswater, Voorne, NL (Leentvaar, 1963); Nat. Res. Neeruti Lakes, Estonia (Määmets A. & A., s.d.); Lake Iacub, Danube delta, Romania (Godeanu, 1976); Lac Luitel, Isère, France (Joly, 1960); Knowsley Park, Liverpool, UK (Galliford, 1964); Aral Sea spawning grounds, R.F. (Yerekeyev & Kazakhbayev, 1972); All over Europe; Caspian area. Cosmopolitan (Bérziņš, 1978); Japanese inland waters (Yamamoto, 1960a); Dozmary pool, Bodwin Moor, Cornwall, UK (Galliford, 1967); River Gracia basin, Poland (Pawlowski, 1960, 1968 & 1970); Lake Balaton, Hungary (Zankai, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b; Coussement, 1976); N, W & SW Iceland (De Ridder, 1966a, 1970 & 1972); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Brackish waters in the Lorraine orientale, France (De Ridder, 1969a); Overmere & Welle, Belgium (De Ridder, 1986); River Danube (Enaceanu & Brezeanu, 1970); Peat bog Chambedaze, Puy-de-Dôme, France (Francez, 1982); Peat bogs in the Auvergne,

France (Francez, 1981 & 1984c); Lake Biwa-ko, Japan (Yamamoto, 1960b); Pond Chylinski Maly II, Pod Padurka pond & Lipowy pond, Poland (Lewkowicz, 1972); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960); Nat. Res. Wellemeeersen, Denderleeuw, Belgium (Dumont *et al.*, 1975); Anatolia, Turkey (Segers *et al.*, 1992); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); In the lower Danube in Romania (Enaceanu, 1964); Mouth areas of the Danube branches, Romania (Popescu-Marinescu, 1974); Lakes in NE Norway (Saeter, 1971); Dübener Heide, N Germany (Wulfert, 1961); All over the Netherlands (Dresscher, 1976); Small river on Rhodos Island, Greece (De Ridder, 1967d); Karelian inland waters, R.F. (Kutikova, 1965); Danube delta, Romania (Popescu-Marinescu & Zinevici, 1967); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Lake Ösbysjön, near Stockholm, Sweden (Pejler, 1961); In pools, ponds & peat bogs in the Ile-de-France, France (Pourriot, 1970); Pool in the Fontainebleau forest, Ile-de-France, Fr. (Pourriot, 1976); Peat bogs in the Paris suburb, France (Pourriot, 1965); In reed borders of the Danube delta, Romania (Rudescu, 1968); Leżna-Włodawa Lake District, E Poland (Kowalczyk & Radwan, 1982); Rés. biol. de Villars-la-Dombes, France (Rostan, 1976); Hamburg area, Germany (Rühmann, 1961); Middle course of river Loire, France (Lair & Sargas, 1981); Fens in Genk, Belgium (Schepens, 1960); Pool, S of Trondheim, Norway (Thomasson, 1979); Lago Trasimeno, Italy (Taticchi, 1968); Serbian section of river Danube (Zivkovic, 1987); All over Spain (Velasco, 1990)

- Nearctic region: Walker Lake & Stoneman Lake, Arizona, USA (Green, 1986); Lake Jasper, W North America (Thomasson, 1962); Brainard Lake, N Colorado, USA (Thomasson, 1964b); Lake Ontario, Canada (Nauwerck, 1978); Columbia Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Colorado, USA (Turner, 1988); Mt. Tremblant Prov. Park, Québec, Canada (Nogrady, 1980); Kingston, Ontario, Canada (Nogrady, 1989); Hayden's Lake, Colorado, USA (Pennak, 1966); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Greens Lake, Beaver Island, Michigan, USA (Foran & King, 1982); Lake Wabamun, Alberta, Canada (Gallup & Hickman, 1975); Atchafalaya river basin, Louisiana, USA (Holland *et al.*, 1983); Long Lake, Washington, USA (Saltero *et al.*, 1975)

- Oriental region: River Yamuna, India (Rai, 1974); India (Sharma & Michael, 1980); Swamp near Mengala, Sumatra (Bērziņš, 1982a); Taiwan, Indonesia & India (Suzuki, 1989); Dal Lake, Kashmir Himalayan Lake, UP, India (Vass & Zutshi, 1983)

- Australian region: In many localities in Australia (Bērziņš, 1982); Tasmania (Koste & Shiel, 1987); NSW & Qld, Aust. (Shiel & Koste, 1979); NW Australia : Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Koste & Shiel, 1983a; Tait, Shiel & Koste, 1984); Widespread, not yet recorded from S Aust. & W Aust. Common (Shiel & Koste, 1992); Mt. Cook & Westland, New Zealand; Chatham Islands (Russell, 1960); New Zealand (Olivier, 1965)

- Neotropical region: ? River Parana, Argentina (Paggi & José de Paggi, 1974); Chile, Argentina, Paraguay, Venezuela & Panama (Koste & José de Paggi, 1982); Rapel reservoir, C Chile (Soto *et al.*, 1984); Chilean lakes (Schmid-Araya, 1991); Prov. Tierra del Fuego, Santa Cruz, Chubut, Rio Negro, Salta, Chaco & Santa Fé, Argentina (José de Paggi, 1990); Prov. Rio Negro, Santa Cruz & Tierra del Fuego, Argentina; Paraguay (Olivier, 1965); Araucarian lakes, W & E of the Andes, Chile (Thomasson, 1963b); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Ecuador : Oriente, Sierra & Costa (Koste & Böttger, 1992)

Trichocerca longistyla (Olofsson, 1918)

- Palaearctic region: Lake Donk, Overmere, Belgium (Coussement, 1976); N Swedish Lapland (Pejler, 1962a); N Eur. tundra (Bērziņš, 1978)

Trichocerca lophoessa (Gosse, 1886)

- Palaearctic region: Peat bog Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Serbian section of river Danube (Zivkovic, 1987); Romania (Rudescu, 1960); Coast of S Finland (Eriksen, 1969); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); All over Europe; Caspian area (Bērziņš, 1978); Peat bogs in the Auvergne, France (Francez, 1981); Lac d'Annecy, France (Balvay & Druart, 1992); In reed borders of the Danube delta, Romania (Rudescu, 1968)

- Nearctic region: Canada (Chengalath, 1977); Illinois, USA (Bērziņš, 1984); Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Lac Echo, Québec, Canada (Nogrady, 1976); Laurentian Great Lakes (Stemberger, 1979); Par pond, S Carolina, USA (Oden, 1979)

- Oriental region: Kalimantan (Borneo), Indonesia (Koste, 1988a); Indonesia (Suzuki, 1989)

- Australian region: Aaieri Beach, New Zealand (Russell, 1960)

- Neotropical region: Varzea near Santarem, Brazil (Koste, 1974b); Brazil (Koste & José de Paggi, 1982)

Trichocerca macera (Jennings, 1886)

- Palaearctic region: N Swedish Lapland (Pejler, 1962a); Romania (Rudescu, 1960); Lake Myvatn, Iceland (Adalsteinsson, 1979); Coast of S Finland (Eriksen, 1969); All over Europe (Bērziņš, 1978); Lake Balaton, Hungary (Zankai, 1968)
- Nearctic region: Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990)
- Australian region: NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); NT, Aust. (Shiel & Koste, 1985); Lake Catani, mountain Buffalo, Aust. (Bērziņš, 1982); Vic & Qld, Aust. (Shiel & Koste, 1985); Rare; NT, Qld & Vic (Shiel & Koste, 1992)

Trichocerca marina (Daday, 1890)

- Palaearctic region: Roscoff, Atlantic coast of Brittany, France (D'Hondt, 1970); Estuary of river Elbe, Germany (Giere, 1968); Lac Léman, Fr./CH (Balvay, 1989); Prov. Drente & Lake Veerse Meer, NL (Dresscher, 1976); Black Sea (Rudescu, 1961); In an inland sea of Japan, around Kasada Island (Suzuki, 1967a & 1976b); Spain : Pontevedra & Valencia (Velasco, 1990); Romania (Rudescu, 1960); In meso- & polyhaline parts of Lake Veerse Meer, NL (Bakker & De Pauw, 1975); Lake Palaeostomi, Georgia, R.F. (Haberman, 1982; Haberman & Laugaste, 1982); Sea & brackish waters in Norway (Eriksen, 1968); Coast of S Finland (Eriksen, 1969); Lake Cullera, near Valencia, Spain (Miracle & Vicente, 1893); N part of Caspian Sea & Lake Aral (Mordukhai-Boltovskoi, 1979); Almost all over Europe; Caucasus (Bērziņš, 1978); Rügener, Hiddenseer & Darszer Boddenwässer, Germany (Schwarz, 1963); Baltic Sea, Belt Sea, Skagerak, N & S North Sea, English Channel W, Bristol Channel & Irish Sea, S & W Ireland, NE Atlantic, Faroe-Shetland area, Faroe-Iceland area, Barentz Sea; Gulf of Riga, Gulf of Finland; Norwegian Sea (Bērziņš, 1960); North Sea in face of Nieuwpoort, Belgium (De Ridder, 1986); River Elbe estuary, Germany (Schulz, 1961); Akkeshi Bay, E District of Hokkaido, Japan (Hada, 1972)
- Nearctic region: St. Andrews Bay system, Florida, USA (Hopkins, 1966); Chesapeake Bay, Potomac river, USA (Heimbokel *et al.*, 1988); Placentia Bay & Conception Bay, Newfoundland, Canada (Davis, 1986)
- Australian region: Menzies Bay, New Zealand (Russell, 1960)

Trichocerca minuta (Olofsson, 1918)

- Palaearctic region: Prov. Zuid-Limburg, NL (Dresscher, 1976); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1975); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989c & 1989d); C Eur. plain, Iceland, Fennoscandia (Bērziņš, 1978); W & N Iceland (De Ridder, 1969b, 1970 & 1972)
- Nearctic region: Bayou de Siard, Monroe, Louisiana, USA (Gallagher, 1965)

Trichocerca mollis Edmondson, 1936

- Palaearctic region: Coast of S Finland (Eriksen, 1969); C Eur. plain, NW Russian taiga (Bērziņš, 1978)
- Nearctic region: N America (Bērziņš, 1978)

Trichocerca montana Hauer, 1956

- Palaearctic region: Peat bogs in the Puy-de-Dôme, France (Francez & Pourriot, 1984); Peat bogs Chambéry & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Southwest islands of Japan (Suzuki *et al.*, 1992)
- Australian region: NT, Australia (Shiel & Koste, 1986)
- Neotropical region: Crescent Lake, Rio Sna Missu, Mato Grosso, C Brazil (Green, 1972); Brokopondo Lake, Surinam (Leentvaar, 1979); Flooded savanna at Mantecal, Venezuela (Michelangeli *et al.*, 1979/1980)

Trichocerca mucosa (Stokes, 1896)

- Australian region: Early record from Qld, not seen in our material (Shiel & Koste, 1992)
- Neotropical region: Brokopondo Lake, Surinam (Leentvaar, 1979); Amazonia, Brazil (Thomasson, 1971); Lago do Castanho, C Amazonia, Brazil (Schaden, 1977b & 1977c); Lago Tefé & Lago Buiú, Amazonia, Brazil (Schaden, 1977c)

Trichocerca multicrinis (Kellicott, 1897)

- Nearctic region: Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Canada (Chengalath, 1977); Ontario, Canada (Chengalath, Fernando & Koste, 1971; Chengalath & Fernando, 1973); Lake Michigan & Milwaukee Harbor, USA (Stemberger, 1974); Northern Michigan inland lakes, USA (Stemberger, 1990); Ontario, Canada; Wisconsin & Kentucky, USA (Bērziņš, 1984); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Illinois river, USA (Brown *et al.*, 1989); Adirondack Lakes, NY State, USA (Siegfried *et al.*, 1984 & 1989); Colorado, USA (Turner, 1988); Acid lakes in Maine, USA (Brett, 1989); Lake Weir, Florida, USA (Beaver & Crisman, 1990); Lake O'Woods, Virginia, USA (Havens & Da Costa, 1987); Miller's Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamootttil, 1975); Newfoundland lakes, Canada (Chengalath *et al.*, 1984); Long Lake, Washington, USA (Soltero *et al.*, 1975); Experimental ponds, C Florida, USA (Fry & Osborne, 1980); Lake Wabamun, Alberta, Canada (Gallup & Hickman, 1975); Atchafalaya river basin, Louisiana, USA (Holland

et al., 1983); Triangle Lake, Portage County, Ohio, USA (Havens, 1989 & 1991); Laurentian Great Lakes (Stemberger, 1979); Clearwater Lake & Swan L., near Sudbury, Ontario, Canada (Yan & Geiling, 1985); Lake Ontario, Canada (Nauwerck, 1978); Bay of Quinte, Lake Ontario, Canada (Nogrady, 1988); Northern Ontario Lakes (Roff & Kwiatkowski, 1977); Dynamite Lake, Vermilion Co, Illinois, USA (Vanni, 1986); Silver Lake, Adirondack mts., NY State, USA (Siegfried, 1991)

- Oriental region: Dal Lake, Kashmir Himalayan Lake, India (Vass & Zutshi, 1983)
- Neotropical region: Panama (Koste & José de Paggi, 1982)

Trichocerca mus Hauer, 1938

- Palaearctic region: Leżna-Włodawa Lake District, E Poland. (Radwan, 1978); Southwest islands of Japan (Suzuki, 1992)
- Nearctic region: ? Mt. Tremblant Prov. Park, Québec, Canada (Nogrady, 1980)
- Oriental region: Kalimantan (Borneo), Indonesia (Koste, 1988a); Singapore & Indonesia (Suzuki, 1989); Lakes of Sunda Expedition (Serruya & Pollingher, 1984)
- Australian region: Vic, Aust. (Koste & Shiel, 1980c); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); NW Aust.: Magela Creek area (Koste, 1981b); NT, Tas & Vic. Rare (Shiel & Koste, 1992)
- Neotropical region: Lago do Macaco, Brazil (Koste, 1989a); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983)

Trichocerca musculus (Hauer, 1935)

- Palaearctic region: Baginice II pond, Poland (Lewkowicz, 1972); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Romania (Rudescu, 1960); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Coast of S Finland (Eriksen, 1969); Minorca, Spain (de Manuel, 1990); Achmer Grasmoor, NW Germany (Koste, 1968b); C Eur. mts., Carpathes, C & E Eur. plains, Baltic province, NW Russian taiga (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1970); Peat bogs in the Puy-de-Dôme, France (Francez & Pourriot, 1984); Peat bogs in the Auvergne, France (Francez, 1981); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960; Dresscher, 1976); River Bistritza & its tributaries, Romania (Miron, 1960); Shores of dammed or flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Peat bog Poiana Stampei, Bukowina, Romania (Rudescu, 1966); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)
- Nearctic region: N America (Bērziņš, 1978); Canada (Chengalath, 1977); Erbsville pond, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975)
- Australian region: ? SW Australia (Koste, Shiel & Brock, 1983); NSW & Southwest W Australia (Shiel & Koste, 1985); 2 records : mt. Koxinsko, NSW & Southwest W Australia (Shiel & Koste, 1992)

Trichocerca myersi (Hauer, 1931)

- Palaearctic region: Peat bog Poiana Stampei, Bukowina, Rom. (Rudescu, 1966); Etang de Coupe-Gorge, near Rambouillet, Yvelines, France (Pourriot, 1965); Pool, S of Trondheim, Norway (Thomasson, 1979); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); C & S Sweden (Bērziņš & Pejler, 1987, 1989a & 1989d); Alps, C Eur. mts., Carpathes, C & E Eur. plains (Bērziņš, 1978); Romania (Rudescu, 1960); River Grabia basin, Poland (Pawlowski, 1968 & 1970); Canals in Hran area, CSR (Terek, 1990a); Pelagic zone of Leżna-Włodawa lakes, Poland (Radwan, 1975)
- Nearctic region: N America (Bērziņš, 1978); Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); Cape Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Acid lakes in Maine, USA (Brett, 1989); Pinehurst Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975)
- Oriental region: India (Sharma & Michael, 1980); Malaysia, Indonesia & India (Suzuki, 1989); W & E Malaysia & Singapore (Fernando & Zankai, 1981); Storage reservoir Ajwa, near Baroda, India (Wulfert, 1966)
- Australian region: Tasmania (Koste & Shiel, 1987); NT, Aust. (Koste & Shiel, 1980c); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Jabiluka, N.T.; Lake St. Clair Nat. Park, Tas; Moorabool river, Vic. Rare (Shiel & Koste, 1992); Westland, New Zealand (Russell, 1960)
- Neotropical region: Ecuador : Oriente (Koste & Böttger, 1992); Venezuela (Koste & José de Paggi, 1982); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980); Amazonia, Brazil (Thomasson, 1971)

Trichocerca nitida Harring, 1914

- Palaearctic region: Leżna & Włodawa lake district, E Poland. (Radwan, 1971)
- Oriental region: ? Nimeta water works, near Baroda, India (Wulfert, 1966); ? Impoundment Kinda, C Burma (Koste & Tobias, 1990); India (Sharma & Michael, 1980; Suzuki, 1989)

- Neotropical region: Panama (Koste & José de Paggi, 1982); (Collado *et al.*, 1984)

Trichocerca obtusidens (Olofsson, 1918)

- Palaearctic region: C & E Eur. plain (Bērziņš, 1978)

- Neotropical region: Mato Grosso, C Brazil (Turner & Da Silva, 1992)

Trichocerca orca (Murray, 1913)

- Australian region: Müller Glacier, New Zealand (Russell, 1960)

Trichocerca ornata Myers, 1934

- Palaearctic region: C Eur. plains (Bērziņš, 1978)

- Nearctic region: N America (Bērziņš, 1978)

- Oriental region: Singapore (Sudzuki, 1989)

Trichocerca parva (Manfredi, 1927)

- Palaearctic region: Italy (Bērziņš, 1978)

Trichocerca parvula Carlin, 1939

- Palaearctic region: Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); ? Coast of S Finland (Eriksen, 1969); C Eur. mts., Carpathes, C & E Eur. plains, Baltic province, NW Russian taiga (Bērziņš, 1978); Romania (Rudescu, 1960); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Achmer Grasmoor, NW Germany (Koste, 1968b); Lapland State Preserve, R.F. (Ghilarov, 1972); Upper part of river Volga delta, R.F. (Chuykov, 1981); Canals in the Hran area, CSR (Terek, 1990c); Peat bogs in the Massif Central, C France (Francez & Pourriot, 1984); Peat bogs Chambédaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Peat bogs Chambédaze, Puy-de-Dôme, France (Francez, 1982); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Fens (acid water bodies) in the Netherlands (Dresscher, 1976); Lac Léman, Fr./CH (Martin, 1961; Balvay, 1989; Balvay & Laurent, 1981); Lake Constance, Germany (Kiefer, 1968); Thingvellir area, W Iceland (De Ridder, 1967a); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Peat bog Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Pool, S of Trondheim, Norway (Thomasson, 1979); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)

Trichocerca pediculus Remane, 1949

- Palaearctic region: C Eur. plain & Baltic province (Bērziņš, 1978)

Trichocerca plaka Myers, 1938

- Palaearctic region: C Eur. plains (Bērziņš, 1978)

- Nearctic region: N America (Bērziņš, 1978)

- Australian region: Lake Mc Gregor, New Zealand (Russell, 1960)

Trichocerca plateissa Myers, 1938

- Nearctic region: Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Ontario, Canada (Chengalath, Fernando & George, 1971; Chengalath & Fernando, 1973); Acid lakes in Maine, USA (Brett, 1989); Cranberry bog, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Newfoundland lakes, Canada (Chengalath *et al.*, 1984); Lac Echo, Québec, Canada (Nogrady, 1976); Mt. Tremblant Prov. Park, Québec, Canada (Nogrady, 1980); Northern Ontario Lakes, Canada (Roff & Kwiatkowski, 1977)

Trichocerca porcellus (Gosse, 1886)

- Palaearctic region: Lake Erken, Sweden (Nauwerck, 1963); Romania (Rudescu, 1960); Lake Myvatn, Iceland (Adalsteinsson, 1979); Lago di Giulianello, Latium, Italy (Bazzanti & Stella, 1977); Lake Bodensee, near Konstanz, Germany (Althaus, 1962); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Impoundments Säidenbach & Neunzehnhain, G. (Althaus, 1976); Lakes in S Sweden (Arnemo *et al.*, 1968); Lake Bielersee, CH (Baugerter *et al.*, 1980; Berner-Fankhauser, 1986); River Klyas'ma, R.F. (Bogoslawski, 1961); River Ybbarspå, Sweden (Bērziņš, 1979); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Afghanistan (Bērziņš, 1961); Lakes in Estonia (Haberman, 1978); Lake Ösbysjön, near Stockholm, Sweden (Pejler, 1961); Lakes in C Sweden (Pejler, 1964); River Moldova, tributary of the Danube, Rom. (Prunescu-Arion, 1968); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Lago Trasimeno, Italy (Taticchi, 1968); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Peat pits in Finland (Selin & Koskinen, 1988); Coast of S Finland (Eriksen, 1969); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Lake Glubokoe, near Moscow, R.F. (Matveeva, 1983, 1986 & 1991); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Buchseen, Bodensee (Einsle, 1966); Mindelsee, Bodensee (Einsle, 1967); C Finland (Eloranta, 1982b); Lake Mikolajskie & L. Sniardy, Mazurian lakeland, Poland (Kosicki, 1960); Lac Léman, Fr./CH (Naef & Martin, 1985; Balvay, 1989; Balvay & Laurent, 1989, 1990a & 1990b; Rougier, 1969); Lac de Nantua, Ain, France (Balvay, 1985); Bog pond near Budapest, Hungary (Nogrady, 1962); Lake Zürichsee, CH (Nipkow, 1961); N Swedish Lapland (Pejler, 1962a); Lakes C Sweden (Pejler, 1965); Lago Maggiore, Italy (Tonolli, 1962); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); Darnsee near Epe, NW Germany (Koste, 1962);

Achmer Grasmoor, NW Germany (Koste, 1968b); Azibo reservoir, near Porto, Portugal (Vascoucelos, 1990); Lake Mälaren, Sweden (Grönberg, 1975); Fish ponds near Warsaw, Poland (Hillbricht-Ilkowska, 1964); Lake Ladoga, R.F. (Nikolayev & Petrova, 1978); All over Europe, Caucasus, Caspian area, Asia Minor. Cosmopolitan (Bērziņš, 1978); Jordan reservoir, Bohemia, CSR (Mitiska *et al.*, 1962); Lower Silesia, Poland (Pawlowski, 1960); River Grabia basin, Poland (Pawlowski, 1968, 1970 & 1973); Lake Balaton, Hungary (Zankai, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b); W Iceland (De Ridder, 1969b & 1972); Peat bogs in the Puy-de-Dôme, C France (Francez & Pourriot, 1984); Peat bogs Chambéry & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Leżnica-Włodawa Lake District, E Poland (Kowalszyk & Radwan, 1982; Radwan, 1984); Japanese inland waters (Yamamoto, 1960a); Lake Constance, Germany (Walz *et al.*, 1987); Peat bog Chambéry, Puy-de-Dôme, France (Francez, 1982); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Lac Luitel, Isère, France (Joly, 1960); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960); Anatolia, Turkey (Segers *et al.*, 1992); Nat. Res. Meyendel; Zuid-Limburg; Nat. Res. Oisterwijkse vennen; Nat. Res. Het Hol; Nat. Res. De Gerritsflesch in Kootwijk; Zuid-Willemsvaart near Den Bosch, NL (Dresscher, 1976); Karelian inland waters, R.F. (Kutikova, 1965); Lac d'Annecy (Balvay & Druart, 1992); In reed borders of the Danube delta, Romania (Rudescu, 1968); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); 3 lakes of Sosnowickie, Poland (Radwan, 1969); Parczew area peat bogs, Poland (Radwan, 1974); Middle course of river Loire, France (Lair & Sargas, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990); Tarn in Värmland, Sweden (Thomasson, 1970); Pool, S of Trondheim, Sweden (Thomasson, 1979); Serbian section of river Danube (Zivkovic, 1987); River Salzach & its tributaries, Austria (Donner, 1970); Dübener Heide, Germany (Wulfert, 1961); Waters in the Szecezin lagoon, Poland (Wiktor, 1960); Spain : Alicante (Velasco, 1990); Lake Biwa-ko, Japan (Yamamoto, 1960b); Impoundment Twistetalsperre, Nordhessen, G. (Meinel *et al.*, 1983)

- Nearctic region: Lake Fernan, Idaho, USA (Buscemi, 1964); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Kingston, Ontario, Canada (Nogrady, 1989); Bay of Quinte, Ontario, Canada (Nogrady, 1988); Lake Michigan & Milwaukee Harbor, USA (Stemberger, 1974); Wisconsin, USA (Bērziņš, 1984); Ross Lake, Florida, USA (Bienert, 1986); Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Canada (Chengalath, 1977); Ontario, Canada (Chengalath, Fernando & George, 1971; Chengalath & Fernando, 1973); NE Canada (Chengalath & Koste, 1983); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); Pinehurst Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Adirondack Lakes, NY State, USA (Siegfried *et al.*, 1984); Lake Wabamun, Alberta, Canada (Gallup & Hickman, 1975); Atchafalaya river basin, Louisiana, USA (Holland *et al.*, 1983); Lac Echo, Québec, Canada (Nogrady, 1976); Mt. Tremblant Prov. Park, Québec, Canada (Nogrady, 1980); Crater L. & Walker L., Arizona, USA (Green, 1986); Silver Lake, Adirondack mts., NY State, USA (Siegfried, 1991)

- Oriental region: India (Sharma & Michael, 1980; Sudzuki, 1989); Lakelet Tan Daha, Kathmandu valley, Nepal (Daems & Dumont, 1974)

- Australian region: NW Aust.; Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); NSW & Qld, Aust. (Shiel & Koste, 1979); Creek & river in Aust. (Bērziņš, 1982); Swamp near Lake Hayes, S Island, New Zealand (Bērziņš, 1982a); Tasmania (Koste & Shiel, 1987); Lake Elingamite, freshwater Aust. crater lake (Green, 1981); Cosmopolitan. May be pancontinental, not yet known from S & W Australia. Common (Shiel & Koste, 1992); Sidney reservoirs, NSW, Aust. (Thomasson, 1973); Mt. Cook, Canterbury & Westland, New Zeal. (Russell, 1960)

- Neotropical region: Lake Llanquihue, Chile (Campos *et al.*, 1988); Ecuador : Costa (Koste & Böttger, 1989); Lago Paroni, Brazil (Koste, 1974a); Varzea near Santarem, Brazil (Koste, 1974b); Brazil & Panama (Koste & José de Paggi, 1982); Near Concepcion, Paraguay (Koste, 1986b); Chilean lakes (Schmid-Araya, 1991); Lake Todos los Santos, Chile (Campos *et al.*, 1990); Prov. Chubut, Rio Negro & Santa Fé, Argentina (José de Paggi, 1990); Amazonia, Brazil (Thomasson, 1971); Araucarian lakes, W & E of the Andes, Chile (Thomasson, 1963b); Mato Grosso, C Brazil (Turner & Da Silva, 1992)

Trichocerca porcellus (Gosse), f. ***major*** Hauer, 1935

- Palaearctic region: Lakes in Estonia (Haberman, 1978); Pool, S of Trondheim (Thomasson, 1979)

Trichocerca pusilla (Lauterborn, 1898)

- Palaearctic region: River Vistula in the Warsaw area, Poland (Klimowicz, 1981); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Romania (Rudescu, 1960); Loch Leven, Scotland (May, 1983, 1986 & 1987); Lake Nuwasawa, Japan (Sudzuki *et al.*, 1983); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Delta of river Danube, Romania (Godeanu & Zinevici, 1983; Prunescu-Arion, 1989; Zinevici & Teodorescu, 1990); Slovenia, ex-Yugoslavia (Markic, 1983); Lakes in Estonia (Mäemets, 1983);

C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Priest Pot, Lake District, UK (Hewitt & George, 1987); C Finland (Eloranta, 1982b); Man-made lake Zegrzynskie, Poland (Dusoge *et al.*, 1985); Lake Balaton, Hungary (Zankai & Kertesz, 1967; Zankai, 1968; Zankai & Ponyi, 1970, 1971a & 1973; Zankai, 1984 & 1989); Lake Biwa, Japan (Hillbricht-Ilkowska, 1983a); Lake Taltowisko, Mazurian lakeland, Poland (Klimowicz, 1972); Impoundment Kninicki, CSR (Kubicek, 1964); Lake Mikolajskie & L. Sniardny, Mazurian lakeland, Poland (Kosicki, 1960); Lakes in C Sweden (Pejler, 1965); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Impoundment Säidenbach, Germany (Althaus, 1976); Oligotrophic Lake Trummen, S Sweden (Andersson *et al.*, 1975); Groszer Müggelsee, Berlin, Germany (Barthelmes, 1960); ? Lake Bielersee, CH (Baugerter *et al.*, 1980); River Klyas'ma, R.F. (Bogoslawski, 1961); Impoundment "Eisernes Tor", Danube, Rom. (Brezeanu & Petén, 1973); Lake Bielersee, CH (Berner-Fankhauser, 1986); Middle course river Po, Italy (Ferrari *et al.*, 1986); Po river, Italy (Ferrari *et al.*, 1989); Etang de Virelles, Belgium (Moureau *et al.*, 1988); Grasmere Lake, Lake District, UK (Elliott, 1977); Spanish reservoirs (Guiset, 1977); Lakes in Estonia (Haberman, 1978); Coast of S Finland (Eriksen, 1969); Moerdijk-Hollands Diep, NL (Leentvaar, 1980); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Lac Léman, Fr./CH (Balvay & Laurent, 1981, 1989, 1990a & 1990b); Canal of river Odra, Poland (Klimowicz, 1979); River Loire, France (Lair *et al.*, 1978; Lair, 1980); Koyame-Ike, a coastal pond in Japan (Sudzuki & Fukuta, 1989); Minorca, Spain (de Manuel, 1990); Azibo reservoir, near Porto, Portugal (Vascoucelos, 1990); River Hase, NW Germany (Koste, 1976b); Lake Dümmer, NW Germany (Koste & Poltz, 1984); Lake Mälaren, Sweden (Grönberg, 1975; Blomquist & Pejler, 1980); Lake Alfsee, NW Germany (Koste & Poltz, 1987); Fish ponds near Warsaw, Poland (Hillbricht-Ilkowska, 1964); River Elbe near Hamburg, Germany (Caspers & Schulz, 1964); "Grote vijver", fort 8, Hoboken (Antwerp), B. (De Smet *et al.*, 1986); Moat of fort Merksem, Antwerp, Belgium (De Smet *et al.*, 1987); All over Europe, Caspian area, Asia Minor. Cosmopolitan (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1960 & 1970); Lake Volvi, Macedonia, Greece (Zarfdjian *et al.*, 1990); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b); Sava Lake, ex-Yugoslavia (Kalafatic, 1978); Yugoslavian section of the Danube river (Zivkovic, 1968); Lake Balaton, Hungary (Zankai & Ponyi, 1972); Leżnica-Włodawa Lake District, E Poland (Radwan, 1984); Japanese inland waters (Yamamoto, 1960a); Southwest islands of Japan (Sudzuki *et al.*, 1992); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Waldviertel area; lower Austria (Naidenow & Wawrik, 1984); River Danube (Enaceanu & Brezeanu, 1970); River Elbe estuary, Germany (Schulz, 1961); Lakes, N of Bucarest, Romania (Godeanu, 1969b); Lakes of the Black Sea shore, Romania (Godeanu, 1970b); Lake Gjersjøen, SE Norway (Hessen & Nilssen, 1985); Anatolia, Turkey (Segers *et al.*, 1992); All over the Netherlands (Dresscher, 1976); Lakes in the Leżnica-Włodawa area, E Poland (Radwan, 1976); Hamburg area, Germany (Rühmann, 1961); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Middle course of river Po, France (Lair & Sargos, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Ponds in the Paris suburb, France (Pourriot, 1965); Aegean lakes, Macedonia, ex-Yugoslavia-Greece (Serafimova-Hadzisce, 1974); Tarn in Värmland, Sweden (Thomasson, 1970); Lago Trasimeno, Italy (Taticchi, 1968); Serbian section of river Danube (Zivkovic, 1987); Lake Biwa, Japan (Yamamoto, 1967); Spain : Barcelona, Huesca (?), Valencia, Vizcaya (Velasco, 1990); Gravel pits along river Meuse, NL (Wibaut-Isebree Moens, 1964); Rybník dam reservoir, Poland (Krzanowski, 1979); Lake Lauerzersee, CH (Odermatt, 1970); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); In the lower Danube in Romania (Enaceanu, 1964); Mouth areas of the Danube branches, Rom. & predeltaic area of the Black Sea (Popescu-Marinescu, 1974); Lake Erken, Sweden (Nauwerck, 1963); Karelian inland waters, R.F. (Kutikova, 1965); Danube delta, Romania (Popescu-Marinescu & Zinevici, 1967); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Lakes in C Sweden (Pejler, 1964); In reed borders of the Danube delta, Romania (Rudescu, 1968); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); Ponds in Oltina, SE Dobrogea, Romania (Popescu-Gorj & Costea, 1961); River Loire in St. Laurent-les-Eaux, France (Pourriot *et al.*, 1982); Parczew area peat bogs, Poland (Radwan, 1974)

- Nearctic region: San Francisco, USA (Nogrady, 1983); Lake Ontario, Canada (Nauwerck, 1978); Arctic N America (Chengalath & Koste, 1989); Bay of Quinte, Lake Ontario, Canada (Nogrady, 1988); Kingston, Ontario, Canada (Nogrady, 1989); Canada (Chengalath, 1977); Wisconsin, Illinois & Kentucky, USA (Bērziņš, 1984); Lakes Adirondack, NY State, USA (Siegfried *et al.*, 1984 & 1989); Northern Michigan inland lakes (Stemberger, 1990); Lakes in Santée, California, USA (Nogrady, 1982); Laurentian Great Lakes (Stemberger, 1979); Bayou de Siard, Monroe, Louisiana, USA (Gallagher, 1965)

- Oriental region: Bangkok, Thailand (De Ridder, 1971); Thailand, Malaysia & Singapore (Sudzuki, 1989); Kalimantan (Borneo), Indonesia (Koste, 1988a); Delhi, India (Sarma, 1988); Singapore. Cosmopolitan (Sudzuki, 1991); Thailand (Boonsom, 1984b); Malaysia & Singapore (Fernando & Zankai, 1981); Impoundment Kinda, C Burma (Koste & Tobias, 1990)

- Australian region: SW Australia (Koste, Shiel & Brock, 1983); Tasmania (Koste & Shiel, 1987); River

Yarra, Aust. (Bērziņš, 1982); W Aust. (Shiel & Koste, 1979); Lower river Murray, S Aust. (Shiel, Walker & Williams, 1982); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, Aust. (Koste & Shiel, 1983a; Tait, Shiel & Koste, 1984); E Aust. (Koste & Shiel, 1980b); Pancontinental, common in billabongs (Shiel & Koste, 1992)

- Neotropical region: Brokopondo Lake, Surinam (Leentvaar, 1979); Argentina (Modenutti & Claps, 1988); Orinoco river basin, Venezuela (Vasquez, 1984a); Caroni river basin, Venezuela (Vasquez, 1984b); Lower Orinoco river & its delta, Venezuela (Vasquez & Rey, 1989); Lago Paroni, Brazil (Koste, 1974a); Amazonia, Brazil (Hauer, 1965b; Thomasson, 1971; Koste, 1972f); Brazil, Chile, Antilles, Venezuela, Surinam & Panama (Koste & José de Paggi, 1982); Ecuador : Oriente & Costa (Koste & Böttger, 1989); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Concepcion, Paraguay (Koste, 1986b); Chilean lakes (Schmid-Araya, 1991); Prov. Neuquen, Argentina (José de Paggi, 1990); Caribbean Islands (Collado *et al.*, 1984); Lake Monte Alegre, SE Brazil (Arcifa *et al.*, 1992); Lago de Valencia, Venezuela (De Infante, 1980 & 1982; Saunders & Lewis, 1988); Antilles françaises (Pourriot, 1975); Araucarian lakes, W & E of the Andes, Chile (Thomasson, 1963b); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Rio Tocantins, Lago Jurucui & Lago Moratu, Amazonia, Brazil (Schaden, 1977c); Lake Cocibolca, L. Masaya & L. Tiscapa, Nicaragua (Moreno *et al.*, 1992)

***Trichocerca pusilla* (Lauterborn), f. *grandis* Koste, 1989**

- Neotropical region: Lago de Macaco, Brazil (Koste, 1989a)

***Trichocerca pygocera* (Wiszniewski, 1932)**

- Palaearctic region: Italy, C & E Eur. plains (Bērziņš, 1978)

***Trichocerca ratus* (O.F. Müller, 1776)**

- Palaearctic region: Romania (Rudescu, 1960); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Bjørnøya (Bear Island), Svalbard (De Smet, 1988); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); Alpine lakes Sierra Nevada, Spain (Morales-Baquero *et al.*, 1989); Minorca, Spain (de Manuel, 1990); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Lake Myvatn, Iceland (Adalsteinsson, 1979); Lago di Giulianello, Latium, Italy (Bazzanti & Stella, 1977); Lake Bodensee, near Konstanz, Germany (Althaus, 1962); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); River Klyas'ma, R.F. (Bogoslowski, 1961); Lake Balaton, Hungary (Zankai & Kertesz, 1967; Zankai, 1968; Zankai & Ponyi, 1971a); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Afghanistan (Bērziņš, 1971); Lakes in Estonia (Haberman, 1978); Sea, brackish & freshwater pools in S Finland (Björklund, 1972a); Coast of S Finland (Eriksen, 1969); Lake Võrtsjärv, Estonia (Kutikova & Haberman, 1986); Delta of river Danube, Romania (Godeanu & Zinevici, 1983; Prunescu-Arion, 1989); Slovenia, ex-Yugoslavia (Markic, 1983); Sierra Nevada, Spain (Morales-Baquero, 1987); District Ostrava, CSR (Dvorakova, 1960b); Hortobagy Nat. Park, Hungary (Forro & Ronkay, 1983); Man-made Lake Zegrzynskie, Poland (Dusoge *et al.*, 1985); Nat. Res. Jursky Sur, CSR (Hudec, 1978); Near Asperg, Württemberg, Germany (Klement, 1960); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Alkal. Péteri-See, Hungary (Kertesz, 1960); Broads & Peat pits in NW Overijssel, NL (Schroevers, 1965); Achmer Grasmoor, NW Germany (Koste, 1968b); River Hase, NW Germany (Koste, 1976b); Lake Dümmer, NW Germany (Koste & Poltz, 1984); Fish ponds near Warsaw, Poland (Hillbricht-Ilkowska, 1964); Volga river, R.F. (V'yushkova & Lakhnova, 1974); All over Europe, Caucasus, Caspian area, Asia Minor. Cosmopolitan (Bērziņš, 1978); Japanese inland waters (Yamamoto, 1960a); Inundation area of Danube delta, Romania (Dorobantu, 1964); River Gracia basin, Poland (Pawlowski, 1968 & 1970); Latorica, E Slovakia lowland, CSR (Terek, 1990a); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b); Camargue, S France (De Ridder, 1960a & 1961); N, W & C Iceland (De Ridder, 1966a, 1969b, 1970 & 1972); Sava Lake, ex-Yugoslavia (Kalafatic, 1978); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Lake Neusiedlersee, Austria (Donner, 1979); Brackish waters near Bruges, Belgium (De Ridder & Verheyen, 1980); Brugge & Welle, Belgium (De Ridder, 1986); Lake Constance, Germany (Walz *et al.*, 1987); River Danube (Enaceanu & Brezeanu, 1970); Peat bogs in the Auvergne, France (Francez, 1981); Water-courses in the Bucegi mts., Romania (Godeanu, 1963b); Lakes N of Bucarest, Romania (Godeanu, 1969b); Lakes of the Black Sea shore, Romania (Godeanu, 1970b); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Temporary waters, small permanent ponds & marshy zones in the Bucegi mts., Romania (Godeanu, 1969a); Nat. Res. Wellemeersen, Belgium (Dumont *et al.*, 1975); Near water works at Okocim, Poland (Klimowicz, 1975); Sewage ponds near Czestochowa, Poland (Klimowicz, 1968); Small water bodies in the Mikolajki area, Poland (Klimowicz, 1967); Pond in the "Forêt de Sénaart", near Paris, France (Chément, 1975); Anatolia, Turkey (Segers *et al.*, 1992); All over the Netherlands (Dresscher, 1976); Dübenner Heide, N Germany (Wulfert, 1960a & 1960b); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Lake Victoria, Oltena area, Romania (Marx *et al.*, 1967); Luga district of the

Leningrad area, R.F. (Kutikova, 1962b); In many acid & alkaline waters of the Ile-de-France, France (Pourriot, 1970); Black Sea (Rudescu, 1961); In reed borders of the Danube delta, Romania (Rudescu, 1968); ? Shores of dammed and free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); Ponds in Oltina, SE Dobrogea, Romania (Popescu-Gorj & Costea, 1961); Rés. biol. de Villars-la-Dombes, France (Rostan, 1976); Lake Chenhu, Changjiang basin, China (Weu, 1992); Hamburg area, Germany (Rühmann, 1961); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Pond in the Paris suburb, France (Pourriot, 1965); Ponds in the Botanical Garden, Gent, Belgium (Segers *et al.*, 1991b); Periphytic biota in sewages & park waters, Japan (Suzuki, 1981); Spain : Granada, Madrid, Orense, Oviedo & Valencia (Velasco, 1990); Lake Biwa-ko, Japan (Yamamoto, 1960b)

- Nearctic region: NE Canada (Chengalath & Koste, 1983); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Colorado, USA (Turner, 1988); Bay of Quinte, Ontario, Canada (Nogrady, 1988); Kingston, Ontario, Canada (Nogrady, 1989); Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Canada (Chengalath, 1977); Parvin Lake, Colorado (Pennak, 1966); Kentucky, USA (Bērziņš, 1984); Laurel Creek reservoir, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Laurentian Great Lakes (Stemberger, 1979)

- Oriental region: Sri Lanka (Mendis & Fernando, 1962; Fernando, 1969 & 1980a); Kalimantan (Borneo), Indonesia (Koste, 1988a); India (Sharma & Michael, 1980); Philippines (Mamaril & Fernando, 1978); Malaysia, Sri Lanka, India & Philippines (Suzuki, 1989); Sri Lanka. Common in lakes & ponds (Chengalath, Fernando & Koste, 1973); W & E Malaysia & Singapore (Fernando & Zankai, 1981); 2 Kumaun Himalayan lakes, UP, India (Sharma & Pant, 1984 & 1985); W Bengal, NE India (Sharma, 1979b); Lake Manguao, Palawan, the Philippines (Davies & Green, 1990); Nimeta water works, near Baroda, India (Wulfert, 1966)

- Australian region: River & Creek in Australia (Bērziņš, 1982); NSW, Qld & Vic, Aust. (Shiel & Koste, 1979); Tasmania (Koste & Shiel, 1987); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Common, widespread in Australia & Tasmania (Shiel & Koste, 1992); Pukeran & Christchurch, New Zealand (Russell, 1960)

- Neotropical region: Ecuador : Oriente & Costa (Koste & Böttger, 1989 & 1992); Araucarian lakes, E of the Andes, Chile (Thomasson, 1963b); Antilles françaises (Pourriot, 1975); Venezuela, Brazil, Argentina, Antilles, Panama (Koste & José de Paggi, 1982); Varzea near Santarem, Brazil (Koste, 1974b); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Galapagos Islands (De Smet, 1987 & 1989); River Parana, Argentina (Paggi & José de Paggi, 1974; José de Paggi, 1985 & 1987a); Eutrophic water Santa Fé, Argentina (José de Paggi, 1976a); ? River Santa Fé, Argentina (José de Paggi, 1976b); River Saladillo basin, Argentina (José de Paggi & Koste, 1988); Argentina (Modenutti & Claps, 1988); Provinces in NW Argentina (José de Paggi, 1989); Prov. Rio Negro, Tucuman, Santiago del Estero, Chaco, Santa Fé & Buenos Aires, Argentina (José de Paggi, 1990); Caribbean Islands (Collado *et al.*, 1984); ? Brokopondo Lake, Surinam (Leentvaar, 1979); Lower course of river Uruguay, Argentina (Di Persia & Neiff, 1986); River Parana, Brazil (Bonetto, 1986); Flooded savanna at Mantecal, Venezuela (Michelangeli *et al.*, 1979/1980)

Trichocerca rattus (O.F. Müller), f. *carinata* (Ehrenberg, 1830)

- Palaearctic region: 3 lakes of Sosnowickie, Poland (Radwan, 1969); Lago Trasimeno, Italy (Taticchi, 1968); Spain : Menorca (Baleares) (Velasco, 1990); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Romania (Rudescu, 1960); River Klyas'ma, R.F. (Bogoslawski, 1961); Coast of S Finland (Eriksen, 1969); Lake Bodensee, near Konstanz, Germany (Althaus, 1962); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989c & 1989d); District Ostrava, CSR (Dvorakova, 1960b); Near Asperg, Württemberg, Germany (Klement, 1960); Lake Palaeostomi, Georgia, R.F. (Haberland & Laugaste, 1982); N Swedish Lapland (Pejler, 1962a); Minorca, Spain (de Manuel, 1990); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Lake Alfsee, NW Germany (Koste & Poltz, 1987); Fish ponds near Warsaw, Poland (Hillbright-Illkowska, 1964); Volga river, R.F. (V'yushkova & Lakhnova, 1974); All over Europe, Caucasus, Caspian area (Bērziņš, 1978); Moat of fort Merksem, Antwerp, Belgium (De Smet *et al.*, 1987); Impoundments in E Bohemia, CSR (Sladeckowa & Sladecek, 1960); Lake Donk, Overmere, Belgium (Dumont, 1968b); Canal Leeds-Liverpool, UK (Galliford, 1960b); Knowsley Park, Liverpool, UK (Galliford, 1964); Side-arm of r. Danube near Bratislava, CSR (Lichardova, 1964); Anatolia, Turkey (Segers *et al.*, 1992); River Bistrizta & its tributaries, Romania (Miron, 1960); S Vecht lakes & Old branch cut off r. Waal; lakes W & E of Amsterdam; peat holes in NW Overijssel; dune lakes on Island of Voorne, NL (Dresscher, 1976); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Karelian inland waters, R.F. (Kutikova, 1965); Luga district of the

Leningrad area, R.F. (Kutikova, 1962b); Pool in Fontainebleau Forest, Ile-de-France, France (Pourriot, 1976); Parczew area peat bogs, Poland (Radwan, 1974); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990)

- Nearctic region: Kingston, Ontario, Canada (Nogrady, 1989); NE Canada (Chengalath & Koste, 1983); Stoneman Lake, Arizona, USA (Green, 1986)
- Oriental region: Singapore (Sudzuki, 1989); Sri Lanka (Mendis & Fernando, 1962); Lakelet Tan Daha, Kathmandu valley, Nepal (Daems & Dumont, 1974)
- Australian region: Lake Euramo, tropical Queensland crater lake (Green, 1981); 2 Creeks in Aust. (Bērziņš, 1982); Vic, Aust. (Shiel & Koste, 1979 & 1985); Tasmania (Koste & Shiel, 1986b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Lake Ellesmere, New Zealand (Russell, 1960)
- Neotropical region: Paraguay (Koste & José de Paggi, 1982); Lake Coca-Cola, near Rio de Janeiro, Brazil (Turner, 1990); Chilean lakes (Schmid-Araya, 1991)

***Trichocerca rutilus* (O.F. Müller), f. *globosa* Dartnall & Hollowday, 1985**

- Antarctic region: Antarctica (Dartnall & Hollowday, 1985)

***Trichocerca rutilus* (O.F. Müller), f. *minor* Fadeev, 1925**

- Palaearctic region: Carpathes, C & E Eur. plains (Bērziņš, 1978); Anatolia, Turkey (Segers *et al.*, 1992); In reed borders of the Danube delta, Romania (Rudescu, 1968); Serbian section of river Danube (Zivkovic, 1987)

***Trichocerca relicta* Donner, 1950**

- Palaearctic region: Romania (Rudescu, 1960); Lake Bodensee, near Konstanz, Germany (Althaus, 1962); Sierra Nevada, Spain (Morales-Baquero, 1987); Alpine lakes, Sierra Nevada, Spain (Morales-Baquero *et al.*, 1989); Bjørnøya (Bear Island), Svalbard (De Smet, 1988); ? Lake Dümmer, NW Germany (Koste & Poltz, 1984); Lake Alfssee, NW Germany (Koste & Poltz, 1987); Alps, C Eur. mts., Pontic province, C Eur. plain & Baltic province (Bērziņš, 1978); Anatolia, Turkey (Segers *et al.*, 1992); Dübener Heide, N Germany (Wulfert, 1961); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); River Salzach & its tributaries, Austria (Donner, 1970); Spain : Granada (Velasco, 1990)
- Nearctic region: Canada (Chengalath, 1977); Illinois, USA (Bērziņš, 1984); Northern Michigan inland lakes (Stemberger, 1990); Cranberry bog, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975)
- Neotropical region: Galapagos Islands (De Smet, 1987 & 1989); Santa Cruz, Galapagos Islands (Segers, 1991a)

***Trichocerca ripli* Bērziņš, 1973**

- Australian region: New Zealand (Bērziņš, 1973)

***Trichocerca rosea* (Stenoos, 1898)**

- Palaearctic region: Small lakes in Estonia (Haberman & Mäemets, 1987); Romania (Rudescu, 1960); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Coast of S Finland (Eriksen, 1969); Nat. Res. Neeruti Lakes, NE Estonia (Mäemets, A. & A., s.d.); Small water bodies in Soviet Karelia, R.F. (Filimonova, 1976); C Eur. mts., Carpathes, C & E Eur. plains, Baltic province, N Sweden, NW Russian taiga (Bērziņš, 1978); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960); Ibid. + Maria-Helenapeel, NL (Dresscher, 1976); Karelian inland waters, R.F. (Kutikova, 1965); Pelagic zone of the Leżna-Włodawa lakes, P. (Radwan, 1975); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966)
- Nearctic region: Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Little Swede n° 1 lake, Colorado, USA (Pennak, 1966); Colorado, USA (Turner, 1988); N America (Bērziņš, 1978); Northern Michigan inland lakes (Stemberger, 1990); ? Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980)
- Australian region: S Tasmania (Koste, Shiel & Tan, 1988); Single record from Tas. (Shiel & Koste, 1992); Westland, New Zealand (Russell, 1960)
- Neotropical region: Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Ecuador : Oriente (Koste & Böttger, 1992)

***Trichocerca rotundata* Myers, 1937**

- Palaearctic region: ? Afghanistan (Bērziņš, 1961)
- Nearctic region: Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980)

***Trichocerca rousseleti* (Voigt, 1902)**

- Palaearctic region: River Moldova, tributary of the Danube, Rom. (Prunescu-Arion, 1968); Lakes in C Sweden (Pejler, 1964); Pelagic zone of the Leżna-Włodawa lakes, P. (Radwan, 1975); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Spain : Lerida (Velasco, 1990); Romania (Rudescu, 1960); Lake Vico, C Italy (Ferrari, 1972); Man-made lake Zegrzynskie, Poland (Dusoge *et al.*, 1985); Lake Balaton, Hungary (Zankai & Ponyi, 1970 & 1971a; Zankai, 1989); Lago Maggiore, Italy (Tonolli, 1962; Goldman *et al.*,

1968); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Lake Stechlinsee & L. Nehmitzsee, Brandenburg, Germany (Kasprzak & Ronneberger, 1982); Impoundment Kninicki, CSR (Kubicek, 1964); Lakes in C Sweden (Pejler, 1965); Lake Sempachersee, CH (Zimmermann, 1974); Darnsee near Epe, NW Germany (Koste, 1962); Lake Mälaren, Sweden (Grönberg, 1975); Lake Alfsee, NW Germany (Koste & Poltz, 1987); Lac Léman, Fr./CH (Martin, 1961; Naef & Martin, 1985;; Balvay, 1989; Balvay & Laurent, 1981, 1989, 1990a & 1990b); Karelian inland waters, R.F. (Kutikova, 1965); Lac de Nantua, Ain, France (Balvay, 1985); Lake Kinselmeer, Amsterdam; Lake Barlosche Kolk near Wijlre, Drente; Lake Ysselmeer, NL (Dresscher, 1978); Volgograd reservoir, R.F. (V'yushkova & Lakhnova, 1974); Lake Ladoga, R.F. (Nikolayev & Petrova, 1978); Lake Zürichsee, CH (Nipkov, 1961); All over Europe (Bērziņš, 1978); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Impoundment Saidenbach, Erzgebirge, Germany (Beuschold, 1961; Rudolf, 1966); Lakes in S Sweden (Arnemo *et al.*, 1968); Small lakes in Estonia (Haberman & Mäemets, 1987); Lake Bielersee, CH (Berner-Fankhauser, 19865 & 1987); Coast of S Finland (Eriksen, 1969); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Lake Krasnoye, Karelian isthmus, R.F. (Alexandrov & Kutikova, 1985); Near Lappeenranta, Finland (Järnefelt, 1961); Leżna-Włodawa Lakes district, Poland (Radwan, 1984); Southwest islands of Japan (Sudzuki *et al.*, 1992); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Vlimmeren, Belgium (De Ridder, 1986); Lake Constance, Germany (Kiefer, 1968; Walz *et al.*, 1987); Lake Lauerzersee, CH (Odermatt, 1970)

- Nearctic region: N America (Bērziņš, 1978); Lake 382, exper. lake area, NW Ontario, Canada (Malley & Chang, 1991); Canada (Chengalath, 1977); Lake Michigan & Milwaukee Harbor, USA (Stemberger, 1974); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); Wisconsin, Kentucky & Illinois, USA (Bērziņš, 1984); Arctic N America (Chengalath & Koste, 1989); Lake Oglethorpe, NE Georgia, USA (Pace & Orcutt, 1981; Orcutt & Pace, 1984); Bayou de Siard, Monroe, Louisiana, USA (Gallagher, 1965); Bay of Quinte, Lake Ontario, Canada (Nogrady, 1988); Singletary Lake, N Carolina, USA (Thomasson, 1963a); ? Lake 122, NW Ontario, Canada (Schindler & Novén, 1971)

- Oriental region: Indonesia (Sudzuki, 1989); Kalimantan (Borneo), Indonesia (Koste, 1988a)

- Australian region: S Australia (Shiel & Koste, 1979); Lower river Murray, S Aust. (Shiel, Walker & Williams, 1982); Creek & river in Aust. (Bērziņš, 1982); Rare, in our collections invariably associated with flowing waters - Darling (NSW), Goulburn (Vic), Murray (SA) - with *Melosira* - blooms (Shiel & Koste, 1992); Rio Negro, Amazonia, Brazil (Schaden, 1977c)

Trichocerca ruttneri (Donner, 1953)

- Palaearctic region: Serbian section of river Danube (Zivkovic, 1987); ? Impoundment in Spain (Velasco, 1990); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); C Eur. mts. & Hungarian plain (Bērziņš, 1978); Austria (Koste, 1972f); Southwest islands of Japan (Sudzuki, 1990; Sudzuki *et al.*, 1992); Lake Neusiedlersee, Austria (Donner, 1979); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978)

- Nearctic region: Canada (Chengalath, 1977); Pinehurst Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975)

- Oriental region: Storage reservoir Ajwa, near Baroda, India (Wulfert, 1966); Punjab State, NW India (Sharma, 1976); India (Koste, 1972f; Sharma & Michael, 1980); Taiwan (Sudzuki, 1988); Taiwan & India (Sudzuki, 1989); Sumatra (Koste, 1972f); Ajwa reservoir, near Baroda, Gujarat, India (Jayangaudar, 1980); Lake Manguao, Palawan, the Philippines (Davies & Green, 1990)

- Australian region: Qld, Aust. (Shiel & Koste, 1985); Known only from dams near Chillagoe, Qld (Shiel & Koste, 1992)

- Neotropical region: Chile (Löffler, 1962); Lago Jurucui, Amazonia, Brazil (Schaden, 1977c); Lago Rotondo, Amazonia, Brazil (Serruya & Pollingher, 1984); Araucarian lakes, W of the Andes (Thomasson, 1963b); Amazonia, Brazil (Hauer, 1965b; Thomasson, 1971;; Koste, 1972f); Brazil (Koste, 1972f); Lower Orinoco river & its delta, Venezuela (Vasquez & Rey, 1989); Brazil, Chile & Honduras, Venezuela (Koste & José de Paggi, 1982); Chilean lakes (Schmid-Araya, 1991); Lake Escondido, prov. Rio Negro, Argentina (Balseiro & Modenutti, 1990); Prov. Chubut, Argentina (José de Paggi, 1990); Lago de Valencia, Venezuela (De Infante, 1980)

Trichocerca scipio Gosse, 1886

- Palaearctic region: Coast of S Finland (Eriksen, 1969); Romania (Rudescu, 1960); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); N Swedish Lapland (Pejler, 1962a); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); Almost all over Europe (Bērziņš, 1978); Japanese inland waters (Yamamoto, 1960a); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Karelian inland waters, R.F. (Kutikova, 1965); In reed borders of the Danube delta, Rom. (Rudescu, 1968); Lipowy pond, Poland (Lewkowicz, 1972)

- Nearctic region: Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Laurentian Great Lakes (Stemberger, 1979)

- Oriental region: India (Sharma & Michael, 1980; Sudzuki, 1989); Sri Lanka (Mendis & Fernando, 1962; Fernando, 1969)
- Australian region: S Tasmania (Koste, Shiel & Tan, 1988); Recorded from 3 localities in Tasmania (Shiel & Koste, 1992); Mt. Cook, New Zealand & Campbell Island (Russell, 1960)
- Neotropical region: Ecuador : Oriente (Koste & Böttger, 1989); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990)

***Trichocerca sejunctipes* (Gosse, 1886)**

- Palaearctic region: Coast of S Finland (Eriksen, 1969); Peat bogs in the Massif Central, France (Batut, 1965); Romania (Rudescu, 1960); All over Europe (Bērziņš, 1978); River Bistritza & its tributaries, Romania (Miron, 1960); Prov. Zuid-Limburg, NL (Dresscher, 1976); Sinaia, S Carpathes, Romania (Rodewald-Rudescu & Godeanu, 1961); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966)

***Trichocerca similis* (Wiersejski, 1893)**

- Palaearctic region: Romania (Rudescu, 1960); Impoundment on river Guadiana, C Spain (Armengol, 1982); Small lakes in Estonia (Haberman & Mäemets, 1987); Lake Bielersee, CH (Berner-Fankhauser, 1986); Middle course of river Po, Italy (Ferrari *et al.*, 1986); Po river, Italy (Ferrari *et al.*, 1989); River Ybbarspå, Sweden (Bērziņš, 1979); Grasmere Lake, Lake District, UK (Elliott, 1977); Banyoles Lake, NE Spain (Miracle & Vicente, 1983; Miracle, 1974 & 1977); Lakes in Estonia (Haberman, 1978; Mäemets, 1983); Asiatic Turkey (Dumont & De Ridder, 1987); C & S Sweden (Bērziņš & Pjeler, 1987, 1989a, 1989b, 1989c & 1989d); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Lake Mikolajskie & L. Taltowisko, Poland (Klimowicz, 1972); Nat. Res. Neeruti Lakes, Estonia (Mäemets A. & A., s.d.); Lakes in C Sweden (Pejler, 1964 & 1965); Lac Léman, Fr./CH (Balvay, 1989; Balvay & Laurent, 1981, 1989, 1990a & 1990b); Loch Rusky, Scotland (Romo, 1990); Volga river, R.F. (V'yushkova & Lakhnova, 1974; Mordukhai-Boltovskoi, 1979); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); River Hase, NW Germany (Koste, 1976b); Lake Dümmer, NW Germany (Koste & Poltz, 1984); ? Lake Volvi, Macedonia, Greece (Zarfdjian *et al.*, 1990); River Grabia basin, Poland (Pawlowski, 1968 & 1970); Lake Balaton, Hungary (Zankai, 1968); Etang du Brochet, Rambouillet, France (Pourriot *et al.*, 1969); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Quéntar reservoir, Sierra Nevada, near Granada, Spain (Leighton, 1981); Southwest islands of Japan (Sudzuki *et al.*, 1992); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Lacuisine, Belgium (De Maeseneer, 1979; De Ridder, 1986); River Danube (Enaceanu & Brezeanu, 1970); Knowsley Park, Liverpool, UK (Galliford, 1964); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Oisterwijkse vennen, NL (De Graaf, 1960); New impoundments in the CSR (Kubicek, 1968); Side-arm of river Danube near Bratislava, CSR (Lichardova, 1964); Anatolia, Turkey (Segers *et al.*, 1992); Shores of dammed or free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Lake Barlosche Kolk, near Wijlre, Drente; Nat. Res. de Oisterwijkse vennen; Lake de Venematen; Maria- & Helenapeel; pool "Vechten" near Utrecht, NL (Dresscher, 1976); Lac d'Annecy, France (Balvay & Druart, 1992); Rés. biol. de Villars-la-Dombes, France (Rostan, 1976); 3 lakes of Sosnowickie, Poland (Radwan, 1969); Parczew area peat bogs, Poland (Radwan, 1974); Gemündener Maar, Weinfelder Maar & Pulvermaar, Eifel, Germany (Oehms & Seitz, 1992); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Middle course of river Loire, France (Lair & Sargos, 1981); ? Lago Trasimeno, Italy (Taticchi, 1968); Ponds in the Paris suburb, France (Pourriot, 1965); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990); Serbian section of river Danube (Zivkovic, 1987); Spain : Badajos, Cuenca, Gerona, Huelva, Léon & Sevilla (Velasco, 1990); C Eur. mts. Carpathes, C & E Eur. plains, Baltic prov., Ireland, UK, N Sweden, NW Russian taiga (Bērziņš, 1978); "Grote vijver", fort 8, Hoboken (Antwerp), B. (De Smet *et al.*, 1986); "Kleine vijver", fort 8, Hoboken (Antwerp), B. (De Smet *et al.*, 1989); Azibo reservoir, near Porto, Portugal (Vascoucelos, 1990); Lake Glubokoe, near Moscow, R.F. (Matveeva, 1986 & 1991); Leżnica & Włodawa Lake district, E Poland (Kowalczyk & Radwan, 1982); (Radwan, 1984); Lake Aydat & L. Bourdouze, Puy-de-Dôme, Fr. (Persoone *et al.*, 1968)

- Nearctic region: Crater Lake, Arizona, USA (Green, 1986); Singleton Lake, N Carolina, USA (Thomasson, 1963a); Massey Lake, 16 km of Beaumont, Texas, USA (Marsh *et al.*, 1978); Acid lakes in Maine, USA (Brett, 1989); NE Canada (Chengalath & Koste, 1983); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Everglades Nat Park, Florida, USA (Nogrady, 1983); Bay of Quinte, Lake Ontario, Canada (Nogrady, 1988); Kingston, Ontario, Canada (Nogrady, 1989); Star Lake, Norwich, Vermont, USA (Gilbert, 1989b); Colorado, USA (Turner, 1988); Lake Brooker, Florida, USA (Scheda & Cowell, 1988); Illinois river, USA (Brown *et al.*, 1989); Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Canada (Chengalath, 1977); Acton Lake, SW Ohio, USA (Chimney *et al.*, 1981); Standley Lake, Colorado, USA

(Pennak, 1966); Ross Lake, Florida, USA (Bienert, 1986); Northern Michigan inland lakes (Stemberger, 1990); Desaulniers reservoir, Québec, Canada (Pinel-Alloul, 1982); Laurentian Great Lakes (Stemberger, 1979); Lakes in Florida, USA (Dawes *et al.*, 1987); Cottonwood pond, Colorado, USA (Herrmann, 1977); Adirondack Lakes, NY State, USA (Siegfried *et al.*, 1984); Experimental ponds, C Florida, USA (Fry & Osborne, 1980); Atchafalaya river basin, Louisiana, USA (Holland *et al.*, 1983)

- Oriental region: Kashmir valley & high mountains of Kashmir Himalaya (Zutshi *et al.*, 1980); India (Sharma & Michael, 1980); Parakrama Samudra reservoir, Sri Lanka (Duncan & Gulati, 1983); Lake Parakrama, Sri Lanka (Fernando, 1984); Sri Lanka (Fernando, 1980a); Victoria reservoir, Sri Lanka (Piyasiri & Jayakody, 1991); Singapore (Karunakaran & Johnson, 1978); Rajasthan, India (Nayar, 1968); Kalimantan (Borneo), Indonesia (Koste, 1988a); Taiwan, Malaysia, Singapore, Indonesia, Sri Lanka & India (Sudzuki, 1989); Punjab State, India (Sharma, 1976; Sharma & Sharma, 1984); Phewa Tal, Pokhara valley, Nepal (Daems & Dumont, 1974); Sri Lanka. Common in ponds & lakes. The commonest species of *Trichocerca* in Sri Lanka (Chengalath, Fernando & Koste, 1973); Singapore & Taiwan. Cosmopolitan (Sudzuki, 1991); Parakrama Samudra, Sri Lanka (Duncan & Gulati, 1981; Duncan, 1983a; Fernando & Rajapaksa, 1983); Reservoir Samudra, Sri Lanka (Duncan, 1984); W & E Malaysia & Singapore (Fernando & Zankai, 1981); Ajwa reservoir, near Baroda, Gujarat, India (Jayangaudar, 1980); Impoundment Kinda, C Burma (Koste & Tobias, 1990); 2 Kumaun Himalayan lakes, UP, India (Sharma & Pant, 1985); Jammu (J & K), India (Jyoti & Sehgal, 1980); Lake Surinsar, Jammu (J & K), India (Jyoti & Sehgal, 1979); West Bengal, NE India (Sharma, 1979b); ? Storage reservoir Ajwa & Nimeta water works, Baroda, India (Wulfert, 1966)

- Australian region: River Murray, Aust. (Koste, 1979b); Billabongs Magela Creek, NW Aust. (Shiel & Koste, 1983); SW Aust. (Koste, Shiel & Brock, 1983); Tasmania (Koste & Shiel, 1987); Lake Barrine & L. Euramo, tropical Queensland crater lakes (Green, 1981); Farm dams near Gloucester, NSW, Aust. (Sudzuki & Timms, 1980); Qld, Vic & NSW, Aust. (Shiel & Koste, 1979); Sheepwash billabongs, near Molesworth, Vic, Aust. (Shiel, 1983); Alligator rivers area, NT, Aust. (Koste & Shiel, 1983a; Tait, Shiel & Koste, 1984); River Murray at Mannum, S Aust. (Koste, 1980); S Tasmania (Koste, Shiel & Tan, 1988); Rivers, billabongs & reservoirs of the Murray-Darling system, Aust. (Shiel, 1986); E Australia (Koste & Shiel, 1980b); Tasmania (Shiel, Koste & Tan, 1989); NW Aust.: Magela Creek area (Koste, 1981b); Billabongs of r.Murray floodplain, NSW, Aust. Particular common in humic acid waters in W Tasmania. Most common *Trichocerca* in Australian waters (Hillman & Shiel, 1991); Lake Alexandrina, New Zealand (Russell, 1960)

- Neotropical region: Embalse Rio Tercero Fish Culture Station, Cordoba province, Argentina (Marinone & Zagarese, 1991); Lago Tefé & L. Castanho, Amazonia, Brazil (Schaden, 1977c); Lago Castanho, L. Redondo, L. Jacaretinga, L. Cristalino & L. Taruma-Mirim, Brazil (Serruya & Pollingher, 1984); Lake Redondo, L. Jacaretinga & L. Cristalino, Amazonia Central, Brazil (Hardy, 1980); Ecuador : Costa & Sierra (Koste & Böttger, 1989); Ecuador : Oriente, Sierra & Costa (Koste & Böttger, 1992); Lake Rio Claro, a floodplain lake of the lower Orinoco, Venezuela (Vasquez & Medina, 1991); Lago do Macaco, Brazil (Koste, 1989a); Provinces in NW Argentina (José de Paggi, 1989); Lago Paroni, Brazil (Koste, 1974a); Amazonia, Brazil (Koste, 1972f); Antilles, Argentina, Brazil, Panama, Paraguay, Surinam & Venezuela (Koste & José de Paggi, 1982); Rio Yatapu & Rio Uatumá, Amazonia, Brazil (Koste & Hardy, 1984); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Panguana, Peru (Koste, 1988b); Near Concepcion, Paraguay (Koste, 1986b); Galapagos Islands (De Smet, 1987 & 1989); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Surinam (Leentvaar, 1975); Lake Llanquihue, Chile (Campos *et al.*, 1988); Orinoco river basin, Venezuela (Vasquez, 1984a); Lower Orinoco & its delta, Venezuela (Vasquez & Rey, 1989); Lake Hemmen, Rio Snia Missu, Mato Grosso, C Brazil (Green, 1972); Amazonia, Brazil (Hauer, 1965b; Thomasson, 1971); ? Argentine Patagonia (Kuczynski, 1987b); Lake Escondido, prov. Rio Negro, Argentina (Balseiro & Modenutti, 1990); All over Argentina (José de Paggi, 1990); Curaçao, Caribbean Islands & Costa Rica (Collado *et al.*, 1984); Aruba (Caribbean part of the Neotropis) (De Ridder, 1977); Lago Orsinera, Orinoco floodplain lake, Venezuela (Vasquez & Rey, 1992); River Parana, Brazil (Bonetto, 1986); Flooded savanna at Mantecal, Venezuela (Michelangelli *et al.*, 1979/1980); Antilles françaises (Pourriot, 1975); Araucarian lakes, W & E of the Andes, + ponds & pools in the same region, Chile (Thomasson, 1963b); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Lago do Castanho, C Amazonia, Brazil (Schaden, 1977b)

Trichocerca similis (Wierzejski), f. *binoculata* Hauer, 1965

- Neotropical region: Amazonia, Brazil (Hauer, 1965b); Brazil (Koste & José de Paggi, 1982)

Trichocerca similis (Wierzejski), f. *grandis* Hauer, 1965

- Australian region: Vic, Australia (Koste & Shiel, 1980c); S Tasmania; NSW, S Aust. & Vic, Aust. (Koste, Shiel & Tan, 1988); This large form is presently regarded as a ssp., *Tr. similis grandis* Hauer, 1965. It is

distinguished by larger, more elongated body and rel. shorter toes than the typical form. Common in Murray-Darling reservoirs (Shiel & Koste, 1992)

- Neotropical region: Rio Yatapu & Rio Uatumá, Amazonia, Brazil (Koste & Hardy, 1984); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Orinoco river basin, Venezuela (Vasquez, 1984a); Caroni river basin, Venezuela (Vasquez, 1984b); Lower orinoco river & its delta, Venezuela (Vasquez & Rey, 1989); Lago do Macaco, Brazil (Koste, 1989a); Amazonia, Brazil (Hauer, 1965b; Thomasson, 1971; Koste, 1972f); Prov. Corrientes, Argentina (José de Paggi, 1990); Pond in tropical rain forest, Peru (Schlüter, 1984); Lago do Castanho, C Amazonia, Brazil (Schaden, 1977b); Lago do Passarinho, L. Jurucui, L. Moratu, L. Tefé, Rio Negro, Lago do Castanho, L. Buiuçu, Amazonia, Brazil (Schaden, 1977c)

Trichocerca simonei De Smet

- Neotropical region: Santa Cruz, Galapagos Islands (Segers, 1991a); Mato Grosso, C Brazil (Turner & Da Silva, 1992)

Trichocerca stenroosi Haberman, 1978

- Palaearctic region: Lakes in Estonia (Haberman, 1978)

Trichocerca stylata (Gosse, 1851)

- Palaearctic region: Karstic Lake Percile, Latium, Italy (Margaritora *et al.*, 1981); Romania (Rudescu, 1960); Lake Balaton, Hungary (Zankai & Kertesz, 1967; Zankai, 1968; Zankai & Ponyi, 1970 & 1971a); All over Europe; Caucasus, Caspian area. Cosmopolitan (Bērziņš, 1978); River Klyas'ma, R.F. (Bogoslowski, 1961); Lake Kinneret, Israel (Serruya & Pollingher, 1984); (Landau *et al.*, 1988); Impoundment "Eisernes Tor", Danube, Romania (Brezeanu & Petén, 1973); Coast of S Finland (Eriksen, 1969); River Loire, France (Lair, 1980); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Delta of river Danube, Romania (Godeanu & Zinevici, 1983; Prunescu-Arion, 1989); Lakes in Estonia (Määemets, 1983); Man-made lake Zegrzynskie, Poland (Dusoge *et al.*, 1985); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Inundation area of Danube delta, Romania (Dorobantu, 1964); Lake Volvi, Macedonia, Greece (Zarfdjian *et al.*, 1990); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b); Lac Aydat, Massif Central, France (Persoone *et al.*, 1968; Lair, 1978); Japanese inland waters (Yamamoto, 1960a); Southwest islands of Japan (Suzuki *et al.*, 1992); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); In 5 localities in Belgium (De Ridder, 1986); River Danube (Enaceanu & Brezeanu, 1970); River Elbe estuary, Germany (Schulz, 1961); Lakes, N of Bucarest, Romania (Godeanu, 1969b); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); River Vistula in the Warsaw area, Poland (Klimowicz, 1981); Canal of river Odra, Poland (Klimowicz, 1979); Small water bodies in the Mikolajski area, P. (Klimowicz, 1967); Rhine waters near Wiesbaden, Germany (Noll, 1972); Lac Léman, Fr./CH (Martin, 1961); Forêt de Sénard, near Paris, France (Ansellem & Clément, 1988); Lake Tüzer See, Neubrandenburg, Germany (Doll, 1964); Lake at Wijk-bij-Duurstede; Zuidelijke Vechtplassen; Kil van Hurwenen; area between Linge & Rhine, id. between Linge & Waal; Loosdrecht Lakes, NL (Dresscher, 1976); Karelian inland waters, R.F. (Kutikova, 1965); Lac d'Annecy, France (Balvay & Druart, 1992); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); Hamburg area, Germany (Rühmann, 1961); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Middle course river Loire, France (Lair & Sargas, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Pond in the Paris suburb, France (Pourriot, 1965); Aegean lakes, Macedonia, ex-Yugoslavia-Greece (Serafimova-Hadzisce, 1974); Serbian section of river Danube (Zivkovic, 1987); Spain : Barcelona (Velasco, 1990); Lake Biwa-ko, Japan (Yamamoto, 1960b)

- Nearctic region: Canada (Chengalath, 1977); Illinois, USA (Bērziņš, 1984); Colorado, USA (Turner, 1988); Laurentian Great Lakes (Stemberger, 1979); Bayou de Siard, Monroe, Louisiana, USA (Gallagher, 1965)

- Oriental region: Tjigombong (Serruya & Pollingher, 1984); Rajasthan, India (Nayar, 1968); Taiwan (Suzuki, 1988); Taiwan, Malaysia, Indonesia, Sri Lanka, India (Suzuki, 1989); Kalimantan (Borneo), Indonesia (Koste, 1988a); India (Sharma & Michael, 1980); Lake Parakrama, Sri Lanka (Fernando, 1984); Sri Lanka (Fernando, 1980a); Sri Lanka. Fairly common. Found in lakes & ponds (Chengalath, Fernando & Koste, 1973); W & E Malaysia & Singapore (Fernando & Zankai, 1981); Parakrama Samudra, Sri Lanka (Fernando & Rajapaksa, 1983)

- Australian region: Vic, Australia (Shiel & Koste, 1979); Lower river Murray, S Aust. (Shiel, Walker & Williams, 1982); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Koste & Shiel, 1983a; Tait, Shiel & Koste, 1984); Rivers, billabongs & reservoirs of the Murray-Darling system, Aust. (Shiel, 1986); Rare; Qld, Tas & Vic (Shiel & Koste, 1992); Lake Alexandrina, New Zealand (Russell, 1960)

- Neotropical region: Chile (Löffler, 1962); Lago Buiuçu, Amazonia, Brazil (Schaden, 1977c); Mato Grosso, C Brazil (Turner & Da Silva, 1992); Prov. Buenos Aires & Tierra del Fuego, Arg.; Brazil (Olivier, 1965);

Araucarian lakes, W & E of the Andes, Chile (Thomasson, 1963b); Chilean lakes (Schmid-Araya, 1991); River Parana, Argentina (Paggi & José de Paggi, 1974); Orinoco river basin, Venezuela (Vasquez, 1984a); Lower Orinoco & its delta, Venezuela (Vasquez & Rey, 1989); Amazonia, Brazil (Hauer, 1965a; Thomasson, 1971; Koste, 1972f); Brazil, Argentina, Chile, Venezuela & Panama (Koste & José de Paggi, 1982); Rio Yatapu, Amazonia, Brazil (Koste & Hardy, 1984); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Prov. Tierra del Fuego, Chubut, Rio Negro, Neuquen, Corrientes, Santa Fé & Buenos Aires, Argentina (José de Paggi, 1990); Panama (Collado *et al.*, 1984); Lago de Valencia, Venezuela (De Infante, 1980; Saunders & Lewis, 1988)

***Trichocerca sulcata* (Jennings, 1894)**

- Palaearctic region: Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Brook in Lao-Shan mountains, China (Wulfert, 1968); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Lakes in S Sweden (Arnemo *et al.*, 1968); River Ybbarspå, Sweden (Bērziņš, 1979); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Coast of S Finland (Eriksen, 1969); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Romania (Rudescu, 1960); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Almost all over Europe (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1960 & 1970); Lake Balaton, Hungary (Zankai, 1968); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960; Dresscher, 1976); In reed border of the Danube delta, Romania (Rudescu, 1968)
- Nearctic region: Canada (Chengalath, 1977); Ross Lake, Florida, USA (Bienert, 1986); Laurentian Great Lakes (Stemberger, 1979)
- Australian region: Avoca river, Vic, Aust. (Bērziņš, 1982); Vic, Aust. (Shiel & Koste, 1985); Recorded by Bērziņš, 1982. No ecological information. Not seen in our collections (Shiel & Koste, 1992); Chatham Islands, New Zealand (Russell, 1960)
- Neotropical region: Argentina (Koste & José de Paggi, 1982); Prov. Rio Negro, Argentina (José de Paggi, 1990)

***Trichocerca sulcata* (Jennings), f. *minuscula* Rodewald, 1940**

- Palaearctic region: Peat bogs Poiana Stampei, Bukowina, Rom. (Rudescu, 1966); Carpathes (Bērziņš, 1978)

***Trichocerca taurocephala* (Hauer, 1931)**

- Palaearctic region: Middle course of river Loire, France (Lair & Sargos, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Lake Ohrid, Balkan peninsula (Stankovic, 1960); River Loire, in St. Laurent-les-Eaux, France (Pourriot *et al.*, 1982); In periphyton of mosses and *Myriophyllum* in the Danube, and in some Alpine brooks, Austria (Donner, 1964); Shores of embanked and free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); Dübener Heide, N Germany (Wulfert, 1961); Lake Bodensee near Konstanz, Germany (Althaus, 1962); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Romania (Rudescu, 1960); Sea & brackish waters in S Finland (Björklund, 1972a); River Loire, France (Lair, 1980); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Asiatic Turkey (Dumont & De Ridder, 1987); River Hase, NW Germany (Koste, 1976b); Courland Spit, R.F. (Zhadin *et al.*, 1972); Alps, W Balkan, C Eur. mts., Pontic prov., C & E Eur. plains, NW Russian taiga (Bērziņš, 1978); River Grabia basin, Poland (Pawlowski, 1960, 1968 & 1970); Lake Balaton, Hungary (Zankai, 1968); Anatolia, Turkey (Segers *et al.*, 1992)
- Nearctic region: In Florida saltwater psammon, USA (Turner, 1990b)

***Trichocerca tenuidens* (Hauer, 1931)**

- Palaearctic region: Lake Balaton, Hungary (Zankai & Ponyi, 1970); C Eur. mountains (Bērziņš, 1978)
- Nearctic region: Canada (Chengalath, 1977); Kentucky, USA (Bērziņš, 1984); Pinehirst Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamootttil, 1975)

***Trichocerca tenuior* (Gosse, 1886)**

- Palaearctic region: Serbian section of river Danube (Zivkovic, 1987); Spain : Granada & Valencia (Velasco, 1990); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Lake Balaton, Hungary (Zankai, 1968); Japanese inland waters (Yamamoto, 1960a); Southwest islands of Japan (Suzuki *et al.*, 1992); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); River Danube (Enaceanu & Brezeanu, 1970); Peat bog Chambedaze, Puy-de-Dôme, France (Francez, 1982 & 1987); Water-courses in the Bucegi mts., Romania (Godeanu, 1963b); Small permanent ponds in the Bucegi mts., Rom. (Godeanu, 1969a); Lakes of the Black Sea shore, Romania (Godeanu, 1970b); Obretin embanked basin, Danube delta, Romania (Godeanu *et al.*, 1973); Small water bodies in the Mikolajki area, Poland (Klimowicz, 1967); Anatolia, Turkey (Segers *et al.*, 1992); In periphyton of mosses and *Myriophyllum* in the Danube, and in some Alpine brooks, Austria (Donner, 1964); Shores of embanked or free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); River Salzach & its tributaries, Austria (Donner, 1970); Lac Léman, Fr./CH (Martin, 1961); (Balvay & Laurent, 1981 & 1990a); Dübener Heide, N Germany (Wulfert,

1961); Peat bog Chambéthane, Massif Central, France (Persoone *et al.*, 1968; Francez, 1988); Lake Watergraafsmeer; ditches in Amsterdam; Nat. Res. Meyendel; Prov. Zuid-Limburg; De Venematen, NL (Dresscher, 19796); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); River Moldova, tributary of the Danube, Rom. (Prunescu-Arion, 1968); In reed border of the Danube delta, Romania (Rudescu, 1968); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); River Loire in St. Laurent-les-Eaux, France (Pourriot *et al.*, 1982); Hamburg area, Germany (Rühmann, 1961); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Middle course of river Loire, France (Lair & Sargas, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979; V'yushkova & Lakhnova, 1974); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990); Lakes around mountain Fuji, Japan (Suzuki, 1971); Mountain Fuji, Japan (Suzuki, 1978a); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Impoundment Saidenbach, Germany (Althaus, 1976); River Klyas'ma, R.F. (Bogoslawski, 1961); Small lakes in Estonia (Haberman & Mäemets, 1987); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Lakes in Estonia (Haberman, 1978); Coast of S Finland (Eriksen, 1969); Romania (Rudescu, 1960); River Loire, France (Lair, 1980); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Asiatic Turkey (Dumont & De Ridder, 1987); Sierra Nevada, Spain (Morales-Baquero, 1987); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); District Ostrava, CSR (Dvorakova, 1960b); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Lake Mikolajskie, Mazurian lakeland, Poland (Klimowicz, 1972); Alpine lakes, Sierra Nevada, Spain (Morales-Baquero *et al.*, 1989); Darnsee near Epe, NW Germany (Koste, 1962); Lake Orog-Nur, Gobi desert, Mongolia (Koste & Wulfert, 1964); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Achmer Grasmoor, NW Germany (Koste, 1968b); River Hase, NW Germany (Koste, 1976b); Lake Dümmer, NW Germany (Koste & Poltz, 1983 & 1984); Fish ponds near Warsaw, Poland (Hillbricht-Ilkowska, 1964); Sphagnum bogs in the Massif Central, France (Batut, 1965); All over Europe; Caucasus, Caspian area. Cosmopolitan (Bērziņš, 1978); Inundation area of the Danube delta, Romania (Dorobantu, 1964); Lower Silesia & river Grabia basin, Poland (Pawlowski, 1960); River Grabia basin, Poland (Pawlowski, 1968, 1970 & 1973); Overmere, Belgium (De Ridder, 1986); Near water works at Okocim, Poland (Klimowicz, 1975)

- Nearctic region: Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); NE Canada (Chengalath & Koste, 1983); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Colorado, USA (Turner, 1988); Kingston, Ontario, Canada (Nogrady, 1989); Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Canada (Chengalath, 1977); Wisconsin, Illinois & Kentucky (Bērziņš, 1984); Miller's Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975)

- Oriental region: Ricefields in NE Thailand (Heckman, 1979); Malaysia & Indonesia (Suzuki, 1989); Kalimantan (Borneo), Indonesia (Koste, 1988a); Fishponds in Thailand (Boonsom, 1984a); Thailand (Boonsom, 1984b); Lake Ranu Lamongau, Indonesia (Green *et al.*, 1976); Sri Lanka (Mendis & Fernando, 1962); (Fernando, 1969)

- Australian region: In many localities in Australia (Bērziņš, 1982); Tasmania (Koste & Shiel, 1987); Bromfield Swamp, tropical Queensland crater lake (Green, 1981); NSW & Qld, Australia (Shiel & Koste, 1979); River Murray, Aust. (Koste, 1979b); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Rare in mainland samples (Vic), more common in Tasmania (Shiel & Koste, 1992); Lincoln, Christchurch & Takahey valley, New Zealand (Russell, 1960)

- Neotropical region: Ecuador : Oriente (Koste & Böttger, 1989); Ecuador : Oriente & Costa (Koste & Böttger, 1992); Chile (Löffler, 1962); Amazonia, Brazil (Hauer, 1965b; Koste, 1972f); Araucarian lakes, E of the Andes (Thomasson, 1963b); Varzea near Santarem, Brazil (Koste, 1974b); Antilles, Brazil, Chile, Panama, Paraguay & Venezuela (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Panguana, Peru (Koste, 1988b); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Chilean lakes (Schmid-Araya, 1991); Orinoco river basin, Venezuela (Vasquez, 1984a); ? Provinces in NW Argentina (José de Paggi, 1989); ? Prov. Chubut, Argentina (José de Paggi, 1990); Panama (Collado *et al.*, 1984); Flooded savanna at Mantecal, Venezuela (Michelangeli *et al.*, 1979/1980); Antilles françaises (Pourriot, 1975); Mato Grosso, C Brazil (Turner & Da Silva, 1992)

Trichocerca tigris (O.F. Müller, 1786)

- Palaearctic region: Slovenia, ex-Yugoslavia (Markic, 1983); Romania (Rudescu, 1960); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Lake Mikolajskie, Mazurian lakeland, Poland (Klimowicz, 1972); N Swedish Lapland (Pejler, 1962a); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); Lake Balaton, Hungary (Zankai & Ponyi, 1970 & 1971a); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, Germany (Koste, 1965); River Hase, NW Germany (Koste, 1976b); Fishponds near Warsaw, Poland

(Hillbricht-Ilkowska, 1964); Sphagnum bogs in the Massif Central, France (Batut, 1965); Volga river, R.F. (V'yushkova & Lakhnova, 1976); All over Europe; Caucasus, Caspian area. Cosmopolitan (Bērziņš, 1978); Peat bog Laptici, Bucegi mts. Romania (Godeanu, 1970a); Lake Myvatn, Iceland (Adalsteinsson, 1979); Lake Bodensee, near Konstanz, Germany (Althaus, 1962); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Coast of S Finland (Eriksen, 1969); Delta of river Danube, Romania (Godeanu & Zinevici, 1983); River Grabia basin, Poland (Pawlowski, 1968 & 1970); Lake Balaton, Hungary (Zankai, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b); N & W Iceland (De Ridder, 1969b, 1970 & 1972); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); River Danube (Enaceanu & Brezeanu, 1970); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Anatolia, Turkey (Segers *et al.*, 1992); In periphyton of mosses and *Myriophyllum* in the Danube, and in some Alpine brooks, Austria (Donner, 1964); ? Shores of dammed or free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); In the lower Danube, Romania (Enaceanu, 1964); Lac Léman, Fr./CH (Balvay & Laurent, 1981); Dübener Heide, N Germany (Wulfert, 1961); R. Rijn, Waal & Yssel; Nat. Res. Het Hol; R. Maas, NL (Dresscher, 1976); River & valley in the Tatra mountains, Poland (Madalinski, 1981); Bog pond near Budapest, Hungary (Nogrady, 1962); In reed border of river Danube delta, Romania (Rudescu, 1968); 3 lakes of Sosnowickie, Poland (Radwan, 1969); Parczew area peat bogs, Poland (Radwan, 1974); Peat bogs Poiana Stampei, Bukowina, Poland (Rudescu, 1966); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Ponds & pools in the Paris suburb, France (Pourriot, 1965 & 1979); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990); Peats in the Caucasus (Tarnogradsky, 1961); Lago Trasimeno, Italy (Taticchi, 1968); Spain : Zamora (Velasco, 1990); Lake Biwa-ko, Japan (Yamamoto, 1960b); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)

- Nearctic region: Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Canada (Chengalath, 1977); NE Canada (Chengalath & Koste, 1983); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Miller's Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Wisconsin, Illinois & Kentucky, USA (Bērziņš, 1984); Laurentian Great Lakes (Stemberger, 1979); Mt. Tremblant Prov. Park, Québec, Canada (Nogrady, 1980); Bay of Quinte, Lake Ontario, Canada (Nogrady, 1988)

- Oriental region: India (Sharma & Michael, 1980); Taiwan, Malaysia & India (Sudzuki, 1989); W & E Malaysia & Singapore (Fernando & Zankai, 1981); West Bengal, NE India (Sharma, 1979b); Sri Lanka (Mendis & Fernando, 1962); (Fernando, 1969)

- Australian region: Tasmania (Koste & Shiel, 1987); NSW, Qld & Vic, Aust. (Shiel & Koste, 1979); Lower river Murray, S Aust. (Shiel, Walker & Williams, 1982); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); In many localities in Australia (Bērziņš, 1982); Common in billabongs, Vic, Tas (Shiel & Koste, 1992); Lincoln, New Zealand (Russell, 1960)

- Neotropical region: Ecuador : Oriente, Costa & Sierra (Koste & Böttger, 1989); Ecuador : Oriente & Sierra (Koste & Böttger, 1992); Galapagos Islands (De Smet, 1987 & 1989); Argentina (Modenutti & Claps, 1988); Lago Paroni, Brazil (Koste, 1974a); Amazonia, Brazil (Hauer, 1965b; Koste, 1972f); Varzea near Santarem, Brazil (Koste, 1974b); Argentina, Brazil, Jamaica, Panama, Paraguay & Venezuela (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Harduy, 1984); Rio Nhamunda, Amazonia, Brazil (Bendorff, Koste & Smirnov, 1982); Near Panguana, Peru (Koste, 1988b); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Chilean lakes (Schmid-Araya, 1991); Prov. Tierra del Fuego, Chubut & R. Negro, Arg. (José de Paggi, 1990); Caribbean Islands & Panama (Collado *et al.*, 1984); Caribbean section of Neotropical region (De Ridder, 1977); Prov. Rio Negro & Tierra del Fuego, Argentina (Olivier, 1965); Araucarian lakes, E of the Andes, Chile (Thomasson, 1963b); Mato Grosso, C Brazil (Turner & Da Silva, 1992)

- Antarctic region: Antarctica (Dartnall, 1983); (Dartnall & Hollowday, 1985); Antarctica & Subantarctica (José de Paggi & Koste, 1984)

Trichocerca tigris (O.F. Müller), f. *monostyla* Russell, 1951

- Australian region: Westland, New Zealand (Russell, 1960)

Trichocerca tortuosa (Myers, 1936)

- Nearctic region: Canada (Chengalath, 1977); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988)

- Australian region: Westland, New Zealand (Russell, 1960)

Trichocerca tropis Hauer, 1938

- Oriental region: India (Sharma & Michael, 1980; Sudzuki, 1989)

Trichocerca uncinata (Voigt, 1902)

- Palaearctic region: In periphyton of mosses and *Myriophyllum* in the Danube, and in some Alpine brooks,

Austria (Donner, 1964); River Salzach and its tributaries, Austria (Donner, 1970); Lake Bodensee near Konstanz, Germany (Althaus, 1962); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Lake Stråken, Aneboda, Sweden (Bérziňš, 1972); Romania (Rudescu, 1960); Coast of S Finland (Eriksen, 1969); C & S Sweden (Bérziňš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); N Swedish Lapland (Pejler, 1962a); Edgeøya, Svalbard (De Smet *et al.*, 1988b); Bjørnøya (Bear Island), Svalbard (De Smet, 1988); Darnsee near Epe, NW Germany (Koste, 1962); All over Europe (Bérziňš, 1978); River Grabia basin, Poland (Pawlowski, 1960); Lake Balaton, Hungary (Zankai, 1968); N & W Iceland (De Ridder, 1967a, 1970 & 1972); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973)

- Nearctic region: N America (Bérziňš, 1978); Canada (Chengalath, 1977); Arctic N America (Chengalath & Koste, 1989)

- Australian region: NSW, Aust. (Shiel & Koste, 1985); Unconfirmed record from Tabrabucca Creek, Barrington, NSW (Bérziňš, 1982). No ecological information. Not seen in our collections (Shiel & Koste, 1992)

- Neotropical region: Amazonia, Brazil (Koste, 1972f); Lago Paroni, Brazil (Koste, 1974a); Brazil (Koste & José de Paggi, 1982)

Trichocerca vargai Wulfert, 1961

- Palaearctic region: Dübener Heide, N Germany (Wulfert, 1961); C Eur. plain (Bérziňš, 1978)

Trichocerca vargai Wulfert var.

- Palaearctic region: L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978)

Trichocerca vernalis (Hauer, 1936)

- Palaearctic region: Lakes of the Black Sea shore, Romania (Godeanu, 1970b); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Anatolia, Turkey (Segers *et al.*, 1992); Romania (Rudescu, 1960); "Altes Land", near Hamburg, Germany (Caspers & Heckman, 1981 & 1982); Near Asperg, Württemberg, Germany (Klement, 1960); C Eur. mts., Hungarian plain, Pontic prov., C Eur. plains (Bérziňš, 1978); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968)

- Nearctic region: N America (Bérziňš, 1978); Canada (Chengalath, 1977); Kingston, Ontario, Canada (Nogrady, 1989); Cranberry bog, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975)

- Oriental region: Lakelet Tan Daha, Kathmandu valley, Nepal (Daems & Dumont, 1974)

- Australian region: Creek in Australia (Bérziňš, 1982); Vic, Aust. (Shiel & Koste, 1985); Rivers, billabongs & ephemeral pools. River Murray & Mitta Mitta river, Vic (Shiel & Koste, 1992)

Trichocerca voluta Murray, 1913

- Neotropical region: Brazil & Panama (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983); Varzea near Santarem, Brazil (Koste, 1974b); Near Concepcion, Paraguay (Koste, 1986b); Panama (Collado *et al.*, 1984)

Trichocerca weberi (Jennings, 1903)

- Palaearctic region: Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Lake Palaeostomi, Georgia, R.F. (Haberman, 1982; Haberman & Laugaste, 1982); Lakes in Estonia (Haberman, 1978); Coast of S Finland (Eriksen, 1969); C & S Sweden (Bérziňš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Pools of upper Elbe, CSR (Dvorakova, 1961); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Nat. Res. Neeeruti Lakes, NE Estonia (Mäemets A. & A., s.d.); Edgeøya, Svalbard (De Smet *et al.*, 1988b); Bjørnøya (Bear Island), Svalbard (De Smet, 1988); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Fish ponds near Warsaw, Poland (Hillbricht-Ilkowska, 1964); All over Europe, Caucasus (Bérziňš, 1978); River Grabia basin, Poland (Pawlowski, 1960 & 1970); Southwest islands of Japan (Suzuki *et al.*, 1992); Lake Balaton, Hungary (Zankai, 1968); N & W Iceland (De Ridder, 1967a, 1969b, 1970 & 1972); Lake Neusiedlersee, Austria (Donner, 1979); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); River Salzach & its tributaries, Austria (Donner, 1970); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Anatolia, Turkey (Segers *et al.*, 1992); Etang de Pommereau, Loir-et-Cher, France (Tassigny *et al.*, 1970); Nat. Res. "Het Hol", Kortenhoef, NL (Dresscher, 1976); Pond in the Paris suburb, France (Pourriot, 1965); Periphytic in ponds in the Ile-de-France (Pourriot, 1970); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); In reed border of the Danube delta, Romania (Rudescu, 1968); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); Parczew area peat bogs, Poland (Radwan, 1974); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Pool, S of Trondheim, Norway (Thomasson, 1979); Lago Trasimeno, Italy (Taticchi, 1968); Serbian section of river Danube (Zivkovic, 1987); Spain : Barcelona, Orense, Pontevedra; Canaries : Gran Canaria (Velasco, 1990); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973)

- Nearctic region: Stoneman Lake, Arizona, USA (Green, 1986); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); Canada (Chengalath, 1977); NW Canada (Chengalath & Koste, 1987); Arctic N America (Chengalath & Koste, 1989); Kingston, Ontario, Canada

(Nogrady, 1989); Wisconsin, USA (Bērziņš, 1984); Miller's Lake, near Waterloo, Ontario, Canada (Chengalath & Mulamoottil, 1975); Experimental ponds, C Florida, USA (Fry & Osborne, 1980)

- Oriental region: Singapore, Indonesia & India (Suzuki, 1989); India (Sharma & Michael, 1980); Singapore. Distribution : warm waters (Suzuki, 1991); Punjab State, NW India (Sharma, 1976; Sharma & Michael, 1984); West Bengal, NE India (Sharma, 1979b)

- Australian region: In many localities in Australia (Bērziņš, 1982); Lake Barrine, tropical Queensland crater lake, Australia (Green, 1981); Qld, Aust. (Shiel & Koste, 1979); Alligator rivers area, NT? Aust. (Tait, Shiel & Koste, 1984); Lake Garcia, Tasmania. Also in Qld & Vic, Aust. (Koste, Shiel & Tan, 1988); Rare : Qld, Tas & Vic (Shiel & Koste, 1992); Lincoln, New Zealand (Russell, 1960)

- Neotropical region: Antilles françaises (Pourriot, 1975); Araucarian lakes, W of the Andes, Chile (Thomasson, 1963b); Antilles, Argentina, Brazil, Panama & Venezuela (Koste & José de Paggi, 1982); Amazonia, Brazil (Hauer, 1965b; Koste, 1972f); Lago Paroni, Brazil (Koste, 1974a); Varzea near Santarem, Brazil (Koste, 1974b); Near Concepcion, Paraguay (Koste, 1986b); Chilean lakes (Schmid-Araya, 1991); Galapagos Islands (De Smet, 1987 & 1989); Argentina (Modenutti & Claps, 1988); Prov. Rio Negro & Santa Fé, Argentina (José de Paggi, 1990); Caribbean Islands & Panama (Collado *et al.*, 1984)

Trichocerca spec.

- Palaearctic region: Lac de Pétichet, Isère, France (Gay, 1979); Lake Meeuwenven, near Nijmegen, NL (Geelen, 1969); Lake Bodensee (Walz, 1986); Lago Scuro Parmense, N Italy (Antonietti *et al.*, 1988); Middle course of river Po, Italy (Ferrari *et al.*, 1986); Lake Glubokoje, near Moscow, R.F. (Ghilarov, 1976); Lake Mikolajskie, Masuria, Poland (Karabin, 1978); Lake Biwa, Japan (Narita & Mori, s.d.); Lake Banyoles, NE Spain (Miracle, 1977); River Loire, France (Lair, 1980); Lake Suwa in Japan (Hanazoto *et al.*, 1989; Hanazoto & Yasumo, 1988); Lake Bielersee, CH (Berner-Fankhauser, 1983 & 1986); Lakes in Estonia (Mäemets, 1983); Lake Königssee, Bavaria, Germany (Laxhuber, 1987); Loch Leven, Scotland (May, 1987); Impoundment Kninicki, near Brünn, CSR (Kubicek, 1964); Lake Quackjeswater & L. Breedewater, Voorne & Lake Vogelmeer, Ymuiden, NL (Leentvaar, 1963b); Lake Laarder Wasmeer, Veluwe, NL (Leentvaar, 1963c); Broads & peat pits in NW Overijssel, NL (Leentvaar, 1965); Lake Zwanewater in Callantsoog & L. Oerd on Ameland, NL (Leentvaar, 1967); Po river, Italy (Ferrari *et al.*, 1989); Lake Numasawa, Japan (Suzuki *et al.*, 1989); Minorca, Spain (de Manuel, 1990); Norfolk Broads, UK (Irvine *et al.*, 1990); Loch Rusky, Scotland (Romo, 1990); Fish-fattening ponds in the Ukraina (Krazhan *et al.*, 1976); Rivers in W Kazakhstan, R.F. (Petrova, 1976); Dniester estuary, R.F. (Polishchuk, 1976); Aral Sea spawning grounds, R.F. (Yerekeyev & Kazakhbayev, 1972); "Kleine vijver", fort 8, Hoboken (Antwerp), B. (De Smet *et al.*, 1989); Terek delta, Astrakhan peninsula, R.F. (Chuykov & Deneskin, 1985); Nat. Res. Neeruti Lakes, NE Estonia (Mäemets A. & A., s.d.); Lake Monterosi, Latium, Italy (Stella & Margaritora, 1966); Luka nad Olzon, near Ostrava, Silesia, CSR (Sladecek, 1969); Latorica, E Slovakia lowland, CSR (Terek, 1990a); Pool, S of Trondheim, Norway (Thomasson, 1979); Lago Trasimeno, Italy (Taticchi, 1968); Saltmarshes of Cadiz, Spain (Yufera *et al.*, 1984); Lake Mikolajskie, Poland (Ejsmont-Karabin, 1974); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Impoundment Weidatalsperrre, E Thüringen, G. (Hoffmann, 1966); New impoundments in the CSR (Kubicek, 1968); Lake in the Leersumse Veld, NL (Leentvaar, 1967b); In periphyton of mosses and *Myriophyllum* in the Danube, and in some Alpine brooks, Austria (Donner, 1964); ? Shores of dammed and free flowing stretches of the Danube, at the German-Austrian border (Donner, 1972); River Salzach & its tributaries, Austria (Donner, 1970); In the lower Danube, Romania (Enaceanu, 1964); Lake Kummerower See, Mecklemburg, Gem. (Kalbe *et al.*, 1965); River Danube & its tributaries (Bothar, 1982); Pond near Besse-en-Chandesse, France (Omaly & Dévaux, 1971); Lake Nimetön, S Finland (Arvola *et al.*, 1986); All over the Netherlands (Dresscher, 1976); Etang de Soustons & Et. de Hardy, SW France (De Ridder, 1967b); Lake Chambon, Puy-de-Dôme, France (Persoone *et al.*, 1968); Pool in Fontainebleau Forest, Ile-de-France, Fr. (Pourriot, 1976); Hamburg area, Germany (Rühmann, 1961 & 1965a); Middle course of river Loire, France (Lair & Sargas, 1981); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990); Garden ponds in Tokyo, Japan (Suzuki, 1975); Fukushima Prefecture, Japan (Suzuki, 1982); River Elbe estuary, Germany (Schulz, 1961); Canals in the Hran area, CSR (Terek, 1990c); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Akgöl, Selçuk-Izmir, Turkey (Ustaoglu & Balik, 1987); Lake Marmara, Turkey (Ustaoglu, in press); Lake Mikolajskie & L. Taltowisko, Poland (Gliwicz *et al.*, 1978); Lake Onega, R.F. (Nikolayev, 1972); Lake Giulianello, Latium, Italy (Stella & Margaritora, 1972); Lake Poletskoye, 73 km of Moscow, R.F. (Smirnov, 1961); Darss-Zingst estuary, S Baltic Sea (Heerkloss *et al.*, 1991); Lough Erne system, NW Ireland (Gibson *et al.*, 1980); Southwest islands of Japan (Suzuki *et al.*, 1992); Highly eutrophic lake Czarna Kuta, Poland (Brinska *et al.*, 1976); Loosdrecht Lakes, NL (Van Tongeren *et al.*, 1992); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Clay pits in Stuivekenskerke, Belgium (Espeel, 1979); Lake Neujahrslöch, Hessen, Germany (Menzel, 1968)

- Nearctic region: Atchafalaya river basin, Louisiana, USA (Holland *et al.*, 1983); Desaulniers reservoir, Québec, Canada (Pinel-Alloul *et al.*, 1982); L. Whitney, Cumberland Island Nat. Seashore, USA (Stoneburger & Schmock, 1980); Lakes in Florida (Dawes *et al.*, 1987); Hogans pond, Newfoundland, Canada (Davis, 1972); Mirror Lake, New Hampshire, USA (Strayer, 1985); Lake Erie, Canada (Davis, 1969); Bauline Long pond, Newfoundland, Canada (Davis, 1973); Lakes at Santee, California, USA (Nogrady, 1982); Lake 122 & 132, NW Ontario, Canada (Schindler & Novén, 1971); Rhode river estuary, Maryland, USA (Dolan & Gallagos, 1992); Long Pond, St. John's, Newfoundland, Canada (O'Connell & Andrews, 1977); Severson Lake, Minnesota, USA (Comita, 1972); Lake Oglethorpe, Georgia, USA (Sanders *et al.*, 1989); Colorado, USA (Turner, 1988); Lake Michigan, USA (Sandgren & Lehman, 1990); Lake St. George, King Township, Ontario, Can. (Malone & Mc Queen, 1983); Degg's Lake, Colorado, USA (Pennak, 1966); Lakes in NE Pennsylvania, USA (Havens & Da Costa, 1988); Lake Brooker, Florida, USA (Scheda & Cowell, 1988); Northern Michigan inland lakes (Stemberger, 1990); Kalamazoo river, Michigan, USA (Prins & Davis, 1966); Vinton County, Ohio, USA (Evans, 1984); Experimental ponds, C Florida, USA (Fry & Osborne, 1980)
- Oriental region: Lake Sat Tal, Utar Pradesh, India (Pant *et al.*, 1985); Montane lakes of Bali (Green *et al.*, 1978); Lake Surinsar, Yammu (Y & K), India (Jyoti & Sehgal, 1979); Malacca peninsula (Dunn, 1970); Parakrama Samudra reservoir, Sri Lanka (Duncan, 1983a & 1984); (Duncan & Gulati, 1983); Victoria reservoir, Sri Lanka (Piyasiri & Jayakodi, 1991); Parakrama Samudra, Sri Lanka (Duncan & Gulati, 1981); Roshamara tank; Naini Lake; Stocking tank, Delhi, India (George, 1966); Lakelet Tan Daha, Kathmandu valley, Nepal (Daems & Dumont, 1974); Impoundment Kinda, C Burma (Koste & Tobias, 1990); Saline crater lake Batur, Bali (Green, 1986); Lake Manguao, Palawan, the Philippines (Davies & Green, 1990); Great Lake & Malayan lakes (Serruya & Pollingher, 1984)
- Australian region: Creek in Australia (Bērziņš, 1982); Billabongs river Murray, SE Aust. (Shiel & Koste, 1983); Lake Bullenmerri (slightly alkaline), Aust. (Timms, 1981); Lake Okaro, New Zealand (Forsyth & James, 1991)
- Neotropical region: Rapel reservoir, C Chile (Soto *et al.*, 1984); Galapagos Islands (De Smet, 1987 & 1989); Surinam (Leentvaar, 1975); Eutrophic water Santa Fé, Argentina (José de Paggi, 1976a); River Santa Fé, Argentina (José de Paggi, 1976b); River Parana, Argentina (José de Paggi, 1985); River Saladillo basin, Argentina (José de Paggi & Koste, 1988); Lower Orinoco river & its delta, Venezuela (Vasquez & Rey, 1989); Broa reservoir, Sao Carlos, Sao Paulo State, Brazil (Matsumara-Tundisi & Tundisi, 1976); Arroyo Rodriguez, prov. de Buenos Aires, Arg. (Modenutti, 1987); Provinces in NW Argentina (José de Paggi, 1989); Lago Camaleão, Amazonia, Brazil (Koste, Robertson & Hardy, 1984); Near Panguana, Peru (Koste, 1988b); Lake Rio Claro, a floodplain lake of the lower Orinoco, Venezuela (Vasquez & Medina, 1991); Billings reservoir, Sao Paulo, Brazil (Sendacz, 1984); Apure river, W Venezuela (Saunders & Lewis, 1988); Amazonia, Brazil (Hauer, 1965b; Gillard, 1967;; Thomasson, 1971); Araucarian lakes, W & E of the Andes & Araucarian ponds & pools, Chile (Thomasson, 1963b); Chile (Löffler, 1962)

Trichotria Bory de Saint Vincent, 1827

Trichotria buchneri Koste, Shiel & Tan, 1988

- Australian region: Strahan, S Tasmania (Koste, Shiel & Tan, 1988); Tasmania (Shiel, Koste & Tan, 1989); W Coast of Tasmania (Koste & Shiel, 1989a)

Trichotria caudata (Lucks, 1912)

- Neotropical region: Ecuador : Sierra (Koste & Böttger, 1992)

Trichotria curta (Skorikov, 1914)

- Palaearctic region: Heiluntsian province, China (Bogoslowski, 1962); Romania (Rudescu, 1960); Asiatic Turkey (Dumont & De Ridder, 1987); Volga river, R.F. (V'yushkova & Lakhnova, 1974; Mordukhai-Boltovskoi, 1979); Pontic province, E Eur. plain, N Sweden, Caspian area (Bērziņš, 1978); Serbian section of river Danube (Zivkovic, 1987)

- Australian region: Not yet recorded from Australia (Koste & Shiel, 1989a)

Trichotria eucosmeta Myers, 1934

- Nearctic region: Labrador (Koste, 1984); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988)

- Australian region: Not yet recorded from Australia (Koste & Shiel, 1989a)

Trichotria pocillum (O.F. Müller, 1776)

- Palaearctic region: Romania (Rudescu, 1960); Bjørnøya (Bear Island), Svalbard (De Smet, 1988); Po river, Italy (Ferrari *et al.*, 1989); Minorca, Spain (de Manuel, 1990); River Hase, NW Germany (Koste, 1962); Lake Dümmer, NW Germany (Koste, 1984; Koste & Poltz, 1984); Fish ponds near Warsaw, Poland

(Hillbricht-Ilkowska, 1964); Lac Léman, Fr./CH (Balvay *et al.*, 1985; Balvay & Laurent, 1988 & 1990a); Lake Derwart, near Tournay, Belgium (Bogaert, 1988; Bogaert & Dumont, 1989); Volga river, R.F. (V'yushkova & Lakhnova, 1974); Rivers in Soviet Karelia, R.F. (Filimonova, 1976); Lapland State Preserve, R.F. (Ghilarov, 1972); Lower Volga river, R.F. (Chuykov & Merkulov, 1986); All over Europe; Caucasus, Caspian area, Asia Minor (Bérziņš, 1978); Lake Quackjeswater & Breedewater (Voorne), & Lake Vogelmeer, Ymuiden, NL (Leentvaar, 1963); Lake Muy (Texel), L. oerd (Ameland) & L. Van Hunenplak (Terschelling), NL (Leentvaar, 1967); Nat. Res. Neeruti lakes, NE Estonia (Mäemets, A. & A., *s.d.*); N Swedish Lapland (Pejler, 1962a); C & S Sweden (Bérziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); 2 lakes in N Sweden (Axelson, 1961); Lake Stråken, Aneboda, Sweden (Bérziņš, 1972); Afghanistan (Bérziņš, 1961); Skopun (Heimara Halsavatn), Faroe (De Smet *et al.*, 1988); Lakes in Estonia (Haberman, 1978); Coast of S Finland (Eriksen, 1969); Sea and brackish waters in S Finland (Björklund, 1972a); River Loire, France (Lair, 1980); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Delta of river Danube, Romania (Godeanu & Zinevici, 1983; Popescu-Marinescu & Zinevici, 1967); Slovenia, ex-Yugoslavia (Markic, 1983); Croatia, ex-Yugoslavia (Erben, 1987); Asiatic Turkey (Dumont & De Ridder, 1987); Lake Erken, Sweden (Nauwerck, 1963); Karelian inland waters, R.F. (Kutikova, 1965); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Lake Ösbysjön, near Stockholm, Sweden (Pejler, 1961); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); Leżnica-Włodawa Lake district, E Poland (Kowalczyk & Radwan, 1982); 3 lakes of Sosnowickie, Poland (Radwan, 1969); Parczew area peat bogs, Poland (Radwan, 1974); Hamburg area, Germany (Rühmann, 1961); Hamburg harbour, Germany (Rühmann, 1962); Middle course of river Loire, France (Lair & Sargas, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Etang de Vaugien, Gif s/ Yvette, Essonne, France (Pourriot, 1965); Lake Doiran, Macedonia, ex-Yugoslavia (Popovska-Stankovic, 1990); Scandinavian coastal waters (Thane-Fenchel, 1968); Serbian section of river Danube (Zivkovic, 1987); All over Spain (Velasco, 1990); Rybník dam reservoir, Poland (Krzanowski, 1979); Waters near river Bodrog, NE Hungary (Dévai *et al.*, 1972/1973); Impoundment Twistetalsperrre, Nordhessen, G. (Meinel *et al.*, 1983); Clay pits in Stuivekenskerke, Belgium (Espeel, 1979); River Danube (Enaceanu & Brezeanu, 1970); River Elbe estuary, Germany (Schulz, 1961); Peat bog Chambedaze, Puy-de-Dôme, France (Francez, 1982); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Lake Iacub, river Danube delta, Romania (Godeanu, 1976); Water-courses in the Bucegi mts., Romania (Godeanu, 1963b); Ellesmere district, UK (Galliford, 1960a); Knowsley park, Liverpool, UK (Galliford, 1964); District Hoge Venen, Belgium (Gijsels *et al.*, 1976); Lakes, N of Bucarest, Romania (Godeanu, 1969b); Lakes of the Black Sea shore, Romania (Godeanu, 1970b); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Marshy zones in the Bucegi mts., Romania (Godeanu, 1969a); River Vistula in the Warsaw area, Poland (Klimowicz, 1981); Canal of river Odra, Poland (Klimowicz, 1979); Small water bodies in the Mikolajki area, Poland (Klimowicz, 1967); Side-arm of river Danube near Bratislava, CSR (Lichardova, 1964); Anatolia, Turkey (Segers *et al.*, 1992); Hilversumse wasmeer & lake in Leersumse veld, NL (Leentvaar, 1967b); Artificial lakes near Köln, Germany (Trahms, 1972); Near Debrecen, Hungary (Dévai, 1972); In periphyton of mosses and *Myriophyllum* in the Danube, and in some Alpine brooks, Austria (Donner, 1964); Shores of dammed or free flowing stretches in the Danube, at the German-Austrian border (Donner, 1972); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); Mouth areas of the Danube branches & predeltaic area of the Black Sea, Romania (Popescu-Marinescu, 1974); Lakes in NE Norway (Saeter, 1971); Dübener Heide, N Germany (Wulfert, 1960b & 1961); All over the Netherlands (Dresscher, 1976); All over Iceland (De Ridder, 1967a, 1969b, 1970 & 1972); Lake Biwa-ko, Japan (Yamamoto, 1960b); Lake Neusiedlersee, Austria (Donner, 1972d & 1979); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); 4 localities in Belgium (De Ridder, 1986); District Ostrava, CSR (Dvorakova, 1960b); C Finland (Eloranta, 1982b); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Lake Mikolajskie, Mazurian lakeland, Poland (Klimowicz, 1972); Lake Myvatn, Iceland (Adalsteinsson, 1979); Lago di Giulianello, Latium, Italy (Bazzanti & Stella, 1977); La Mancha, Castill., Spain (Armengol *et al.*, 1975); Lake Bodensee near Konstanz, Germany (Althaus, 1962); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Impoundments Sädenbach & Neunzehnhain, G. (Althaus, 1976); River Klyas'ma, R.F. (Bogoslawski, 1961); "Altes Land" near Hamburg, Germany (Caspers & Heckman, 1981 & 1982); Lake Bielersee, CH (Berner-Fankhauser, 1986); Middle course of Po river, Italy (Ferrari *et al.*, 1986); River Ybbarsån, Sweden (Bérziņš, 1979); Lake Monterosi, Latium, Italy (Stella & Margaritora, 1966); Beysehir Lake, Turkey (Emir, 1991); Peat bogs Chambedaze & la Godivelle, Massif Central, France (Francez & Dévaux, 1985); Japanese inland waters (Yamamoto, 1960a); River Grabia basin, Poland (Pawlowski, 1968, 1970 & 1973); Lake Balaton, Hungary (Zankai & Kertesz, 1967; Zankai, 1968); Canals in Hran area, CSR (Terek, 1990a); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b); Meuse Liégeoise, Belgium (Baiverlin, 1984); Akgöl, Selçuk-Izmir, Turkey (Ustaoglu & Balik, 1987); Lake Marmara (Ustaoglu, in press); Camargue, S France (De Ridder, 1960a & 1961)

- Nearctic region: Stoneman Lake, Arizona, USA (Green, 1986); Canada (Chengalath, 1977); Ontario, Canada (Chengalath, Fernando & George, 1971; Chengalath & Fernando, 1973); NE Canada (Chengalath & Koste, 1983); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); Kingston, Ontario, Canada (Nogrady, 1989); Lakes in Santee, California, USA (Nogrady, 1982); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980); Lake Wabamun, Alberta, Canada (Gallup & Hickman, 1975); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990)
- Oriental region: Sri Lanka (Mendis & Fernando, 1962; Fernando, 1969 & 1980a); Malaysia, Sri Lanka & India (Sudzuki, 1989); W & E Malaysia & Singapore (Fernando & Zankai, 1981); Sri Lanka; found in ponds (Chengalath, Fernando & Koste, 1973); Two Kumaun Himalayan Lakes, UP, India (Sharma & Pant, 1984 & 1985)
- Australian region: River Yarra, Aust. (Bērziņš, 1982); Australia (Koste & Shiel, 1989a); NSW & Qld, Aust. (Shiel & Koste, 1979)
- Neotropical region: Paraguay, Argentina & Surinam (Koste & José de Paggi, 1982); Chilean lakes (Schmid-Araya, 1991); Ecuador : Sierra (Koste & Böttger, 1989 & 1992); Argentina (Modenutti & Claps, 1988); Argentine Patagonia (Kuczynsly, 1987b); Prov. Tierra del Fuego, Chubut, Rio Negro & Neuquen, Argentina (José de Paggi, 1990); Prov. Tierra del Fuego, Argentina (Olivier, 1965)

***Trichotria pocillum bergi* (Meiszner, 1908)**

- Palaearctic region: C & E Eur. plains, Baltic province (Bērziņš, 1978); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Serbian section of river Danube (Zivkovic, 1987)

***Trichotria pocillum hudsoni* Varga, 1930**

- Palaearctic region: Alkal. lake Péteri-See, Hungary (Kertesz, 1960); Lake Balaton, Hungary (Zankai, 1968); Hungarian plain (Bērziņš, 1978)

***Trichotria pocillum majus* Varga, 1930**

- Palaearctic region: Hungarian plain (Bērziņš, 1978)

***Trichotria pseudocurta* Koste, Shiel & Tan, 1988**

- Australian region: Lake Pedder, W of Strathgordon, S Tasmania (Koste, Shiel & Tan, 1988); Tasmania (Shiel, Koste & Tan, 1989; Koste & Shiel, 1989a)

***Trichotria tetractis* (Ehrenberg, 1830)**

- Palaearctic region: Romania (Rudescu, 1960); Lake Stråken, Aneboda, Sweden (Bērziņš, 1972); Lakes in Estonia (Haberman, 1978); Coast of S Finland (Eriksen, 1969); River Loire, France (Lair, 1980); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Delta of river Danube, Romania (Godeanu & Zinevici, 1983; Zinevici & Teodorescu, 1990); Slovenia, ex-Yugoslavia (Markic, 1983); Asiatic Turkey (Dumont & De Ridder, 1987); Sierra Nevada, Spain (Morales-Baquero, 1987); C & S Sweden (Bērziņš & Pejler, 1987, 1989a, 1989b, 1989c & 1989d); Korea (Turner, 1986); Lake Bodensee near Konstanz, Germany (Althaus, 1962); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Impoundment Neunzehnhain, Germany (Althaus, 1976); Lake Bielersee, CH (Baugerter *et al.*, 1980; Berner-Fankhauser, 1986); River Klyas'ma, R.F. (Bogoslawski, 1961); Heiluntsian province, China (Bogoslawski, 1962); Lago Scuro Parmense, N Italy (Antonietti *et al.*, 1988); Lake Kisajno, Mazurian lakeland, Poland (Klimowicz, 1964); Impoundment Kninicki, CSR (Kubicek, 1964); Lake Quackjeswater, Voorne, NL (Leentvaar, 1963); N Swedish Lapland (Pejler, 1962a); Lake Lunzer Obersee, Austria (Mikschi, 1989); Alpine lakes, Sierra Nevada, Spain (Morales-Baquero *et al.*, 1989); Darnsee near Epe, NW Germany (Koste, 1962); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); River Hase, NW Germany (Koste, 1969b); Fishponds near Warsaw, Poland (Hillbright-Ilkowska, 1964); Sphagnum bogs in the Massif Central, France (Batut, 1965); Lac Léman, Fr./CH (Balvay *et al.*, 1985; Balvay & Laurent, 1989 & 1990a); Volga river & Volgograd reservoir, R.F. (V'yushkova & Lakhnova, 1974); Rivers in Soviet Karelia, R.F. (Filimonova, 1976); Aral Sea spawning grounds, R.F. (Yerekeyev & Kazakbayev, 1972); Lower Volga, R.F. (Chuykov & Merkulov, 1986); All over Europe; Caucasus, Caspian area, Asia Minor. Cosmopolitan (Bērziņš, 1978); Tingwall Loch, Mainland, Shetland (Bērziņš, 1982a); Lake Balaton, Hungary (Zankai, 1968 & 1989); District Ostrava, CSR (Dvorakova, 1960b); Karatos Lake & Gelendost, Turkey (Emir, 1991); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Japanese inland waters (Yamamoto, 1960a); ? Garden ponds, Tokyo, Japan (Sudzuki, 1975); Anatolia, Turkey (Segers *et al.*, 1992); In periphyton of mosses & *Myriophyllum*, in the Danube, and in some Alpine brooks, Austria (Donner, 1964); Shores of embanked and free flowing stretches in the Danube, at the German-Austrian border (Donner, 1972); L. Wallsee & r. Lobau, Austrian Danube basin (Donner, 1978); River Salzach & its tributaries, Austria (Donner, 1970); Dübener Heide, N Germany (Wulfert, 1960a & 1961); Peat bog La Godivelle, Massif Central, France (Francez, 1988); All over the Netherlands (Dresscher, 1976); N & W Iceland (De Ridder, 1966a, 1967a, 1969b, 1970 & 1972); Bolshoe Lake, White Sea (Ghilarov, 1965); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); Karelian inland waters, R.F. (Kutikova, 1965);

Danube delta, Romania (Popescu-Marinescu & Zinevici, 1967); Lake Montcineyre, Puy-de-Dôme, France (Persoone *et al.*, 1968); River Moldova, tributary to the Danube, Rom. (Prunescu-Arion, 1968); River Gracia basin, Poland (Pawlowski, 1960, 1968, 1970 & 1973); Canals in Hran area, CSR (Terek, 1990c); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968); Lake Donk, Overmere, Belgium (Dumont, 1968b); Marismas, Spain (De Ridder, 1962b); Pond Stifterteich, near Karlstift, lower Austria (Wawrik, 1969); In acid ponds of lower Austria (Wawrik, 1972); Southwest islands of Japan (Sudzuki *et al.*, 1992); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Waldviertel area, lower Austria (Naidenow & Wawrik, 1984); Fagne de la Mochette, Samrée, Belgium (De Ridder, 1980); In 3 localities in Belgium (De Ridder, 1986); Interstitial fauna of the Elbe estuary, Germany (Riemann, 1966); River Danube (Enaceanu & Brezeanu, 1970); River Elbe estuary, Germany (Schulz, 1961); Peat bog Chambedaze, Puy-de-Dôme, France (Francez, 1982); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Peat bog Laptici, Bucegi mountains, Romania (Godeanu, 1970a); Knowsley Park, Liverpool, UK (Galliford, 1964); Lakes of the Black Sea shore, Romania (Godeanu, 1970b); Lac Luitel, Isère, France (Joly, 1960); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960); New impoundments in the CSR (Kubicek, 1968); River Vistula in the Warsaw area, Poland (Klimowicz, 1981); Small water bodies in the Mikolajki area, Poland (Klimowicz, 1967); Side-arm of river Danube near Bratislava, CSR (Lichardova, 1964); Pool in Fontainebleau forest, Ile-de-France, Fr. (Pourriot, 1976); Pond Heinrichsweiher, Bonn, Germany (Reck, 1981); Duszatynskie Lakes, Poland (Kowalczyk & Radwan, 1962); River Loire in St. Laurent-les-Eaux, France (Pourriot *et al.*, 1982); Lake Agigea, Romania (Rodewald-Rudescu & Godeanu, 1961); 3 lakes of Sosnowickie, Poland (Radwan, 1969); Parczew area peat bogs, Poland (Radwan, 1974); Hamburg harbour, Germany (Rühmann, 1962); Middle course of river Loire, France (Lair & Sargas, 1981); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Pond and peat bogs in the Paris suburb, France (Pourriot, 1965); Peats in the Caucasus (Tarnogradsky, 1961); Pool, S of Trondheim, Norway (Thomasson, 1979); Lago Trasimeno, Italy (Taticchi, 1968); Serbian section of river Danube (Zivkovic, 1987); Ditch near Ichun, China (Wulfert, 1968); Forest pond, Lembeek, Belgium (Van Elslande, 1975); Spain : Barcelona, Granada, Huelva, Oviedo, Orense, Santander; Mallorca (Velasco, 1990); Lake Biwa-ko, Japan (Yamamoto, 1960b)

- Nearctic region: Great Dismal Swamp, border Virginia - N Carolina, USA (Anderson, Benfield & Buikema, 1977); Canada (Chengalath, 1977); Ontario, Canada (Chengalath, Fernando & George, 1971; Chengalath & Fernando, 1973); NE Canada (Chengalath & Koste, 1983); NW Canada (Chengalath & Koste, 1987); Cap Breton Island, Nova Scotia, Canada (Chengalath & Koste, 1988); Arctic N America (Chengalath & Koste, 1989); 9 mountain lakes in W Canada (Anderson & De Henau, 1980); Degg's Lake, Colorado, Canada (Pennak, 1966); Ross Lake, Florida, USA (Bienert, 1986); Cape Herschel, Ellesmere Island, NWT, Canada (Nogrady & Smol, 1989); Illinois river, USA (Brown *et al.*, 1989); Colorado, USA (Turner, 1988); Kingston, Ontario, Canada (Nogrady, 1989); Mud pond, S Grafton County, New Hampshire, USA (Wallace & Starkweather, 1983); Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); Vinton County, Ohio, USA (Evans, 1984); Lake Wabamun, Alberta, Canada (Gallup & Hickman, 1975); Atchafalaya river basin, Louisiana, USA (Holland *et al.*, 1983); Lac Echo, Québec, Canada (Nogrady, 1976); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980); Desaulniers reservoir, Québec, Canada (Pinel-Alloul *et al.*, 1982); Newfoundland lakes, Canada (Chengalath *et al.*, 1984); Triangle Lake, Portage County, Ohio, USA (Havens, 1991); James bay area, Canada (Pinel-Alloul *et al.*, 1979); Bay of Quinte, Lake Ontario, Canada (Nogrady, 1988); Walker Lake, Arizona, USA (Green, 1986); Papoose Lake, L. Jenny & Grizzly L., W North America (Thomasson, 1962); Brainard Lake & Redrock L., N Colorado, USA (Thomasson, 1964b); ? Lake 132, NW Ontario, Canada (Schindler & Novén, 1971)

- Oriental region: 2 sewage channels of Gwalior, India (Sakseena & Kulkarni, 1986); Parakrama Samudra, Sri Lanka (Fernando & Rajapaksa, 1983); Prata-pura reservoir & Nimeta water works, near Baroda, India (Wulfert, 1966); Taiwan. Cosmopolitan (Sudzuki, 1991); W & E Malaysia & Singapore (Fernando & Zankai, 1981); Indonesia, Sri Lanka, India & Philippines (Sudzuki, 1989); Kalimantan (Borneo), Indonesia (Koste, 1988a); West Bengal, NE India (Sharma, 1979a); Philippines (Mamaril & Fernando, 1978); Lake Ranu Lamongau, Indonesia (Green *et al.*, 1976); Sri Lanka (Fernando, 1980a); Singapore & Malaysia (Karunakaran & Johnson, 1978); Near Nagpur, India (Arora, 1966a, 1966c & 1966d); Punjab State, India (Sharma, 1976; Sharma & Sharma, 1984); Lakelet Tan Daha, Kathmandu valley, Nepal (Daems & Dumont, 1974); Sri Lanka : occurs in ponds (Chengalath, Fernando & Koste, 1973); Andhra Pradesh, India (Dhanapathi, 1974b); Impoundment Kinda, C Burma (Koste & Tobias, 1990); Orissa, E India (Sharma, 1980b)

- Australian region: In many localities in Australia (Bērziņš, 1982); Tasmania (Koste & Shiel, 1987); Lake Euramoo & Bromfield Swamp, tropical Queensland crater lakes (Green, 1981); Australia (Koste & Shiel, 1989a); NSW, Qld & Vic, Aust. (Shiel & Koste, 1979); Billabongs of the Murray-Darling system, Aust.

(Shiel, 1986); Lower river Murray, S Aust. (Shiel, Walker & Williams, 1982); River Murray, Aust. (Koste, 1979b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); New Zealand (Olivier, 1965); Taieri Beach, Roturna & mountain Cook, New Zealand; Chatham Islands (Russell, 1960)

- Neotropical region: Crescent Lake, Rio Snia Missu, Mato Grosso, C Brazil (Green, 1972); River Saladillo basin, Argentina (José de Paggi & Koste, 1988); River Parana, Argentina (José de Paggi, 1985); Ecuador : Oriente & Costa (Koste & Böttger, 1989 & 1992); Argentina (Modenutti & Claps, 1988); Lower Orinoco river & its delta, Venezuela (Vasquez & Rey, 1989); Lago do Macaco, Brazil (Koste, 1989a); Provinces in NW Argentina (José de Paggi, 1989); Amazonia, Brazil (Hauer, 1965; Thomasson, 1971); (Koste, 1972f); Lago Paroni, Brazil (Koste, 1974a); Varzea near Santarem, Brazil (Koste, 1974b); Argentina, Venezuela, Brazil, Chile, Surinam & Panama (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983; Koste, Robertson & Hardy, 1984); Prov. Rio Negro, Buenos Aires & Tierra del Fuego, Argentina; Chile (Olivier, 1965); Suape area, Pernambuco, Brazil (Neumann-Leitao *et al.*, 1992); Chile (Löffler, 1962); Lake Jacaretinga, Amazonia Central, Brazil (Hardy, 1980); Araucarian lakes, E of the Andes, and pools & ponds in the same region, Chile (Thomasson, 1963b); Brokopondo Lake, Surinam (Leentvaar, 1979); Rio Nhamunda, Amazonia, Brazil (Brandorff, Koste & Smirnov, 1982); Near Concepcion, Paraguay (Koste, 1986b); Island of Maraca, Roraima, Brazil (Koste & Robertson, 1990); Chilean lakes (Schmid-Araya, 1991); Argentine Patagonia (Kuczynski, 1987b); Panama (Collado *et al.*, 1984); Flooded savanna at Mantecal, Venezuela (Michelangeli *et al.*, 1979/1980)

***Trichotria tetractis* (Ehrenberg), f. *caudata* (Lucks, 1912)**

- Palaearctic region: Peat bog Chambedaze, Massif Central, France (Francez, 1987 & 1988); Lake Sträken, Aneboda, Sweden (Bérzinš, 1972); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); C & S Sweden (Bérzinš & Pejler, 1987, 1989a & 1989d); Achmer Grasmoor, NW Germany (Koste, 1968b); Engelbergs Moor, Druchhorn, NW Germany (Koste, 1965); Pontic province, C Eur. plain & Baltic province (Bérzinš, 1978); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985); Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Nat. Res. Oisterwijkse vennen & Nat. Res. De Gerritsfles, NL (Dresscher, 1976); Nat. Res. Oisterwijkse vennen, NL (De Graaf, 1960)

- Nearctic region: Cap Breton Island, Nova Scotia, Canada (Koste, 1984; Chengalath & Koste, 1988)

- Oriental region: Kerala, S India (Nayar & Nair, 1969); Taiwan (Sudzuki, 1989)

- Australian region: Not yet recorded from Australia (Koste & Shiel, 1989a)

***Trichotria tetractis* (Ehrenberg), f. *paupera* (Ehrenberg, 1830)**

- Palaearctic region: C Eur. mts., Carpathes, Pontic province, C & E Eur. plains; Caucasus (Bérzinš, 1978); Romania (Rudescu, 1960); Peat bog Laptici, Bucegi mts., Romania (Godeanu, 1970); Side-arm of river Danube, near Bratislava, CSR (Lichardova, 1964); Pelagic zone of Leżnica-Włodawa lakes, Poland (Radwan, 1975); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Serbian section of river Danube (Zivkovic, 1987)

***Trichotria tetractis* (Ehrenberg), f. *similis* (Stenoos, 1898)**

- Palaearctic region: Peat bogs in the Auvergne, France (Francez, 1981 & 1984c); Dübener Heide, N Germany (Wulfert, 1961); W Iceland (De Ridder, 1969b & 1972); In reed border of river Danube delta, Romania (Rudescu, 1968); Peat bogs Poiana Stampei, Bukowina, Rom. (Rudescu, 1966); Pool, S of Trondheim, Norway (Thomasson, 1979); Rybnik dam reservoir, Poland (Krzanowski, 1979); Romania (Rudescu, 1960); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); Croatia, ex-Yugoslavia (Erben, 1987); Lake Stechlinsee, Brandenburg, Germany (Althaus, 1963; Ronneberger & Schönborn, 1985); Lake Sträken, Aneboda, Sweden (Bérzinš, 1972); Coast of S Finland (Eriksen, 1969); Baltic province, UK, Iceland, NW Russian taiga, C Eur. mountains, Carpathes, Pontic province, C & E Eur. plains (Bérzinš, 1978); Peat bogs Chambedaze & La Godivelle, Massif Central, France (Francez & Dévaux, 1985)

- Nearctic region: NE Canada (Chengalath & Koste, 1983)

- Oriental region: W & E Malaysia & Singapore (Fernando & Zankai, 1981); Malaysia (Sudzuki, 1989); Taiwan. Cosmopolitan (Sudzuki, 1991)

- Australian region: NT, Aust.: Magela Creek area (Koste, 1981b); NT, Aust. (Shiel & Koste, 1985); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); S Tasmania (Koste, Shiel & Tan, 198)

- Neotropical region: Ecuador : Oriente & Costa (Koste & Böttger, 1992)

***Trichotria tetractis* (Ehrenberg), f. *turfacea* Petr, 1890**

- Palaearctic region: Romania (Rudescu, 1960); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); C Eur. mts., Carpathes, Pontic province, C Eur. plains, Ireland, UK, Boreal highland (Bérzinš, 1978); Inundation area of Danube delta, Romania (Dorobantu, 1964); Lower Silesia, Poland (Pawlowski, 1960); River Grabia basin, Poland (Pawlowski, 1960 & 1970)

***Trichotria truncata* (Whitelegge, 1889)**

- Palaearctic region: Lakes in NE Norway (Saeter, 1971); Dübener Heide, N Germany (Wulfert, 1960a); Umbo river system, White Sea basin, R.F. (Kruglova, 1991); Fens in the Netherlands (Prov. Drente)

(Dresscher, 1976); Luga district in the Leningrad area, R.F. (Kutikova, 1962b); River Moldova, tributary of the Danube, Rom. (Prunescu-Arion, 1968); Peat bogs Poiana Stampei, Bukowina, Rom. (Rudescu, 1966); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Romania (Rudescu, 1960); Heiluntsian province, China (Bogoslawski, 1962); Coast of S Finland (Eriksen, 1969); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); C & S Sweden (Bērziņš & Pjeler, 1987); N Swedish Lapland (Pejler, 1962a); Mt. Hoverla, Maramarosch Alps, Hungary (Varga, 1962); In bog ponds, C Europe (Koste, 1984); Small water bodies, Soviet Karelia, R.F. (Filimonova, 1976); Aral Sea spawning grounds, R.F. (Yerekeyev & Kazakhbayev, 1972); Kiev reservoir (Polishchuk, 1987); C Eur. mts., Carpathes, Hungarian plain, Pontic prov., W, C & E Eur. plains, Baltic prov., UK, Boreal highland, NW Russian taiga (Bērziņš, 1978); Nat. Res. Neeruti lakes, NE Estonia (Mäemets A. & A., s.d.); Pond Stifterteich, near Karlstift, Austria (Wawrik, 1969); In acid ponds in lower Austria (Wawrik, 1972); Japanese inland waters (Yamamoto, 1960a); Waldwinkel area, lower Austria (Naidenow & Wawrik, 1984); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973)

- Nearctic region: Mesotrophic fen, Newfoundland, Canada (Bateman & Davis, 1980); Canada (Chengalath, 1977); Illinois & Missouri, USA (Chengalath & Koste, 1983); Lac Echo, Québec, Canada (Nogrady, 1976)

- Australian region: Lake Mc Gregor & Chatham Isl., New Zealand (Russell, 1961)

***Trichotria truncata* (Whitelegge), f. *aspinosa* Rodewald, 1934**

- Palaearctic region: Water-courses in the Bucegi mts., Romania (Godeanu, 1963b); Marshy zones in the Bucegi mts., Romania (Godeanu, 1969a); Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966)

- Australian region: Lake Barracoota, Aust. (Bērziņš, 1982); NSW, Aust. (Shiel & Koste, 1979); NW Aust.: Magela Creek area (Koste, 1981b); Alligator rivers area, NT, Aust. (Tait, Shiel & Koste, 1984); Australia (Koste & Shiel, 1989a)

***Trichotria zanclum* Bērziņš, 1982**

- Australian region: Mt. Paparoa, South Island, New Zealand (Bērziņš, 1982a)

Trichotria spec.

- Palaearctic region: River Loire, France (Lair, 1980); Norfolk Broads, UK (Irvine *et al.*, 1990); Fish fattening ponds in the Ukraina (Krazhan *et al.*, 1976); Terek delta, Astrakhan peninsula, R.F. (Chuykov & Deneskin, 1985); Canal de Séclin, Nord, France (Angeli, 1972); Prov. Zuid Limburg; Nat. Res. De Gerritsfles at Kootwijk; Hilversumse Wasmeer, NL (Dresscher, 1976); Periphytic biota in sewages & park waters, Japan (Suzuki, 1981)

- Nearctic region: Mirror Lake, New Hampshire, USA (Strayer, 1985); Kalamazoo river, Michigan, USA (Prins & Davis, 1966); Long Pond, St. John's, Newfoundland, Canada (O'Connell *et al.*, 1977); The Sacramento-San Joaquin delta estuary, USA (Siegfried & Kopache, 1980)

- Oriental region: Lake Surinsar, Jammu (J & K), India (Jyoti & Sehgal, 1979)

- Neotropical region: River Parana, Argentina (Paggi & José de Paggi, 1974); Amazonia, Brazil (Hauer, 1965b; Thomasson, 1971)

***Tripleuchlanis* Myers, 1930**

***Tripleuchlanis plicata* (Levander, 1894)**

- Palaearctic region: Gulf of Finland & Gulf of Riga (Bērziņš, 1960); Camargue, S France (De Ridder, 1960a & 1961); Marismas, Spain (De Ridder, 1962); Southwest islands of Japan (Suzuki *et al.*, 1992); Balearic Archipelago, Spain (de Manuel *et al.*, 1992); Obretin embanked basin, Danube delta, Rom. (Godeanu *et al.*, 1973); Anatolia, Turkey (Segers *et al.*, 1992); Brackish polder ditches, NL (Oosten, 1972); River Moldova, tributary to the Danube, Rom. (Prunescu-Arion, 1968); Pelagic zone of the Leczna-Wladawa lakes, P. (Radwan, 1975); Parczew area peat bogs, Poland (Radwan, 1974); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Spain : Valencia & Huelva (Velasco, 1990); Romania (Rudescu, 1960); Lake Palaeostomi, Georgia, R.F. (Haberland & Laugaste, 1982); Sea and brackish water pools in S Finland (Björklund, 1972a); Coast of S Finland (Eriksen, 1969); Asiatic Turkey (Dumont & De Ridder, 1987); Volga river, R.F. (V'yushkova & Lakhnova, 1974); All over Europe & Caspian area (Bērziņš, 1978); Rügener, Hiddensee & Darszer Boddengewässer, NE Germany (Schwarz, 1963); Obretin Lake, Danube delta, Romania (Neagu-Godeanu *et al.*, 1968)

- Nearctic region: Illinois, USA (Bērziņš, 1984); Experimental ponds near Bakersfield, Kern County, California (Hurlbert *et al.*, 1972)

- Oriental region: West Bengal, NE India (Sharma, 1979b); Bangkok & pool near river Kwai at Kanchanaburi, Thailand (De Ridder, 1971); Thailand (Boonsom, 1984b); Philippines (Mamaril & Fernando, 1978); Sri Lanka (Fernando, 1980a); Taiwan (Suzuki, 1988); Taiwan, Thailand, Indonesia, Sri Lanka, India, Philippines (Suzuki, 1989); Sri Lanka, rare. Found in ponds (Chengalath, Fernando & Koste, 1973);

Taiwan. Warm waters (Sudzuki, 1991); Municipal tank, Kaikalur, India (Dhanapathi, 1974b)
 - Australian region: Creek in Australia (Bērziņš, 1982); Australia (Koste & Shiel, 1989a); Qld, Aust. (Shiel & Koste, 1979)

- Neotropical region: Antilles françaises (Pourriot, 1975); Amazonia, Brazil (Thomasson, 1971); Ecuador : Oriente, Costa & Sierra (Koste & Böttger, 1989); Orinoco river basin (Vasquez, 1984a); Venezuela & Antilles (Koste & José de Paggi, 1982); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983); Caribbean section of the Neotropical region (De Ridder, 1977); Aruba, Bonaire, Curaçao, Caribbean Islands, Cuba, Haiti, Panama (Collado *et al.*, 1984)

Tripleuchlanis plicata (Levander), f. *razelmi* Rodewald, 1940

- Palaearctic region: Pontic province (Bērziņš, 1978); Lakes of the Black Sea shore, Romania (Godeanu, 1970b); Lake Agigea, Romania (Rodewald-Rudescu & Godeanu, 1961); River Volga, R.F. (Mordukhai-Boltovskoi, 1979)

Tripleuchlanis spec.

- Oriental region: Fishponds in Thailand (Boonsom, 1984a)
- Neotropical region: Province San Juan, Argentina (José de Paggi, 1990)

Trochosphaera Semper, 1972

Trochosphaera aequatorialis Semper, 1872

- Palaearctic region: Delta of river Danube, Romania (Zinevici & Teodorescu, 1991)
- Nearctic region: Lake Wabamun, Alberta, Canada (Gallup & Hickman, 1975)
- Oriental region: Sri Lanka (Fernando, 1969 & 1980a; Serruya & Pollingher, 1984); Sri Lanka & Philippines (Sudzuki, 1989); Sri Lanka. Only a single record from Senanayake Samudra (Chengalath, Fernando & Koste, 1983); Fishponds in Thailand (Boonsom, 1984a); Thailand (Boonsom, 1984b)
- Neotropical region: Surinam (Leentvaar, 1975); S America : Surinam & Amazonia (Koste, 1976a); Surinam, Brazil & Argentina (Koste & José de Paggi, 1982); Flood valley of river Parana, Argentina (Paggi, 1978); Prov. Santa Fé, Argentina (José de Paggi, 1990)

Trochosphaera solstitialis Thorpe, 1893

- Palaearctic region: SE Europe (Koste, 1967a); E Balkan & Pontic prov. Tropical & subtropical (Bērziņš, 1978); Lake Iacub, river Danube delta, Romania (Godeanu, 1976)
- Nearctic region: N America (Koste, 1976a); Kingston, Ontario, Canada (Nogrady, 1989); Temporary pond in the Stephen F. Austin experimental forest, 15 km SW of Nacagoloches, Texas, USA (Mc Cullaugh & Lee, 1980)
- Oriental region: E Asia (Koste, 1976a)

Trichosphaera spec.

- Oriental region: In rice-fields of N Kriaú, Perak, Malaysia (Ali, 1990)
- Neotropical region: Surinam (Leentvaar, 1975); River Parana, Argentina (José de paggi, 1985); Lago Camaleão, Amazonia, Brazil (Koste & Robertson, 1983)

Tylotrocha Harring & Myers, 1922

Tylotrocha monopus (Jennings, 1894)

- Palaearctic region: Lake Ladoga, R.F. (Nikolayev & Petrova, 1978); Romania (Rudescu, 1960); Lake Balaton, Hungary (Zankai, 1968); Karelian inland waters, R.F. (Kutikova, 1965); Hungarian plain, W Eur. plain, Baltic province, NW Russian taiga (Bērziņš, 1978)
- Nearctic region: Laurentian Great Lakes (Stemberger, 1979); Northern Michigan inland lakes (Stemberger, 1990); NE Canada (Chengalath & Koste, 1983); Lac Echo, Québec, Canada (Nogrady, 1976); Mt. Tremblant Prov. park, Québec, Canada (Nogrady, 1980)

Wierzejskiella Wiszniewski, 1934

Wierzejskiella elongata (Wiszniewski, 1932)

- Palaearctic region: River Hase, NW Germany (Koste, 1976b); Courland Spit, R.F. (Zhadin *et al.*, 1972); Peat bogs Poiana Stampei, Bukowina, Rom. (Rudescu, 1966); Carpathes, C & E Eur. plains, Ireland (Bērziņš, 1978)

- Nearctic region: N America (Bērziņš, 1978)

Wierzejskiella marina Remane, 1949

- Palaearctic region: C Eur. plain (Bērziņš, 1978)

Wierzejskiella ricciae (Harring, 1913)

- Palaearctic region: River Hase, NW Germany (Koste, 1976b); S Sweden & C Eur. plain (Bērziņš, 1978)

- Nearctic region: N America (Bērziņš, 1978)

Wierzejskiella sabulosa (Wiszniewski, 1932)

- Palaearctic region: Romania (Rudescu, 1960); Shores of Black Sea, C & E Eur. plains (Bērziņš, 1978); Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986)

Wierzejskiella subterranea Remane, 1949

- Palaearctic region: C Eur. plain (Bērziņš, 1978)

Wierzejskiella vagneri Koniar, 1935

- Palaearctic region: Peat bog Laptici, Bucegi mountains, Romania (Godeanu, 1970); Carpathes (Bērziņš, 1978)

Wierzejskiella velox (Wiszniewski, 1932)

- Palaearctic region: Lake Vörtsjärv, Estonia (Kutikova & Haberman, 1986); River Hase, NW Germany (Koste, 1976b); Peat bogs in the Massif Central, France (Batut, 1965); Italy, W Balkan, Carpathes, C & E Eur. plains (Bērziņš, 1978); Lake Ohrid, Balkan peninsula (Stankovic, 1960)

- Nearctic region: N America (Bērziņš, 1978); Laurentian Great Lakes (Stemberger, 1979); NE Canada (Chengalath & Koste, 1983)

- Australian region: SW Australia (Koste, Shiel & Brock, 1983); W Australia (Shiel & Koste, 1985)

Wierzejskiella spec.

- Palaearctic region: 3 different species in mt. Fuji lava caves, Japan (Sudzuki, 1971a); Lakes around mountain Fuji, Japan (Sudzuki, 1971b); Mountain Fuji area, Japan (Sudzuki, 1978a); Iriomote Island, Japan (Sudzuki, 1979); ? Interstitial fauna of the Elbe estuary, Germany (Riemann, 1966)

- Nearctic region: 2 different species in Vinton County, Ohio, USA (Evans, 1984); Mirror Lake, New Hampshire, USA (Strayer, 1985)

- Oriental region: Malaysia (Sudzuki, 1989)

Wigrella Wiszniewski, 1932

Wigrella amphora (Remane, 1929)

- Palaearctic region: C Eur. plain (Bērziņš, 1978)

Wigrella depressa Wiszniewski, 1932

- Palaearctic region: Peat bogs Poiana Stampei, Bukowina, Romania (Rudescu, 1966); Carpathes, C & E Eur. plains (Bērziņš, 1978)

Wigrella spec.

- Palaearctic region: In an inland sea of Japan, round Kasado Island (Sudzuki, 1976a & 1976b)

- Nearctic region: The Sacramento-San Joaquin delta estuary, USA (Siegfried & Kopache, 1980)

Wolga Skorikov, 1903

Wolga spinifera (Western, 1894)

- Palaearctic region: Heiluntsian province, China (Bogoslowski, 1962); Asiatic Turkey (Dumont & De Ridder, 1987); Volga & Volgograd reservoir, R.F. (V'yushkova & Lakhnova, 1974); Almost all over Europe (Bērziņš, 1978); Latorica, E Slovakia lowland, CSR (Terek, 1990a & 1990b); Side-arm of river Danube near Bratislava, CSR (Lichardova, 1964); Anatolia, Turkey (Segers *et al.*, 1992); Luga district of the Leningrad area, R.F. (Kutikova, 1962b); River Volga, R.F. (Mordukhai-Boltovskoi, 1979); Serbian section of river Danube (Zivkovic, 1987); Pond Chylinski Mali II, Poland (Lewkowicz, 1972); Spain : Guadalajara, Huesca, Jaen & Pamplona (Velasco, 1990)

- Nearctic region: Canada (Chengalath, 1977); Illinois & Kentucky, USA (Bērziņš, 1984); NW Canada (Chengalath & Koste, 1987); Arctic N America (Chengalath & Koste, 1989); Laurentian Great Lakes

(Stemberger, 1979); Bay of Quinte, Lake Ontario, Canada (Nogrady, 1988)

- Oriental region: Yamuna river, near Wazirabad, Delhi, India (Sarma, 1988)

- Australian region: Tasmania (Koste & Shiel, 1987); NSW, Aust. (Koste & Shiel, 1980c)

- Neotropical region: Argentina (José de Paggi, 1982; Koste & José de Paggi, 1982; Modenutti & Claps, 1988); Provinces in NW Argentina (José de Paggi, 1989); Prov. Chubut, Tucuman, Cordoba, Santa Fé & Buenos Aires, Argentina (José de Paggi, 1990); Argentine Patagonia (Kuczynski, 1987b)

Wulfertia Donner, 1943

Wulfertia kindensis Koste & Tobias, 1990

- Oriental region: Impoundment Kinda, C Burma (Koste & Tobias, 1990)

Wulfertia ornata Donner, 1943

- Palaearctic region: Romania (Rudescu, 1960); C Eur. mountains & plains, Carpathes (Bērziņš, 1978); Dübener Heide, N Germany (Wulfert, 1960b)

Wulfertia spec.

- Antarctic region: Antarctica & Subantarctica (José de Paggi & Koste, 1984); Antarctica (Dartnall & Hollowday, 1985)

References

- ACKEFORS, H., 1969. Seasonal and vertical distribution of the zooplankton in the Askö area (Northern Baltic proper) in relation to hydrographical conditions. *Oikos* 20: 480-492
- ADALSTEINSSON, H., 1974. Plankton i Myvatn. *Scripta Limnol. Upsal.* 372. 46pp.
- ADALSTEINSSON, H., 1979. Zooplankton and its relation to available food in Lake Myvatn. *Oikos* 32: 162-194
- ADAMKIEWICZ-CHOJNACKA, B & Z. ROZANSKA, 1990. Seasonal variation of the biomass of Copepoda and Rotatoria: permanent components of the zooplankton of the Vistula lagoon. *Limnologica (B)* 20: 71-73
- AFANAS'JEVA, E.L., 1975. Littoral zooplankton in the open zone of Lake Baikal. *Hydrobiol. Jnl.* 11 (3): 19-23
- AGUESSE, P., 1960. Observations floristiques et faunistiques sur un étang de Moyenne Camargue: la Basse Salée de la Tour de Valat. *Vie et Milieu XI*: 284-307
- AL'KAMM, K.S., 1984. New Rotifer species from the Black Sea. *Hydrobiol. Jnl.* 20 (6): 39-43
- ALBERTOVA, D., 1960. Eine neue Art der Gattung *Polyarthra* (Rotatoria). *Acta Soc. zool. Bohemoslov.* 24: 16-18
- ALEKSANDROV, V. V., & L. A. KUTIKOVA, 1985. Connections between Diel Migration of Zooplankton and fluctuations in the Earth's Electrical and Magnetic Fields. *Hydrobiol. Jnl.* 21 (4): 59-69
- ALI, A. B., 1990. Seasonal dynamics of microcrustacean and rotifer communities in Malaysian ricefields used for rice-fish farming. *Hydrobiologia* 206: 139-148
- ALLEN, A. A., 1968. Morphology of the planktonic rotifer *Polyarthra vulgaris*. *Trans. amer. microscop. Soc.* 87: 60-69
- ALONSO, M., 1984. Crustaceos y rotiferos de la laguna del Tesoro (Cuba). *Limnetica* 1: 70-77
- ALTHAUS (B. KOCH-), 1961. *Cephalodella theodora*, ein neues Rotator aus dem Stechlin-see. *Zool. Anz.* 166: 219-221
- ALTHAUS (B. KOCH-), 1962a. Weitere neue und bemerkenswerte Rotatorien aus dem Stechlinsee. *Limnologica* 1: 65-82
- ALTHAUS (B. KOCH-), 1962b. Beitrag zur Rotatorienfauna des Bodensees. *Limnologica* 1: 93-110
- ALTHAUS (B. KOCH-), 1963. Systematische und ökologische Studien an Rotatorien des Stechlinsees. *Limnologica* 1: 375-456
- ALTHAUS (B. KOCH-), 1976. Die Rotatorienfauna der Saidenbachtalsperre und der oberen Neunzehnhainer Talsperre. *Limnologica* 11: 17-46
- AMREN, H., 1964a. Temporal variation of the Rotifer *Keratella quadrata* (Müller) in some ponds on Spitsbergen. *Zool. Bidr. Uppsala* 36: 193-208
- AMREN, H., 1964b. Studies on the ecology and taxonomy of zooplankton, especially the Rotifers from Spitsbergen. *Acta Univ. Uppsala* 42: 5-10
- AMREN, H., 1964c. Ecological studies of zooplankton populations in some ponds on Spitsbergen. *Zool. Bidr. Uppsala*, 36: 161-191
- AMREN, H., 1964d. Ecological and taxonomical studies on zooplankton from Spitsbergen. *Zool. Bidr. Uppsala* 36: 209-276
- ANDERSON, K. B., F. BENFIELD & A. L. BUIKEMA, 1977. Zooplankton of a swamp water ecosystem. *Hydrobiologia* 55: 177-185
- ANDERSON, R. S., 1968. The zooplankton of five small mountain lakes in south-western Alberta. *Nat. Mus. Can. Nat. Hist. Paper* n° 39. 18pp.
- ANDERSON, R. S., 1970a. Predator-prey relationships and predation rates for crustacean zooplankters from some lakes in western Canada. *Can. J. Zool.* 48: 1229-1240
- ANDERSON, R. S., 1970b. Effects of Rotenone on zooplankton communities and a story of their recovery in two Mountain Lakes in Alberta. *J. Fish. Res. Bd. Canada* 27: 1335-1336
- ANDERSON, R. S., 1972. Zooplankton composition and change in an alpine lake. *Verh. intern. Ver. Limnol.* 18: 264-268
- ANDERSON, R. S., 1977. Rotifer populations in mountain lakes in relation to fish and species of copepoda present. *Arch. Hydrobiol. Beih.* 8: 130-134
- ANDERSON, R. S., 1980. Relationships between trout and invertebrate species as predators and the structure of the crustacean and rotiferan plankton in mountain lakes. in W. C. Kerfoot, 1980: *Evolution and ecology of plankton communities*. The University Press of New England
- ANDERSON, R. S., & A. M. DE HENAU, 1980. An assessment of the meiobenthos from nine mountain lakes in Western Canada. *Hydrobiologia* 70: 257-264

- ANDERSON, R. S., & R. B. GREEN, 1975. Zooplankton and Phytoplankton studies in the Waterton Lakes, Alberta, Canada. *Verh. intern. Ver. Limnol.* 19: 571-579
- ANDERSON, R. S., & R. B. GREEN, 1976. Limnological and planktonic studies in the Waterton Lakes, Alberta. *Can. Wildlife Serv. Occas. Paper n° 27.* 39pp.
- ANDERSSON, G., H. BERGGREN & S. HAMRIN, 1975. Lake Trummen restoration project III. Zooplankton, macrobenthos and fish. *Verh. intern. Ver. Limnol.* 19: 1097-1106
- ANDREW, T. E., S. CABRERA & V. MONTECINO, 1989. Diurnal changes in zooplankton respiration rates and the phytoplankton activity in two Chilean lakes. *Hydrobiologia* 175: 121-135
- ANDREW, T. E. & A. G. FITZSIMONS, 1992. Seasonality, population dynamics and production of planktonic rotifers in Lough Neagh, Northern Ireland. *Hydrobiologia* 246: 147-164
- ANGELI, N., 1972. Esquisse du profil biologique (plancton) d'un canal influencé par une pollution complexe d'origine industrielle: le canal de Séclin-Nord. *Biol. Jb. Dodonaea* 40: 77-115
- ANGELI, N., 1975. Migrations verticales: quelques données sur le comportement automnal du zooplancton d'un canal de navigation. *Annls Limnol.* 11: 239-253
- ANSELLEM, J. & P. CLEMENT, 1988. The muscle of a monogont Rotifer, *Trichocerca rattus*, II. The central retractor muscles. *Tissue and cell* 20: 89-108
- ANTONIETTI, R., I. FERRARI, G. ROSSETTI, L. TAROZZI & P. VIAROLI, 1988. Zooplankton structure in an oligotrophic mountain lake in Northern Italy. *Verh. intern. Ver. Limnol.* 23: 545-552
- ARANJO-LIMA, C. A. R. M. & E. HARDY, 1987. Aspectos biológicos de peixes amazônicos, VIII. A alimentação dos alevinos do Jaraqui, *Semiprochilodus insignis*. *Amazoniana*, 10: 127-136
- ARCIFA, M. S., 1984. Zooplankton composition of ten reservoirs in southern Brazil. *Hydrobiologia* 113: 137-145
- ARCIFA, M. S., E. A. T. GOMES & A. J. MESSCHIATTI, 1992. Composition and fluctuations of the zooplankton of a tropical Brazilian reservoir. *Arch. Hydrobiol.* 123: 479-494
- ARMENGOL, J., 1982. Ecología del zooplancton de los embalses. *Mundo Científico (La Recherche)* 11: 168-179
- ARMENGOL, J., N. E. ESTRADA, A. GUISET, R. MARGALEF, D. PLANAS, J. TOYA & F. VALLESPINOS, 1975. Observaciones limnológicas en las lagunas de la Mancha. *Boletín de la Estación de Ecología IV*, n°8, Madrid: 11-27
- ARMENGOL, J. G. MOREAU & D. PLANAS, 1988. Evolution, à court terme, des communautés planctoniques de deux rivières du nord Québécois, soumises à une forte réduction de débit. *Can. J. Biol.* 61: 2011-2020
- ARMENGOL, J., F. SABATER, J. L. RIERA & M. A PUIG, 1988. Longitudinal changes in the zooplankton communities along a series of reservoirs in the Guadiana River (W Spain). *Verh. intern. Ver. Limnol.* 23: 1006-1010
- ARMITAGE, K. B. & H. B. HOUSE, 1962. A limnological reconnaissance in the area of Murdo Sound, Antarctica. *Limnol. Oceanogr.* 7: 36-41
- ARNDT, H., 1991. Population ecology of estuarine rotifers in an inlet of the Southern Baltic. VI *International Rotifer Symposium - Book of Abstracts*: 83
- ARNDT, H., I. DEBUS, R. HEERKLOSS & W. SCHULTE, 1984. Diurnal changes in the matter flux of a shallow-water system in a Baltic inlet. *Ophelia, suppl.* 3: 1-9
- ARNDT, H. & R. HEERKLOSS, 1989. Diurnal variations in feeding and assimilation rates of planktonic rotifers and its possible ecological significance. *Int. Revue ges. Hydrobiol.* 74: 261-272
- ARNDT, H., R. HEERKLOSS & W. SCHNESE, 1984. Seasonal and spatial fluctuations of estuarine rotifers in a Baltic inlet. *Limnologica* 15: 377-385
- ARNDT, H., W. SCHNESE & R. HEERKLOSS, 1981. Untersuchungen zur räumlichen Verteilung des Zooplanktons in einem flachen inneren Küstengewässer der Ostsee. *Wiss. Z. Univ. Rostock, Math.-nat. Reihe* 30 (4/5): 43-48
- ARNDT, H., C. SCHRÖDER & W. SCHNESE, 1990. Rotifers of the genus *Synchaeta* - an important component of the zooplankton in the coastal waters of the Southern Baltic. *Limnologica* 21: 233-235
- ARNEMO, R., B. BĒRZIŅŠ, B. GRÖNBERG & I. MELLGREN, 1968. The dispersal in Swedish waters of *Kellicottia bostoniensis* (Rousselet) (Rotatoria). *Oikos* 19: 351-358
- ARORA, H. C., 1962a. Studies on Indian Rotifera. Part I. On a small collection of illoricate Rotifera from Nagpur, India, with notes on their bionomics. *J. Zool. Soc. India* 14: 33-44
- ARORA, H. C., 1962b. Studies on Indian Rotifera. Part II. On some species of the genus *Brachionus* from Nagpur, India. *J. Zool. Soc. India* 15: 112-121
- ARORA, H. C., 1963. Studies on Indian Rotifera. Part IV. On some species of sessile Rotifera from India. *Arch. Hydrobiol.* 59: 502-507

- ARORA, H.C., 1965. Studies on Indian Rotifera. Part VI. On a collection of Rotifera from Nagpur, India, with four new species and a new variety. *Hydrobiologia* 26: 444-456
- ARORA, H. C., 1966a. Studies on Indian Rotifera. Part V. On species of some genera of the family Brachionidae, subfam. Brachioninae, from India. *Arch. Hydrobiol.* 61: 482-493
- ARORA, H. C., 1966b. Cyclomorphosis (form variation) in some species of Indian planktonic Rotatoria. *Int. Revue ges. Hydrobiol.* 51: 623-632
- ARORA H. C., 1966c. Rotifera as indicators of trophic nature of environments. *Hydrobiologia* 27: 146-159
- ARORA, H. C., 1966d. Responses of Rotifera to variations in some ecological factors. *Proc. Indian Acad. Sci. Sect. B* 63: 57-66
- ARORA, H. C., 1967. Photodynamic action of light on Rotifera. *J. zool. Soc. India* 17: 37-47
- ARVOLA, L., T.-R. METSÄLÄ, A. SIMILÄ & M. RASK, 1990. Phyto- and zooplankton in relation to water pH and humic content in small lakes in Southern Finland. *Ver. intern. Ver. Limnol.* 24: 688-692
- ARVOLA, L., K. SALONEN, I. BERGSTRÖM, A. HEINÄNEN & A. OJALA, 1986. Effects of experimental acidification on phyto-, bacterio- and zooplankton in enclosures of a highly humic lake. *Int. Revue ges. Hydrobiol.* 71: 737-758
- ARVOLA, L., K. SALONEN, R. I. JONES, A. HEIMÄNEN & I. BERGSTRÖM, 1987. A three day study of the diel behaviour of plankton in a highly humic and deeply stratified lake. *Arch. Hydrobiol.* 109: 89-106
- AXELSON, J., 1961. Zooplankton and impoundment of two lakes in Northern Sweden (Ransaren and Kultsjön). *Rep. Inst. Freshw. Res. Drott.* 42: 84-167
- BADALAMENTE, M. & S. H. VERNICK, 1973. The ultrastructure of *Bertramia asperospora* Fritsch, a microsporidan parasite on the Rotifer *Brachionus plicatilis* Müller. *Arch. Protistenk.* 115: 1-9
- BADINO, G. & C. ROBOTTI, 1975. Selection in parthenogenetic lines of *Asplanchna sieboldi* (Leydig, 1854) (Rot.). *Experientia* 31: 298-299
- BAIVERLIN, P., 1984. Etude des Rotifères planctoniques de la Meuse en relation avec le complexe nucléaire de Tihange et l'agglomération Liégeoise. *Mémoire de licence, Sci. zool.*, Univ. Liège. 57pp.
- BAKER, R. L., 1979a. Birth rate of planktonic rotifers in relation to food concentration in a shallow, eutrophic lake in Western Canada. *Can. J. Zool.* 57: 1206-1214
- BAKER, R. L., 1979b. Specific status of *Keratella cochlearis* (Gosse) and *K. earlinae* Ahlstrom (Rotifera: Brachionidae): morphological and ecological considerations *Can. J. Zool.* 57: 1719-1722
- BAKKER, C., 1964. Planktonuntersuchungen in einem holländischen Meeresarm vor und nach der Abdeichung. *Helgol. wiss. Meeresunters.* 10: 456-472
- BAKKER, C. & N. DE PAUW, 1975. Comparison of plankton assemblages of identical salinity ranges in estuarine tidal, and stagnant environments. II. Zooplankton. *Neth. J. Sea Res.* 9: 145-165
- BALSEIRO, E. G. & B. E. MODENUTTI, 1990. Zooplankton dynamics of Lake Escondido (Rio Negro, Argentina), with special reference to a population of *Boeckelia gracilipes* (Copepoda, Calanoidea). *Int. Revue ges. Hydrobiol.* 75: 475-491
- BALVAY, G., 1972. L'évolution de la biocénose planctonique du lac d'Annecy. *Ann. Hydrobiol.* 3: 93-116
- BALVAY, G., 1984. Zooplankton, in Feuillade, J., ed.: Caractérisation et essais de restauration d'un écosystème dégradé: le lac de Nantua. *INRA publ.* Paris. 165pp.
- BALVAY, G., 1989. Evolution de la biocénose rotatorienne au cours des variations de l'état trophique du Léman et comparaison avec le lac de Constance. *Rev. sci. de l'eau* 2: 739-753
- BALVAY, G., 1991. What is the best reference to express biological results? The calendar year or the ecological year? *Verh. int. Ver. Limnol.* 24: 928-930
- BALVAY, G. & J.-C. DRUART, 1992. Le plancton du lac d'Annecy: historique et inventaire. *Arch. Sci. Genève* 45: 135-169
- BALVAY, G. , J.-C. DRUART & M. LAURENT, 1985. Premier complément à l'inventaire de la biocénose du Lac Léman. *Schweiz. Z. Hydrol.* 47: 76-80
- BALVAY, G. & M. LAURENT, 1981. The Rotifers of Lake Geneva (Lake Léman). *Schweiz. Z. Hydrol.* 43: 126-139
- BALVAY, G. & M. LAURENT, 1989. Incidences des variations de l'état trophique sur les caractéristiques de la biocénose planctonique rotatorienne du Lac Léman. *Arch. Sci. Genève* 42: 341-365
- BALVAY, G. & M. LAURENT, 1990a. Evolution qualitative à long terme des rotifères du Lac Léman. *Aquatic Science* 52: 156-161
- BALVAY, G. & M. LAURENT, 1990b. Evolution quantitative à long terme des rotifères au cours de l'eutrophisation du Lac Léman. *Aquatic Science* 52: 162-175
- BARRON, G. L. & E. SZIJARTO, 1990. A new genus of hyphomycetes endoparasitic in rotifers. *Mycologia* 82: 134

- BAUGERTER, B. R., E. BÖLEN & H. FANKHAUSER, 1980. Untersuchungen über das pelagische Phyto-, Rotatorien- und Crustacean- Plankton im Bielersee. *Lic. Thesis Zool. Inst. Un. Bern, Schweiz.*
- BARTHELMES, D., 1960. Die horizontale Planktonverteilung im Groszen Müggelsee. I. Einige Beobachtungen über die Verhältnisse im Eulimnion. *Z. Fisch. Hilfswiss. IX, N. F.*, H. 5/6: 333-352
- BATEMAN, L. E. & CH. C. DAVIS, 1980. The rotifers of Hummock-Hollow Formations in a poor (mesotrophic) Fen in New-foundland. *Int. Rev. ges. Hydrobiol.* 65: 127-153
- BATTES, K., I. CARAUSU, C. MAZAREANU, I. MIRON, R. RUJINSCHI, C. RUJINSCHI, FL. SIMALESIK & F. R. SIMALESIK, 1971. Hydrobiological studies upon Bicaz water power reservoir during 1970. *Lucr. Stat. "Stearul"* 4: 375-398
- BATUT, J., 1965. Etude de la faune submicroscopique de quelques tourbières à *Sphagnum*. *Hydrobiologia* 25: 239-276
- BAYLY, I. A. E., 1967. The fauna and chemical composition of some athalassic saline waters in New Zealand. *N. Z. J. Mar. Freshwat. Res.* I: 105-117
- BAYLY, I. A. E. & W. D. WILLIAMS, 1966. Chemical and biological studies on some saline lakes of SE Australia. *Austral. J. Mar. Freshw. Res.* 17: 177-228
- BAZZANTI, M. & E. STELLA, 1977. Popolamento teutonico, in Alfinito, S et al.: Indagini eco-logistiche sul Lago di Giuliancello (Lazio). *Annali di botanica* 35-36 (1976-1977): 314-331
- BEACH, N. W., 1960. A study of the planctonic rotifers of the Ocqueoc River system, Presque Isle County, Michigan. *Ecol. Monogr.* 30: 339-357
- BEAVER, J. R. & T. L. CRISMAN, 1990. Use of microzooplankton as an early indicator of advancing cultural eutrophication. *Verh. intern. Ver. Limnol.* 24: 532-537
- BENNETT, W. N. & M. E. BORAAS, 1989. Comparison of population dynamics between slow-growing and fast-growing strains of the Rotifer *Brachionus calyciflorus* Pallas in continuous culture. *Oecologia* 81: 494-500
- BERGHahn, R., 1979. Schädigung limnischer Rotatorien bei der Passage durch das Kühlssystem eines Kohlekraftwerkes. *Arch. Hydrobiol. / Suppl.* 43: 225-235
- BERMAN, TH., U. POLLINGHER & M. GOPHEN, 1972. Lake Kinneret: planktonic populations during seasons of high and low phosphorus availability. *Verh. intern. Ver. Limnol.* 18: 588-598
- BERNER-FANKHAUSER, H., 1983. Abundance, dynamics and succession of planktonic rotifers in Lake Biel, Switzerland. *Hydrobiologia* 104: 349-352
- BERNER-FANKHAUSER, H., 1986. Populationsdynamik der Plankton-Rotatorien im Bielersee, 1981/1982. *Doctor's thesis Un. Bern, CH*
- BERNER-FANKHAUSER, H., 1987. The influence of sampling strategy on the apparent population dynamics of plankton rotifers. *Hydrobiologia* 147: 181-188
- BĒRZINŠ, B., 1960. Rotatoria. *Conseil intern. Explor. Mer. Zooplankton.* Sheets 84-88
- BĒRZINŠ, B., 1961a. New Rotatoria (Rotifera) from Victoria, Australia. *Proc. Roy. Soc. Victoria* 74: 83-86
- BĒRZINŠ, B., 1961b. Two new *Keratella* subspecies from Canada. *J. Quek. micr. Club*, ser. 4, vol. V: 437-438
- BĒRZINŠ, B., 1961c. Rotatoria. Contribution à l'étude de la faune d'Afghanistan, 23. *Förhandl. Fysiogr. Sällsk. i Lund* 31, n° 2: 6-16
- BĒRZINŠ, B., 1962. Revision der Gattung *Anuraeopsis* Lauterborn (Rot.). *K. fysiogr. Sällsk. Lund Förh.* 32: 33-47
- BĒRZINŠ, B., 1963. Two new *Keratella*, Rotatoria, from Australia. *Hydrobiologia* 21: 380-383
- BĒRZINŠ, B., 1972. Eine Mikrobenthosstudie an Rotatorien. *Hydrobiologia* 40: 447-542
- BĒRZINŠ, B., 1973. Eine neue *Trichocerca*-Art, Rotatoria, aus Neuseeland. *Zool. Anz.* 190: 123-124
- BĒRZINŠ, B., 1973a. Some Rotifers from Cambodia. *Hydrobiologia* 41: 453-459
- BĒRZINŠ, B., 1976. Notes on the Rotifers from Aneboda, Sweden. Lund, nov. 1976. 16pp
- BĒRZINŠ, B., 1978. Rotatoria in J. Illies: Limnofauna Europeae: 54-91. *G. Fischer Verlag, Stuttgart & New York*, 1978
- BĒRZINŠ, B., 1979. Limnological investigations in the river Ybbarspå, 1979. Periphyton and plankton. Private ed. *Inst. Limnol. Lund*. 42 pp.
- BĒRZINŠ, B., 1982. Contribution to the knowledge of Rotatoria of Australia. Private ed. *Inst. Limnol. Lund*. 24pp.
- BĒRZINŠ, B., 1982a. Short notes on rotatoria. Private ed. *Inst. Limnol. Lund*. 8 pp.
- BĒRZINŠ, B., 1984. Some Rotifers from North America: a reminiscence. Private ed. *Inst. Limnol. Lund*. 13 pp.
- BĒRZINŠ, B. & B. PEJLER, 1987. Rotifer occurrence in relation to pH. *Hydrobiologia* 147: 107-116

- BĚRZINŠ, B. & B. PEJLER, 1989a. Rotifer occurrence in relation to temperature. *Hydrobiologia* 175: 223-231
- BĚRZINŠ, B. & B. PEJLER, 1989b. Rotifer occurrence and trophic degree. *Hydrobiologia* 182: 171-180
- BĚRZINŠ, B. & B. PEJLER, 1989c. Rotifer occurrence in relation to oxygen content. *Hydrobiologia* 183: 165-172
- BĚRZINŠ, B. & B. PEJLER, 1989d. Rotifer occurrence in relation to water colour. *Hydrobiologia* 184: 23-28
- BEUSCHOLD, E., 1961. Limnische Untersuchungen am Hauptbecken der Säidenbach-Talsperre. *Int. Revue ges. Hydrobiol.* 46: 18-42
- BICK, H. & E. SCHOLTYSECK, 1960. Ökologische Untersuchungen an Abwasserteichen. Beitrag zur Oekologie der Planktonorganismen aus Abwasserteichen. *Arch. Hydrobiol.* 57: 196-216
- BIENERT, R. W. Jr., 1986. A new species of *Lecane* (Rotifera - Lecanidae) from subtropical Florida. *Hydrobiologia* 141: 175-177
- BILIO, M., 1966. Die aquatische Bodenfauna von Salzwiesen der Nord- und Ostsee. II. Ökologische Faunen-analyse: Hydrozoa, Nematodes, Rotatoria, Gastrotricha, Nemertini, Polychaeta, Oligochaeta, Halacaridae, Ostracoda, Copepoda. *Int. Revue ges. Hydrobiol.* 51: 147-195
- BIRKY, C. W., 1964. Studies on the physiology and genetics of the Rotifer *Asplanchna*. I. Methods and physiology. *J. exp. zool.* 155: 273-288
- BIRKY, C. W., 1965. Studies on the physiology and genetics of the Rotifer *Asplanchna*. II. The genic base of a case of male sterility. *J. exper. zool.* 158: 349-356
- BIRKY, C. W., 1967. Studies on the physiology and genetics of the Rotifer *Asplanchna*. III. Results of outcrossing, selfing and selection. *J. exp. zool.* 164: 105-116
- BIRKY, C. W. & J. A. POWER, 1969. The development genetics of polymorphism in the Rotifer *Asplanchna*. II. A method for quantitative analysis of changes in morphogenesis induced by vitamin E and the short-term inheritance of the effects of vitamin E. *J. exp. zool.* 170: 157-166
- BJÖRKLUND, B. G., 1972a. The rotifer fauna of rock-pools in Tvärrminne archipelago, southern Finland. *Acta zool. fenn.* 135: 1-30
- BJÖRKLUND, B. G., 1972b. Taxonomical and ecological studies of species of *Notholca* (Rotatoria) found in sea and brackish water, with description of a new species. *Sarsia* 51: 25-65
- BLAKAR, I. A. & O. J. JACOBSEN, 1979. Zooplankton distribution and abundance in seven lakes from Jotunheimen, a Norwegian high mountain area. *Arch. Hydrobiol.* 85: 227-290
- BLOMQUIST, P. & B. PEJLER, 1980. Experiments on phototaxis in limnetic zooplankters. *Limnologica* (B) 12: 7-13
- BLOUIN, A. C., T. M. COLLINS & J. J. KEREKES, 1984. Plankton of an acid-stressed lake (Kejim-kujik National Park, Nova Scotia, Canada). Part 2. Population dynamics of an enclosure experiment. *Verh. intern. Ver. Limnol.* 22: 401-405
- BOGAERT, G., 1988. Zooplankton distributie in een ondergelopen steengroeve bij Doornik. *Lic. thesis, RU Gent.*
- BOGAERT, G. & H. J. DUMONT, 1989. Community structure and coexistence of the rotifers of an artificial crater lake. *Hydrobiologia* 186/187: 167-179
- BOGDAN, K. G., J. J. GILBERT & P. L. STARKWEATHER, 1980. *In situ* clearance rates of planktonic rotifers. *Hydrobiologia* 73: 73-77
- BOGDAN, K. G. & J. J. GILBERT, 1982. Seasonal patterns of feeding by natural populations of *Keratella*, *Polyarthra* and *Bosmina*: clearance rates, selectivities and contributions to community grazing. *Limnol. Oceanogr.* 27 (5): 918-934
- BOGDAN, K. G. & J. J. GILBERT, 1987. Quantitative comparison of food niches in some freshwater zooplankton. A multi-tracer approach. *Oecologia* 72: 331-340
- BOGOSLOWSKI, A. S., 1961. Distribution of Rotatoria in a section of the river Klyas'ma, its pools and a pond as influenced by a barrage. *Trud. User. Gidrobiol. Obslch.* 11: 54-81
- BOGOSLOWSKI, A. S., 1962. Material to the study of the Rotifer Fauna of China. *Zool. Zh.* 41: 1327-1333
- BOLTOVSKOY, D. & H. E. MAZZONI, 1988. The effect of sampling gear and environmental conditions on the abundance estimates of freshwater zooplankton. *Rev. Hydrobiol. Trop.* 21: 21-34
- BOLTOVSKOY D., F. L. PEDROZO & P. A. BATTISTONI, 1984. The effects of wind and diel vertical migrations on the distribution of freshwater zooplankton. *Studies on Neotropical Fauna and Environment* 19: 137-154
- BOLTOVSKOY, D. & R. URREJOLA, 1977. Dos nuevos especies del genero *Keratella* (Rotatoria) de Tierra del Fuego, Argentina. *Limnobios* 1 (6): 181-187

- BONETTO, A. A., 1986. The Parana River system, in Davies, B. R. & K. F. Walker (eds.): The ecology of River Systems. W Junk Publishers, Dordrecht, the Netherlands
- BOONSOM, J., 1984a. Zooplankton feeding in the fish *Trichogaster pectoralis* Regan. *Hydrobiologia* 113: 217-221
- BOONSOM, J., 1984b. The freshwater zooplankton of Thailand (Rotifera and Crustacea). *Hydrobiologia* 113: 223-229
- BORAAS, M. E. & W. N. BENNETT, 1988. Steady-state rotifer growth in a two-stage, computer-controlled turbidostat. *J. Plankton Res.* 10: 1023-1038
- BOSSELMAN, S., 1979a. Population dynamics of *Keratella cochlearis* in Lake Esrom. *Arch. Hydrobiol.* 87: 152-165
- BOSSELMAN, S., 1979b. Production of *Keratella cochlearis* in Lake Esrom. *Arch. Hydrobiol.* 87: 304-313
- BOSSELMAN, S., 1981. Population dynamics and production of *Keratella hiemalis* and *K. quadrata* in Lake Esrom, DK. *Arch. Hydrobiol.* 90: 427-447
- BOSSET, E., 1962. Le lac de Joux. *Schweiz. Z. Hydrol.* 24: 90-151
- BOTHAR, A., 1982. Die Entwicklung des Zooplanktons und -benthos unter dem Einfluss des Abwassers. *Schweiz. Z. Hydrol.* 44: 230-242
- BOTOSANEANU, L. & St. NEGREA, 1961. Une oasis aquatique à faune relique dans la plaine du Danube inférieur. *Hydrobiologia* 18: 199-218
- BOTTRELL, H.H., A. DUNCAN, Z.-M. GLIWICZ, E. GRYGIEREK, A. HERZIG, A. HILLBRICHT-ILKOWSKA, H. KURASAWA, P. LARSSON & T. WEGLENSKA, 1976. A review of some problems in zooplankton production studies. *Norw. J. Zool.* 24: 419-456
- BRANDL, Z. & C. H. FERNANDO, 1979. The impact of predation by the copepod *Mesocyclops edax* (Forbes) on zooplankton in three lakes in Ontario, Canada. *Can. J. Zool.* 57: 940-942
- BRANDL, Z. & C. H. FERNANDO, 1981. The impact of predation by cyclopoid copepodes on zooplankton. *Verh. int. Verein. Limnol.* 21: 1573-1577
- BRANDL, Z. & C. H. FERNANDO, 1984. Feeding and food consumption by *Mesocyclops edax*. *Proc. 2nd Intern. Conf. Copepoda*: 254-258
- BRANDORFF, G. O., W. KOSTE & N. N. SMIRNOV, 1982. The composition and structure of Rotifera and Crustacean communities of the Lower Rio Nhamunda, Brazil. *Studies on neotropical Fauna and Environment* 17: 69-121
- BRETSCHKO, G., 1966. Der grüne See bei Tragös, Steiermark. Ein Beitrag zur Karstlimnologie. *Int. Revue ges. Hydrobiol.* 51: 699-726
- BRETT, M. Y., 1989. The rotifer communities of acid-stressed lakes of Maine. *Hydrobiologia* 186/187: 181-189
- BREZEANU, GH. & V. MARINESCU-POPESCU, 1964. Hydrobiologisches Studium des Cerna-Beckens. *Hidrobiol. Bucur.* 5: 65-94
- BREZEANU, GH. & M. PETEN, 1973. Biologische Grundlagen der Bildung und Evolution des Zooplanktons im Stausee des Eisernen Tors im ersten Jahr. *Hidrobiol. Buc.* 14: 177-187
- BRINGMANN, G. & R. KÜHN, 1962. Biomassenriter und Saproben - Eine Hydrobiologische Vergleichungsanalyse an Nieder-Rhein, Fulda und Havel. *Int. Revue ges. Hydrobiol.* 47: 123-145
- BRINSKA, M., A. HILLBRICHT-ILKOWSKA, Z. KAJAK, T. WEGLENSKA & B. ZDANOWSKI, 1976. Influence of mineral fertilization on lake ecosystem functioning. *Limnologica* 10: 255-267
- BRINSON, M. M. & F. G. NORDLIE, 1975. Lake Izabal, Guatemala. *Verh. intern. Verein. Limnol.* 19: 1468-1479
- BROMLEY, H. L. & F. D. POR, 1975. The metazoan fauna of a sewage-carrying wadi, Nahal Soreq (Judean Hills, Israel). *Freshwat. Biol.* 5: 121-133
- BROWN, A. V., R. L. LIMBECK & M. D. SCHRAM, 1989. Trophic importance of zooplankton in streams with alluvial riffle and pool geomorphometry. *Arch. Hydrobiol.* 114: 349-367
- BRYAS, J. A., 1960. A freshwater pond in New Zealand. *Austr. J. Mar. Freshw. Res.* 11: 222-240
- BRZEK, G., C. KOWALCZYK, W. LECEWICZ, S. RADWAN, WL. WOJCIECHOWSKA & I. WOJCIEKOWSKI, 1975. Influence of abiotic environmental factors on plankton in lakes of different trophy. *Pol. Arch. Hydrobiol.* 22: 123-139
- BUCHNER, H., 1982. Untersuchungen über die Bedingungen der heterogenen Fortpflanzungsarten bei den Rädertern. II. Über die Bedingungen der Parthenogenese bei *Brachionus urceolaris*. *Arch. Hydrobiol./Suppl.* 64 (3): 399-441
- BUCHNER, H., 1987. Untersuchungen... III. Über den Verlust der miktischen Potenz bei *Brachionus urceolaris*. *Arch. Hydrobiol.* 109: 333-354

- BUCHNER, H., 1992. Untersuchungen... IV. Über die Reaktivierung der miktischen Potenz bei *Brachionus urceolaris*. *Zool. Jb. Physiol.* 96: 97-165
- BUCHNER, H., H. KIECHLE & L. TIEFENBACHER, 1969. Untersuchungen über die Bedingungen der heterogenen Fortpflanzungarten bei den Rädertieren. *Zool. Jb. Physiol.* 74: 329-426
- BUCHNER, H. & F. MULZER, 1961. Untersuchungen über die Variabilität der Rädertiere. II. Der Ablauf der Variation im Freien. *Z. Morph. Ökol. Tiere* 50: 330-374
- BUCHNER, H., Chr. MUTSCHLER & H. KIECHLE, 1967. Die Determination der Männchen- und Dauereiproduktion bei *Asplanchna sieboldi*. *Biol. Zentralbl.* 86: 599-621
- BUIKEMA, A. L. Jr. & P. H. LOEFFELMAN, 1978. Effects of pumped storage operations on rotifer populations. *Verh. intern. Ver. Limnol.* 20: 1597-1603
- BUIKEMA, A. L. Jr., J. D. MILLER & W. H. YOUNGE Jr., 1978. Effects of algae and protozoans on the dynamics of *Polyarthra vulgaris*. *Verh. intern. Ver. Limnol.* 20: 2395-2399
- BÜRGI, H. R., H. BÜHRER, J. BLOESCH & E. SZABO, 1979. Der Einfluss experimentell varierter Zooplanktondichte auf die Produktion und Sedimentation im hocheutrophen See. *Schweiz. Z. Hydrol.* 41: 38-63
- BURNS, C. W. & J. J. GILBERT, 1986a. Effects of daphnid size and density of interference between *Daphnia* and *Keratella cochlearis*. *Limnol. Oceanogr.* 31: 848-858
- BURNS, C. W. & J. J. GILBERT, 1986b. Direct observations of the mechanisms of interference between *Daphnia* and *Keratella cochlearis*. *Limnol. Oceanogr.* 31: 859-866
- BUSCERNI, P. A., 1964. Some aspects of the regional limnology of northern and central Idaho. *Verh. intern. Ver. Limnol.* 15: 709-718
- BUTLER, M. N., 1983. Substrate selection and larval settlement by *Cupelopagis vorax*. *Hydrobiologia* 104: 317-323
- CAJANDER, V.-R., 1983. Productions of planktonic Rotatoria in Ormajärvi, an eutrophicated lake in southern Finland. *Hydrobiologia* 104: 329-333
- CAMPBELL, C. E. & R. KNOECHEL, 1988. Zooplankton communities and trophic structure of lakes on the Avalon Peninsula, Newfoundland. *Verh. intern. Ver. Limnol.* 23: 297-305
- CAMPOS, H., W. STEFFEN, G. AGÜERO, O. PARRA, & L. ZUNIGA, 1988. Limnological study of Lake Llanquihue (Chile). Morphometry, physics, chemistry, plankton and primary productivity. *Arch. Hydrobiol./Suppl.* 81: 37-67
- CAMPOS, H., W. STEFFEN, G. AGÜERO, C. PARRA & L. ZUNIGA, 1990. Limnological study of lake Todos los Santos (Chile). Morphometry, physics, chemistry, plankton and primary productivity. *Arch. Hydrobiol.* 117: 453-484
- CAMPOS, H., W. STEFFEN, G. AGÜERO, O. PARRA & L. ZUNIGA, 1992. Limnology of Lake Raco (Chile). *Limnologica* 22: 337-353
- CAPUZZO, J. M., 1979. The effect of halogen toxicants on survival, feeding and egg production of the rotifer *Brachionus plicatilis*. *Estuar. coast. mar. Sci.* 8: 307-316
- CARMONA, M. J. & M. SERRA, 1991. Comparative total protein and demographic patterns of mictic and amictic female rotifers. *Verh. intern. Ver. Limnol.* 24: 2754-2759
- CARRILLO, P., L. CRUZ-PIZZARO & R. MORALES-BAQUERO, 1989. Empirical evidence for a complex diurnal movement in *Hexarthra bulgarica* from an oligotrophic high mountain lake (La Caldera, Spain). *Hydrobiologia* 186/187: 103-108
- CASPERS, H., 1964. Characteristics of hypertrophic lakes and canals in cities. *Verh. intern. Ver. Limnol.* 15: 631-638
- CASPERS, H. & Ch. W. HECKMAN, 1981. Ecology of orchard drainage ditches along the freshwater section of the Elbe estuary. *Arch. Hydrobiol./Suppl.* 43: 347-486
- CASPERS, H. & Ch. W. HECKMAN, 1982. The biota of a small standing water ecosystem in the Elbe flood plain. *Arch. Hydrobiol./Suppl.* 61: 227-316
- CASPERS, H. & H. SCHULZ, 1960. Studien zur Wertung der Saprobiensysteme. Erfahrungen an einem Stadtkanal Hamburgs. *Int. Revue. ges. Hydrobiol.* 45: 535-565
- CASPERS, H. & H. SCHULZ, 1964. Die biologischen Verhältnisse der Elbe bei Hamburg. Beitrag zu einer Diskussion der Güterwertung. *Arch. Hydrobiol.* 60: 53-88
- CATHEY, D. D., B. C. PARKER, G. M. SIMMONS Jr., W. H. YOUNGE Jr. & M. R. VAN BRUNT, 1981. The microfauna of algal mats and artificial substrates in Southern Victoria lakes of Antarctica. *Hydrobiologia* 85: 3-15
- CAUTHON, St. P. & J. V. WARD, 1981. Benthos and zooplankton of coal strip mine ponds in the mountains of northwestern Colorado, USA. *Hydrobiologia* 85: 23-31

- CHAMP, P., 1974. Etude des pullulations de Rotifères périphytiques dans la Loire à l'aide de supports artificiels. *Bull. Fr. Pisc.* 252: 119-123
- CHAMP, P., 1978. Dynamique d'une population d'un Rotifère épiphytique thermophile (*Sinantherina socialis*) en présence de pollution thermique. *Arch. Hydrobiol.* 83: 213-231
- CHAMP, P. & R. POURRIOT, 1977a. Reproductive cycle in *Sinantherina socialis*. *Arch. Hydrobiol. Beih.* 8: 184-1860
- CHAMP, P. & R. POURRIOT, 1977b. Particularités biologiques et écologiques du Rotifère *Sinantherina socialis* (L.). *Hydrobiologia* 55: 55-64
- CHAPMAN, A., 1973. *Camamoecia lucasi* (Copepoda, Calanoidea) and other zooplankters in Two Rotorna, New Zealand, lakes. *Int. Revue ges. Hydrobiol.* 58: 79-104
- CHENGALATH, R., 1976. Littoral Rotifera of Ontario: the genus *Lepadella* Bory de St. Vincent. *Can. J. Zool.* 54: 901-907
- CHENGALATH, R., 1977. A list of Rotifera recorded from Canada, with synonyms. *Syllogeus* 11: 1-30
- CHENGALATH, R., 1978. A new species of the genus *Notholca* Gosse, 1886 (Brachionidae, Rotifera) from the Great Slave Lake, N. W. T. *Can. J. Zool.* 56: 363-364
- CHENGALATH, R., 1983. Rotifer and crustacean community in lakes of Newfoundland, Canada. *Abstracts XXIInd Congress S. I. L.*: 62
- CHENGALATH, R., 1984. Synopsis speciorum. Rotifera. *Bibliographia intervertebratorum aquatorium canadensis*, Vol. 3, 102pp. *Nat. Museum Canada*
- CHENGALATH, R., 1985. The Rotifera of the Canadian Arctic ice sea, with description of a new species. *Can. J. Zool.* 63: 2212-2216
- CHENGALATH, R., W. J. BRUCE & D. A. SCRUTON, 1984. Rotifer and crustacean plankton communities of lakes in insular Newfoundland. *Verh. intern. Ver. Limnol.* 22: 419-430
- CHENGALATH, R. & C. H. FERNANDO, 1973a. The planctonic rotifers of Ontario with records of distribution notes on some morphological variations. *Can. Field Nat.* 87: 267-277
- CHENGALATH, R. & C. H. FERNANDO, 1973b. Rotifers from Sri Lanka (Ceylon). I. The genus *Lecane* with description of two new species. *Bull. Fish. Stn Sri Lanka (Ceylon)* 24: 13-27
- CHENGALATH, R., C. H. FERNANDO & M. G. GEORGE, 1971. Planktonic Rotifera from Ontario. *University of Waterloo, Biology Series*, n°2: 40 pp, 94 figs.
- CHENGALATH, R., C. H. FERNANDO & W. KOSTE, 1973. Rotifera from Sri Lanka (Ceylon). II. Further studies on the Eurotatoria, including new records. *Bull. Fish. Res. Stn Sri Lanka (Ceylon)* 24: 29-62
- CHENGALATH, R., C. H. FERNANDO & W. KOSTE, 1974. Rotifera from Sri Lanka (Ceylon). III. New species and records with a list of Rotifera recorded and their distribution in different habitats from Sri Lanka. *Bull. Fish. Res. Stn Sri Lanka (Ceylon)* 25: 83-96
- CHENGALATH, R. & W. KOSTE, 1983. Rotifera from northeastern Québec, Newfoundland and Labrador, Canada. *Hydrobiologia* 104: 49-56
- CHENGALATH, R. & W. KOSTE, 1987. Rotifera from Northwestern Canada. *Hydrobiologia* 147: 49-56
- CHENGALATH, R. & W. KOSTE, 1988. Composition of littoral rotifer communities of Cape Breton Island, Nova Scotia, Canada. *Verh. Internat. Ver. Limnol.* 23: 2019-2027
- CHENGALATH, R. & W. KOSTE, 1989. Composition and distributional patterns in arctic rotifera. *Hydrobiologia* 186/187: 191-200
- CHENGALATH, R. & G. MULAMOOTTIL, 1974. Littoral Rotifera of Ontario - genus *Lecane*, with description of two new species. *Can. J. Zool.* 52: 947-957
- CHENGALATH, R. & G. MULAMOOTTIL, 1975. Littoral Rotifers of Ontario. Genus *Trichocerca*. *Can. J. Zool.* 53: 1403-1411
- CHIMNEY, M. J., R. W. WINNER & S. K. SEILKOP, 1981. Prey utilization by *Chaoborus punctipennis* Say in a small, eutrophic reservoir. *Hydrobiologia* 85: 193-199
- CHODOROWSKA, W. & A CHODOROWSKI, 1966. Quelques observations sur les rock-pools dans les Pyrénées Centrales. *Verh. internat. Ver. Limnol.* 16: 554-561
- CHUYKOV, YU. S., 1979. Zooplankton of treatment ponds in Astrakhan Province. *Hydrobiol. Jnl.* 15 (6): 51-53
- CHUYKOV, YU. S., 1981. Zooplankton of the Lower Volga and its delta. *Hydrobiol. Jnl.* 17 (3): 103-104
- CHUYKOV, YU. S. & P. G. DENESKIN, 1985. Zooplankton of the Terek delta. *Hydrobiol. Jnl.* 21 (2): 110-111
- CHUYKOV, YU. S. & A. G. MERKULOV, 1986. Association of Intervertebrate Species in the Plankton of the Lower Volga. *Hydrobiol. Jnl.* 22 (1): 76-77

- CIANFICCONI, F., G. P. MORETTI & Q. PIRISINU, 1985. Peuplements lotiques et benthiques dans le système hydrographique de la Plaine de Rieti (Latium, Italie). *Verh. internat. Ver. Limnol.* 22: 3306-3315
- CISNEROS, R. O. & E. I. MANGAS, 1991. Zooplankton studies in a tropical lake (Lake Xolotlan, Nicaragua). *Verh. internat. Ver. Limnol.* 24: 1167-1170
- CLAUS, C. & U. DALL' ASTA, 1972. Limnologische karakterisatie van het bekkencomplex tussen de Schelde, Tweegeootsloot (riolering Overmere) en het Donkmeer (O. VI.). *Licentiaatsthesis, RU Gent.*
- CLEMENT, P., 1968. Ultrastructures d'un Rotifère, *Notommata copeus*. I. La cellule-flamme. Hypothèses physiologiques. *Z. Zellforsch. Mikrosk. Anat.* 89: 478-498
- CLEMENT, P., 1969. Premières observations sur l'ultrastructure compacte des téguments des Rotifères. *Vie et Milieu* XX, fasc. 3-A: 461-483
- CLEMENT, P., 1975. Ultrastructure de l'oeil cérébral d'un Rotifère: *Trichocerca rutilus*. *J. Microsc. Biol. Cell* 22: 69-86
- CLEMENT, P. & R. POURRIOT, 1973a. Influence de la densité de population sur la production de femelles mictiques induites par la photopériode chez *Notommata copeus* (Rotifère). *C. R. Acad. Sci. Paris*, 276 D: 3151-3154
- CLEMENT, P. & R. POURRIOT, 1973b. Mise en évidence d'un effet de masse et d'un effet de groupe dans l'apparition de phases de reproduction sexuée chez le rotifère *Notommata copeus*. *C. R. Acad. Sci. Paris*, 277 D: 2533-2536
- CLEMENT, P. & R. POURRIOT, 1975. Influence du groupement et de la densité de population sur le cycle de reproduction de *Notommata copeus* (Rotifère). I. Mise en évidence et essai d'interprétation. *Arch. Zool. exp. gén.* 116: 375-422
- CLEMENT, P. & R. POURRIOT, 1976a. Photopériode et cycle hétérogonique chez le Rotifère *Notommata copeus*. IV. Influence de l'intensité d'éclairement en lumière monochromatique. Variation du pourcentage de femelles au cours de la ponte de femelles parentales. *Arch. Zool. exp. gén.* 117: 205-224
- CLEMENT, P. & R. POURRIOT, 1976b. Influence du groupement et de la densité de population sur le cycle de reproduction des Rotifères. II. Comparaison de deux souches de *Notommata copeus* Ehrb. *Arch. Zool. exp. gén.* 117: 5-14
- CLEMENT, P. & R. POURRIOT, 1977. Cytoplasmic and chromosomal inheritance of a mictic reaction in a parthenogenetic clone of *Notommata copeus*. *Arch. Hydrobiol. Beihefte* 8: 205-206
- CLEMENT, P. & R. POURRIOT, 1979. Influence de l'âge des grands-parents sur l'apparition de mâles chez le Rotifère *Notommata copeus* Ehrb. *Intern. Jnl. Invertebr. Reproduction* 1: 89-98
- COINEAU, Y. & M. KUNST, 1964. Une nouvelle espèce de Rotifère parasite d'Oligochète: *Albertia soyeri*. *Vie et Milieu* XV, fasc. 4: 1007-1015
- COMIN, F. A. & M. ALONSO, 1988. Spanish salt lakes: their chemistry and biota. *Hydrobiologia* 158: 237-245
- COMIN, F. A., M. ALONSO, P. LOPEZ & M. COMELLES, 1983. Limnology of Gallocanta Lake, Aragon, Northwestern Spain. *Hydrobiologia* 105: 207-221
- COLLADO, C., C. H. FERNANDO & D. SEPHTON, 1984. The freshwater zooplankton of Central America and the Caribbean. *Hydrobiologia* 113: 105-115
- COMITA, G. W., 1972. The seasonal zooplankton cycles, production and transformations of energy in Severson Lake, Minnesota. *Arch. Hydrobiol.* 70: 14-66
- COOPER, S. D. & Ch. R. GOLDMAN, 1980. Opussum Shrimp (*Mysis relicta*) predation on zooplankton. *Can. J. Fish. Aquat. Sci.* 37: 909-919
- CORIJN, E., 1969. Studie van het periphyton in een ondiep mesohalien brakwater. *Lic. thesis, RUG*
- CORNEANU, G., 1972. Nouveaux éléments dans le zooplancton des lacs de barrage de la vallée de Bistrița (Roumanie) - Note II. *Hidrobiol. Bucur.* 13: 109-114
- CORRILLO, P., P. SANCHEZ-CASTILLO & L. CRUZ-PIZARRO, 1991. Coincident zooplankton and phytoplankton diel migration in a high mountain lake (La Caldera, Sierra Nevada, Spain). *Arch. Hydrobiol.* 122: 57-67
- COUSSEMENT, M., 1976. Nieuwe gegevens omtrent de rotatoriënfauina van het Donkmeer in Oost-Vlaanderen. *Natuurwet. Tijdschr.* 58: 138-146
- COUSSEMENT, M., 1982. Limnologisch en visserijbiologisch onderzoek van enkele hengelwateren in Oost-Vlaanderen. *Doctoral diss. RUG*
- COUSSEMENT, M. & A. M. DE HENAU, 1975. Een vergelijkende studie van het noordelijk en zuidelijk bekken van het Donkmeer aan de hand van het periphyton. *Lic. thesis, RUG*

- COUSSEMENT, M., A. M. DE HENAU & H. J. DUMONT, 1976. *Brachionus mirabilis* Hempel and *Asplanchna girodi* de Guerne, two rotifer species new to Europe and Belgium, respectively. *Biol. Jb. Dodonaea* 44: 118-122
- COUSSEMENT, M. & H. J. DUMONT, 1980. Some peculiar elements in the rotifer fauna of the atlantic Sahara and of the Atlas mountains. *Hydrobiologia* 73: 249-254
- CRUZ-PIZARRO, L., 1978. Comparative vertical zonation and diurnal migration among Crustacea and Rotifers in the small mountain lake La Caldera (Granada, Spain). *Verh. internat. Ver. Limnol.* 20: 1026-1032
- CRUZ-PIZARRO, L. & R. MORALES, 1987. Taxonomic and ecological notes on *Hexarthra bulgarica* from high mountain lakes and ponds in the Sierra Nevada (Spain). *Hydrobiologia* 147: 91-96
- CRYER, M. & C. R. TOWNSEND, 1988. Spatial distribution of zooplankton in a shallow eutrophic lake, with a discussion of its relation to fish predation. *J. Plankt. Res.* 10: 487-501
- CULVER, D. A., 1988. Plankton ecology in fish hatchery ponds in Narrandera, NSW, Australia. *Verh. internat. Ver. Limnol.* 23: 1085-1089
- CULVER, D. A. & G. J. BRUNSKILL, 1969. Fayetteville Breen Lake, N.Y.-V. Studies of primary production and zooplankton in a meromictic marl lake. *Limnol. & Oceanogr.* 14: 862-873
- DABORN, G. R., 1974. Biological features in an aestival pond in Western Canada. *Hydrobiologia* 44: 287-299
- DAEMS, G. & H. J. DUMONT, 1973. Some interesting Rotifera from the periphyton in Central Belgium. *Biol. Jb. Dodonaea* 41: 116-119
- DAEMS, G. & H. J. DUMONT, 1974. Rotifers from Nepal, with the description of a new species of *Scardium* and a discussion of the Nepalese representatives of the genus *Hexarthra*. *Biol. Jb. Dodonaea* 42: 61-81
- DARTNALL, H. J. G., 1977. Antarctic freshwater Rotifera. *Arch. Hydrobiol. Beih.* 8: 240-242.
- DARTNALL, H.J.G., 1983. Rotifers from the Antarctic and the Subantarctic. *Hydrobiologia* 104: 57-60
- DARTNALL, H. J. G., 1992. The reproductive strategies of two antarctic Rotifers. *J. Zool. Lond.* 227: 145-162
- DARTNALL, H. J. G. & E. HOLLOWDAY, 1985. Antarctic Rotifers. *Br. Antarct. Surv. Sci. Repts* 100: 1-46
- DAVIES, J. & J. GREEN, 1990. A preliminary survey of Lake Mangnao, Palawan, The Philippines. *Asian Wetland Bureau, Publ.* n° 59. 39pp + app.
- DAVIS, Ch. C., 1969. Seasonal distribution, constitution and abundance of zooplankton in Lake Erie. *J. Fish. Res. Board Can.* 26: 2460-2476
- DAVIS, Ch. C., 1972a. Plankton dynamics in a Newfoundland Lake. *Verh. internat. Ver. Limnol.* 18: 278-283
- DAVIS, Ch. C., 1972b. Plankton succession in a Newfoundland Lake. *Internat. Rev. ges. Hydrobiol.* 57: 367-395
- DAVIS, Ch. C., 1973. A seasonal quantitative study of the plankton of Banline Long Pond, a Newfoundland Lake. *Nat. Can.* 100: 85-105
- DAVIS, Ch. C., 1974. A study of the Hatching Process in aquatic invertebrates. XXVI. The Eclosion from the Egg of *Kellicottia longispina* (Kellicott) (Rotifera). *Hydrobiologia* 44: 29-35
- DAVIS, Ch. C., 1978. Notes on zooplankton from some Labrador Lakes. *Verh. internat. Ver. Limnol.* 20: 233-239
- DAVIS, Ch. C., 1986. A Comparison of the Zooplankton in two Newfoundland Bays with different influences from Major Current. *Intern. Rev. ges. Hydrobiol.* 71: 11-47
- DAWES, C. J., B. C. COWELL, W. È. GARDINER & S. M. SCHEDA, 1987. Limnological characteristics of two eutrophic and four mesotrophic Lakes in westcentral Florida. *Internat. Rev. Ges. Hydrobiol.* 72: 171-203
- DAWIDOWICZ, P. & J. PIJANOWSKA, 1991. Competitive interactions between cladocera and rotifers: an experimental study. *Verh. internat. Ver. Limnol.* 24: 917-920
- DE BEAUCHAMP, P. 1961. Sur *Asplanchna henrietta* (Rotifera). *Bull. Soc. zool. Fr.* 86: 701-704
- DE BEAUCHAMP, P., 1963. Mizukawari: Rotifères contre anguilles. *Bull. Fr. Piscicult.* 208: 96-97
- DE BEAUCHAMP, P., 1966. Observations à propos de Rotifères. *Bull. Soc. zool. France* 91: 55-58
- DE BEAUCHAMP, P. & R. POURRIOT, 1961. A propos de *Microcodides chlaena* (Rotifère). *Bull. Soc. zool. France* 86: 705-709
- DE BENARDI, R., G. GIUSSANI, M. MAUCA & D. RUGGIN, 1988. Long-term dynamics of plankton communities in Lago Maggiore (N Italy). *Verh. internat. Ver. Limnol.* 23: 729-733

- DEEVEY, E. S., 1964. Preliminary account of fossilization of zooplankton in Rogers Lake. *Verh. internat. Ver. Limnol.* 15: 981-990
- DE GRAAF, F., 1960. De Rotatoria van het Voorste Choorven, Witven, Van Esschenven en het Achterste Choorven. Hydrobiologie van de Oisterwijkse vennen. *Publ. n° 5 Hydrobiol. Ver. Amsterdam*: 59-73
- DE INFANTE, A., 1978. Zooplankton of Lake Valencia, Venezuela. I. Species composition and abundance. *Verh. internat. Ver. Limnol.* 20: 1186-1191
- DE INFANTE, A., 1980. Los Rotiferos del Lago de Valencia. *Acta Cient. Venezolana* 31: 30-47
- DE INFANTE, A., 1982. Annual variations in abundance of zooplankton in Lake Valencia (Venezuela). *Arch. Hydrobiol.* 93: 194-208
- DE INFANTE, A. & W. RIEHL, 1984. The effects of Cyanophyta upon zooplankton in an eutrophic tropical lake (Lake Valencia, Venezuela). *Hydrobiologia* 113: 293-298
- DE LA CRUZ, A. S. & D. NORIS MILLARES, 1974. Metodo de cultivo masivo de *Brachionus plicatilis* (Rotifera) a escala experimental. *Cienc. Investig. Mar.* n° 11. 29pp.
- DELBECQUE, E. J. P. & R. E. M. SUYKERBUYK, 1988. A comparison of the periphyton of *Nuphar lutea* and *Nymphaea alba*. Spatial and temporal changes in the occurrence of sessile microfauna. *Arch. Hydrobiol.* 112: 541-566
- DE MAESENEER, J., 1972. Limnological observations on the "Nationale Watersportbaan G. Nachez" at Ghent in 1968, 1969, 1970 and 1971. *Meded. Fakult. Landbouw* 37: 289-313
- DE MAESENEER, J., 1973. Zooplanktonpopulaties in de Gentse "Nationale Watersportbaan G. Nachez" tijdens 1972, een jaar van voortdurende waterbloei. *Natuurwet. Tijdschr.* 55: 193-201
- DE MAESENEER, J., 1976a. Waarnemingen betreffende *Filinia* spp. in de Watersportbaan te Gent. *Natuurwet. Tijdschrift* 58: 226-231
- DE MAESENEER, J., 1976b. *Brachionus* and *Keratella* in fish ponds at Malacca. *Meded. Fakult. Landbouw* 410: 1-10
- DE MAESENEER, J., 1979. Morfologische en ekologische waarnemingen betreffende enkele Rotatoriën in België. *Natuurwet. Tijdschrift* 61: 108-131
- DE MAESENEER, J., N. DE PAUW & D. WAEGEMANS, 1978. Influence of the mud layer of the "Watersportbaan" at Ghent on some aquatic life forms, especially chironomid larvae and *Filinia* spp. *Hydrobiologia* 60: 151-158
- DE MANUEL, J., 1990. Contribution to the rotifer fauna (Rotifera: Monogonta) from Minorcan in-land waters (Baleares Isles, Spain). *Limnetica* 6: 119-130
- DE MANUEL, J., 1991. Distribution of Brachionidae (Rotifera: Monogonta) in Spanish reservoirs. *Verh. internat. Ver. Limnol.* 24: 2741-2744
- DE MANUEL, J., J. L. L. PRETUS & D. JAUME, 1992. Rotifera from the Balearic archipelago. *Hydrobiologia* 239: 33-41
- DEPOORTERE, H., & N. MAGIS, 1967. Mise en évidence, localisation et dosage de la chitine dans la coque des oeufs de *Brachionus leydigii* Cohn et d'autres Rotifères. *Ann. Soc. roy. Zool. Belg.* 97: 187-195
- DE RIDDER, M., 1960a. Ecologisch-faunistische studie van de raderdertjes van de Camargue (Zuid-Fr.). *Verh. Kon. Vl. Acad. Wet., Lett. en Sch. Kunsten België* n° 65. 200pp.
- DE RIDDER, M., 1961. Etude faunistique et écologique des Rotifères de la Camargue. *Biol. Jb. Dodonaea* 29: 169-232
- DE RIDDER, M., 1962. Recherches sur les Rotifères des eaux saumâtres. VIII. Quelques Rotifères des Marismas espagnoles. *Hydrobiologia* 20: 92-109
- DE RIDDER, M., 1964. Enkele raderdertjes uit de Spaanse Pyreneëen. *Biol. Jb. Dodonaea* 32: 185-201
- DE RIDDER, M., 1965. Onderzoek van een watermonster uit Syrië. *Biol. Jb. Dodonaea* 33: 234-242
- DE RIDDER, M., 1966a. Wetenschappelijke resultaten van de studiereis van Prof. Dr. P. Van Oye op IJssel. XVII. Raderdieren. *Biol. Jb. Dodonaea* 35: 121-161
- DE RIDDER, M., 1966b. Rotifers from Nicaragua. *Hydrobiologia* 27: 238-247
- DE RIDDER, M., 1967a. Rotatoriën van IJssel, II. *Natuurwet. Tijdschr.* 49: 216-228
- DE RIDDER, M., 1967b. Un étang dans la lande. *Bull. Nat. Belg.* 48: 298-324
- DE RIDDER, M., 1967c. Quelques Rotifères de Majorque. *Bull. Nat. Belg.* 48: 409-418
- DE RIDDER, M., 1967d. Raderdieren uit Griekenland. *Biol. Jb. Dodonaea* 35: 188-194
- DE RIDDER, M., 1969a. Recherches sur les Rotifères des eaux saumâtres de la Lorraine orientale (France). *Hydrobiologia* 32: 340-380
- DE RIDDER, M., 1969b. Raderdieren uit de omgeving van Budir. *Biol. Jb. Dodonaea* 37: 140-185
- DE RIDDER, M., 1970. Raderdieren van IJssel. *Verh. Kon. Vl. Acad. Wet., Lett. en Sch. Kunsten België, Klasse Wet. Jg.* 32, n° 116. 44pp.

- DE RIDDER, M., 1971. Raderdieren uit het Verre Oosten. *Biol. Jb. Dodonaea* 39: 361-391
- DE RIDDER, M., 1972. Zoology of Iceland, Vol II; part 13: Rotatoria. 102 pp. *Zoology of Iceland*, Kopenhagen & Reykjavik
- DE RIDDER, M., 1973. Atlas provisoire des Rotifères de Belgique. Cartes 1 à 234. *Cartographie des invertébrés européens. Fac. Sci. agron. Etat*, Gembloux
- DE RIDDER, M., 1977. Rotatoria of the Caribbean Region, in *Studies of the fauna of Curaçao and other Caribbean Islands* (Ed.: P. Wagenaar Hummelinck & L. J. Van der Steen). Vol. 52: 72-134
- DE RIDDER, M., 1980. Nieuwe gegevens in verband met de verspreiding der raderdieren in België. *Natuurwet. Tijdschr.* 62: 124-128
- DE RIDDER, M., 1981. Some considerations on the geographical distribution of Rotifers. *Hydro-biologia* 85: 209-225
- DE RIDDER, M., 1986. Aanvullende gegevens betreffende het voorkomen van raderdieren in België. *Natuurwet. Tijdschr.* 68: 21-33
- DE RIDDER, M., 1987 (for 1986). Annotated checklist of non-marine Rotifers (Rotifera) from African inland waters. *Zool. Doc. n° 21* - Kon. Museum voor Midden-Afrika, Tervuren, België
- DE RIDDER, M., 1991. Additions to the "Annotated checklist of non-marine Rotifers from African inland waters". *Rev. Hydrobiol. trop.* 24 (1): 25-46
- DE RIDDER, M., 1992. Distribution of Belgian Rotifera. *Proc. 8th. internat. Colloquium Eur. Inver-tebr. Survey*: 199-212
- DE RIDDER, M., 1993. Additions II to the "Annotated checklist of non-marine Rotifers from African inland waters". *Biol. Jb. Dodonaea* 61: 99-153
- DE RIDDER, M. & H. VERHEYE, 1980. Bijdrage tot de kennis van de raderdierfauna van de brakke wateren uit de Belgische kuststreek. *Natuurwet. Tijdschr.* 62: 129-138
- DERONDE, L., T. LOMEMA, I. VAN ELSLANDE, D. VAN DE GUCHT, H. BORGOO & J. J. SYMOENS, 1984. Some limnological characteristics of a temperate forest pond. *Verh. internat. Ver. Limnol.* 22: 1663-1671
- DE SMET, W. H., 1987. Rotifera uit de Galapagoseilanden. *Natuurwet. Tijdschr.* 69: 110-131
- DE SMET, W. H., 1988. Rotifers from Bjørnøya (Svalbard) with the description of *Cephalodella evabroedi* n. sp. and *Synchaeta lakowitziana arctica* n. subsp. *Fauna Norv. Ser. A* 9: 1-18
- DE SMET, W. H., 1989. Addendum bij: Rotifera uit de Galapagoseilanden. *Natuurwet. Tijdschr.* 71: 80-81
- DE SMET, W. H., 1990. Notes on the monogonont rotifers from submerged mosses collected on Hopen (Svalbard). *Fauna Norv. Ser. A* 11: 1-8
- DE SMET, W. D., A. DAS, E. A. VAN ROMPU & J. A. DE GREEF, 1988 (for 1986). Hydrobio-logisch onderzoek van oppervlaktewaters in Antwerpen. I. De grote vijver van het fort 8 van Hoboken. *Natuurwet. Tijdschr.* 68: 153-203
- DE SMET, W. H., A. DAS; E. A. VAN ROMPU & J. A. DE GREEF, 1987. Hydrobiologisch onderzoek van oppervlaktewaters in Antwerpen. 2. Het vestingwater van het fort van Merksem. *Natuurwet. Tijdschr.* 69: 156-213
- DE SMET, W. H., A. DAS, E. A. VAN ROMPU & J. A. DE GREEF, 1989. Hydrobiologisch onderzoek van oppervlaktewaters in Antwerpen. 3. De kleine vijver van het Fort 8 van Hoboken. *Natuurwet. Tijdschr.* 71: 67-113
- DE SMET, W. H. & F. J. M. C. EVENS, 1970. The influence of light and air in the purification of polluted water. *Hydrobiologia* 35: 494-544
- DE SMET, W. H., E. A. VAN ROMPU & L. BEYENS, 1988a. Rotifera, Gastrotricha en Tardigrada uit Shetland, de Faröer en Spitsbergen. *Natuurwet. Tijdschr.* 69: 81-102
- DE SMET, W. H., E. A. VAN ROMPU & L. BEYENS, 1988b. Contribution to the rotifers and aquatic tardigrada of Edgeøya (Svalbard). *Fauna Norv. Ser. A* 9: 19-30
- DEVAI, G., 1972. Hydrobiologische Untersuchungen an den Binnengewässern in der Umgebung von Debrecen. Der Einfluss der Binnengewässer auf die Wasserqualität des Ost-Hauptkanals für die Trink- und Brauchwasserversorgung von Debrecen. *Arch. Hydrobiol.* 70: 206-209
- DEVAI, G., I. BAUSCI, I. DEVAI, R. HORVATH, A. SZABO & M. TOTH, 1972-1973. Angaben zur Kenntnis der hydrobiologischen Verhältnisse des toten Fluszmarmes des Dobrog bei Sarospatak. III. Flora und Fauna. *Acta biol. debrecina* 10-11: 117-127
- DEVAUX, J., 1980. Contribution à l'étude limnologique du lac Pavin. (France). II. Relations entre paramètres abiotiques, phyto- et zooplancton dans la zone 0-20 mètres. *Hydrobiologia* 68: 17-34
- DHANAPATHI, M. V. S. S. S., 1974a. A new brachionid rotifer *Platyas quadricornis andhraensis* subsp. nov. from India. *Curr. Sci.* 43: 358
- DHANAPATHI, M. V. S. S. S., 1974b. Rotifers from Andhra Pradesh, India. I. *Hydrobiologia* 45: 357-372

- DHANAPATHI, M. V. S. S., 1974c. On the occurrence of the rare Rotifer *Rotaria neptunia* (Ehrenberg) in India. *Curr. Sci.* 42: 770
- DHANAPATHI, M. V. S. S., 1974d. New record of the rotifer *Tripleuchlanis plicata* (Levander) from India. *Curr. Sci.* 44: 130-131
- DHANAPATHI, M. V. S. S. S., 1975. Rotifers from Andhra Pradesh, India. II. *Zool. J. Linn. Soc. London* 57: 85-94
- DHANAPATHI, M. V. S. S. S., 1976a. Rotifers from Andhra Pradesh, India. III. Family Lecanidae, including two new species. *Hydrobiologia* 48: 9-16
- DHANAPATHI, M. V. S. S. S., 1976b. A new Lecanid rotifer from India. *Hydrobiologia* 50: 191-192
- DHANAPATHI, M. V. S. S. S., 1977. Studies on the distribution of *Brachionus calyciflorus* in India. *Verh. internat. Ver. Limnol.* Beih. 8: 226-229
- D'HONDT, J. L., 1970. Inventaire de la faune marine de Roscoff. Gastrotriches, Kinorhynches, Rotifères, Tardigrades. Editions de la station biologique de Roscoff. *Cah. biol. mar.*, 1970. 28pp.
- DI PERSIA, D. F. & J. J. NEIFF, 1986. The Uruguay River system, in DAVIES, B. R. & K. F. WALKER: *The ecology of River systems*: 599-621. W Junk Publishers, Dordrecht, The Netherlands
- DOLAN, J. R. & C. L. GALLEGOS, 1991. Trophic coupling of rotifers, microflagellates and bacteria during fall months in the Rhode River Estuary. *Mar. Ecol.* 77: 147-156
- DOLAN, J. R. & C. L. GALLEGOS, 1992. Trophic role of planktonic rotifers in the Rhode River Estuary, spring summer, 1991. *Mar. Ecol.* 85: 187-199
- DOLL, R., 1984. Der Tüzer See im Kreis Altentreptow. *Limnologica* 15: 179-189
- DONNER, J., 1964. Die Rotatorien-Synusien submerser Makrophyten der Donau bei Wien und mehrerer Alpenbäche. *Arch. Hydrobiol./Suppl.* XXVII: 227-324
- DONNER, J., 1968. Zwei neue Schlam-Rotatorien aus dem Neusiedlersee, *Paradicranophorus sud-zukii* und *Paradicranophorus sordidus*. *Anz. math. nat. Kl. Oesterr. Akad. Wiss.* 10: 224-232
- DONNER, J., 1970a. Die Räderterbestände submerser Moose der Salzach und anderer Wasser-Biotope des Flusgebietes. *Arch. Hydrobiol./Suppl.* XXXVI: 110-252
- DONNER, J., 1970b. Rotatorien aus einigen Böden und Moosen Spaniens und seiner Inseln. *Rev. Ecol. Biol. Sol.* 7: 501-532
- DONNER, J., 1971. Rotatorien aus einigen Auböden der Donau, aus ostmediterranen Böden und aus Kiew. *Arch. Hydrobiol./Suppl.* XXXVI: 352-376
- DONNER, J., 1972a. Die Räderterbestände submerser Moose and weiterer Merotope im Bereich der Stauräume der Donau an der deutsch-österreichischen Landesgrazie. *Arch. Hydrobiol./Suppl.* 44: 49-114
- DONNER, J., 1972b. Rotatorien aus einigen Böden des östlichen Oesterreich. *Rev. Ecol. Biol. Sol.* 9: 371-682
- DONNER, J., 1972c. Berichte über Funde von Räderterien aus der Antarktis. *Pol. Arch. Hydrobiol.* 19: 251-252
- DONNER, J., 1972d. Räderterie der Grenzschicht Wasser-Sediment aus dem Neusiedlersee. *Sitzungsber. Oesterr. Akad. Wiss., Math. -naturw. Kl.*, Abt. I, 180: 49-63
- DONNER, J., 1975. Seltene und auffallende sessile und notommatide Rotatorien aus dem Schilfgürtel des Neusiedlersees. *Sitzungsber. Oesterr. Akad. Wiss., math.-naturwiss. Kl.*, Abt. I: 131-148
- DONNER, J. 1978. Material zur saprobiologischen Beurteilung mehrerer Gewässer des Danau-Systems bei Wallsee und in der Lobau, Oesterreich, mit besonderer Berücksichtigungen der litoralen Rotatorien. *Arch. Hydrobiol./Suppl.* 52: 117-228
- DONNER, J., 1979. The Rotifers of Neudiedlersee, in W. LÖFFLER (ed.): Neusiedlersee: the limnology of a shallow lake in central Europe: 411-421. W Junk, The Hague, the Netherlands.
- DONNER, J., 1980. Einige neue Forschungen über Bdelloide Rotatorien, besonders in Boden. *Rev. Ecol. Biol. Sol* 17: 125-143
- DOR, I., H. SCHECHTER & H.-J. BROMLEY, 1987. Limnology of a hypertrophic reservoir storing wastewater effluent for agriculture at Kibbutz Na'an, Israel. *Hydrobiologia* 150: 225-241
- DOROBANTU, C., 1964. Beitrag zur Kenntnis der Rotiferenfauna des Flachseecomplexes "Crupina Jijila" (Ueberschwemmungsgebiet der Donau). *Hidrobiol. bucur.* 5: 124-133
- DOTTNE-LINDGREN, A., 1973. Horizontal variation of plankton in lakes. *Acta Univ. Uppsala* 258: 3-13
- DOUGHERTY, E. C., 1964. Cultivation and nutrition of micrometazoa. I. The antarctic Rotifer *Philodina gregaria*. *Trans. Am. microsc. Soc.* 83: 1-6
- DRABKOVA, V. G., G. I. LETANSKAYA, Ye. S. MAKARTSEVA & E. E. SHARMAN, 1979. Changes in the ecological and production indicators of Lakes, consequent upon anthropogenic eutrophication. *Hydrobiol. Jnl.* 15 (2): 1-5

- DRESSCHER, Th. G. N., 1960. Description of a community of microorganisms in purified sewage mixed with rain water from store-sewers. *Hydrobiologia* 14: 189-204
- DRESSCHER, Th. G. N., 1976. Index of names and location of algae and some groups of micro-organisms found in the Netherlands waters. *North Holl. Publ. Cie, A'dam.* 808pp.
- DULMA, A., 1979. Hydrobiological outline of the Mongolian Lakes. *Internat. Rev. ges. Hydrobiol.* 64: 709-736
- DUMONT, H. J., 1968a. A study of a man-made freshwater reservoir in Eastern Flanders (Belgium) with special reference of the migration of the plankton. *Hydrobiologia* 32: 97-130
- DUMONT, H. J., 1968b. Een limnologische studie van het Donkmeer (O.-VI.) *Doctor's dissertation*, RUG
- DUMONT, H. J., 1972. A competition-based approach of the reverse vertical migration in zooplankton and its implications, chiefly based on a study of the interactions of the Rotifer *Asplanchna priodonta* (Gosse) with several Crustacea Entomostraca. *Internat. Rev. ges. Hydrobiol.* 57: 1-38
- DUMONT, H. J., 1981. Kratergöll, a deep hypersaline crater-lake in the steppic zone of western Anatolia (Turkey), subject to occasional limnometereological perturbations. *Hydrobiologia* 82: 281-289
- DUMONT, H. J., 1983. Biogeography of rotifers. *Hydrobiologia* 104: 19-30
- DUMONT, H. J., M. COUSSEMENT & R. S. ANDERSON, 1978. An examination of some *Hexar-thra* species (Rotatoria) from western Canada and Nepal. *Can. J. Zool.* 56: 440-445
- DUMONT, H. J., M. COUSSEMENT & A. M. DE HENAU, 1977. A tentative estimation of annual production for some periphytic rotifers with a unimodal phenology. *Arch. Hydrobiol. Beih.* 8: 67-70
- DUMONT, H. J., A. M. DE HENAU & M. COUSSEMENT, 1977. Douze années de recherches hydrobiologiques sur le lac d'Overmere, *in* Productivité biologique en Belgique: 503-549. Ed. P. DUVIGNEAUD & P. KESTEMONT: *Travaux de la section belge du programme biologique international*
- DUMONT, H. J. & M. DE RIDDER, 1987. Rotifers from Turkey. *Hydrobiologia* 147: 65-73
- DUMONT, H. J., I. VAN DE VELDE & S. DUMONT, 1975. The dry weight estimate of biomass in a selection of Cladocera and Rotifera from the plankton, periphyton and benthos of continental waters. *Oecologia (B)* 19: 75-97
- DUMONT, H. J. & A. VAN VOSSEL, 1973. Een diurnaal ritme in het voorkomen van plankton in de Beneden-Samber (Mornimont), *in* Modèle mathématique de pollution d'eaux intérieures. Rapport de synthèse. Services du premier Ministre. *Commission interministérielle de la Politique Scientifique*: 274-288
- DUNCAN, A., 1983a. The composition, density and distribution of the zooplankton in Parakrama Samudra. *Devel. Hydrobiol.* 12: 85-94
- DUNCAN, A., 1983b. The influence of temperature upon the duration of embryonic development of tropical *Brachionus* species (Rotifera). *Devel. Hydrobiol.* 12: 107-115
- DUNCAN, A., 1984. Assessment of factors influencing the composition, body size and turnover rate of zooplankton in Parakrama Samudra, an irrigation reservoir in Sri Lanka. *Hydrobiologia* 113: 201-215
- DUNCAN, A. & R. D. GULATI, 1981. Parakrama Samudra (Sri Lanka) project, a study of a tropical lake ecosystem. III. Composition, density and distribution of the zooplankton in 1979. *Verh. internat. Ver. Limnol.* 21: 1001-1008
- DUNCAN, A. & R. D. GULATI, 1983a. A diurnal study of the planktonic rotifer populations in Parakrama Samudra reservoir, Sri Lanka, *in* SCHIEMER, F. ed.: *Limnology of Parakrama Samudra - Sri Lanka*: 96-106. Junk Publishers, The Hague, the Netherlands
- DUNCAN, A. & R. D. GULATI, 1983b. Feeding studies with natural food particles on tropical species of planktonic rotifers. *Devel. Hydrobiol.* 12: 117-125
- DUNN, I. G., 1970. Recovery of a tropical pond zooplankton community after destruction of algal bloom. *Limnol. Oceanogr.* 15: 373-379
- DÜRRENFELDT, A., 1978. Untersuchungen zur Besiedlungsbiologie von Kalktuff - faunistische, ökologische und elektronenmikroskopische Befunde. *Arch. Hydrobiol./Suppl.* 54: 1-79
- DUSOGE, K., L. BOWNIK-DYLINSKA, J. EJSMONT-KARABIN, I. SPADIEWSKA & T. WEGLENSKA, 1985. Plankton and benthos of man-made lake Zegrzynskie. *Ecol. Polska* 33: 455-479
- DUSSART, B., 1966. Limnologie. L'étude des eaux continentales. Paris, *Gauthier-Villars*. 677pp.
- DUTHIE, H. C. & M. L. OSTROVSKY, 1974. Plankton, chemistry and physics of lakes in the Churchill falls region of Labrador. *J. Fish. Res. Bd. Can.* 31: 1105-1117
- DVORAKOVA, M., 1960a. Einige neue und interessante Räderterre aus der Tschechoslowakei. *Acta Soc. zool. bohemoslav.* 24: 215-221
- DVORAKOVA, M., 1960b. Rotatoria des rivières des environs de Sunbark et Senov, district Ostrava (Tsjecho-Slov.). *Pr. Rodov. Casop. Slensky* 21: 425-432

- DVORAKOVA, M., 1961. Einige interessante Rotatorienarten aus den Kolken des Elbegebietes. *Acta Soc. zool. bohemoslav.* 25: 289-296
- DVORAKOVA, M., 1963. Rotatorien des Flusses Yamuna (Indien). *Acta Soc. zool. bohemoslav.* 27: 170-177
- EDMONDSON, W. T., 1960. Reproductive rates of Rotifera in natural populations. *Mem. Inst. Ital. Idrobiol. Marchi* 12: 21-77
- EDMONDSON, W. T., 1965. Reproductive rate of zooplanktonic rotifera, as related to food and temperature in nature. *Ecol. Monogr.* 35: 61-111
- EDMONDSON, W. T., 1977. Lake Washington, in SEYB, L. & R. RANDOLPH: North American Project - A study of U.S. Water Bodies - *A report for the organization for economic co-operation and development*: 288-300
- EDMONDSON, W. T. & A. H. LITT, 1987. *Conochilus* in Lake Washington. *Hydrobiologia* 147: 157-162
- EDMONDSON, W. T. & A. H. LITT, 1989. Morphological variation in *Kellicottia longispina*. *Hydrobiologia* 186/187: 109-117
- EGGERT, M. S., 1973. Daily vertical distribution of winter zooplankton in the pelagic zone of Lake Baikal. *Hydrobiol. Jnl.* 9 (1): 7-14
- EGLOFF, D. A., 1986. Effects of *Olisthodiscus luteus* on the feeding and reproduction of the marine rotifer *Synchaeta cecilia*. *J. Plankt. Res.* 8: 263-274
- EGLOFF, D. A., 1988. Food and growth relations of the marine microzooplankter, *Synchaeta cecilia* (Rotifera). *Hydrobiologia* 157: 129-141
- EINSLE, U., 1966. Die Buchseen bei Radolfzell (Bodensee) und ihr Zooplankton. *Mitt. bad. Landesver. Naturk. u. Naturschutz*, NF 9: 27-63
- EINSLE, U., 1967. Ueber das jahreszeitliche Auftreten der Planktonrotatorien im Mindelsee. *Mitt. bad. Landesver. Naturk. u. Naturschutz*, NF 9: 513-527
- EJSMONT-KARABIN, J., 1974. Studies on the feeding of planktonic polyphage *Asplanchna priodon-ta* Gosse (Rotatoria). *Ekol. Polska* 22: 311-317
- EJSMONT-KARABIN, J., R. D. GULATI & J. ROOTH, 1989. Is food availability the main factor controlling the abundance of *Euchlanis dilatata lucksiana* Hauer in a shallow, hypertrophic lake? *Hydrobiologia* 186/187: 29-34
- ELGMARK, K., 1966. On the relation between lake and pond zooplankton. *Verh. internat. Ver. Limnol.* 16: 216-221
- ELLIOTT, J. I., 1977. Seasonal changes in the abundance and distribution of planktonic rotifers in Grasmere (English Lake District). *Freshw. Biol.* 7: 147-166
- ELMORE, J. L., C. BRUCE & D. S. VODOPICH, 1984. Biological communities of three subtropical Florida lakes of different trophic character. *Arch. Hydrobiol.* 100: 455-478
- ELORANTA, P., 1981. Winter plankton in one eutrophic, ice-free pond in C Finland. *Verh. internat. Ver. Limnol.* 21: 1187-1193
- ELORANTA, P., 1982a. Notes on the morphological variation of the rotifer species *Keratella cochlearis* (Gosse) s. l. in one eutrophic pond. *J. Plankt. Res.* 4: 299-312
- ELORANTA, P., 1982b. Zooplankton in the Vasikkalampi pond, a warm water effluent recipient in Central Finland. *J. Plankton Res.* 4: 813-837
- EMIR, N., 1989a. A note on four rotifer species new to Turkey. *Biol. Jb. Dodonaea* 57: 78-80
- EMIR, N., 1989b. Samsun bafra Gölü rotatoria surlarının mevsimsel değişimi üzerine ekolojik birçalisma. *Doga-Tu Zool.* D. C. 13 S: 220-227
- EMIR, N., 1991. Some rotifer species from Turkey. *Tr. J. Zool.* 15: 39-45
- ENACEANU, V., 1964. Das Donauplankton auf rumänischem Gebiet. *Arch. Hydrobiol. /Suppl.* 27: 442-456
- ENACEANU, V. & G. BREZEANU, 1970. Die Verteilung und der Bestand der Flora und Fauna der Donau von der Quelle bis zu Mündung. *Hidrobiol. bucur.* 11: 227-264
- EPP, R. W. & W. M. LEWIS, 1980. Metabolic uniformity over the environmental temperature rate in *Brachionus plicatilis* (Rotifera). *Hydrobiologia* 73: 145-147
- EPP, R. W. & W. M. LEWIS, 1984. Cost and speed of locomotion for rotifers. *Oecologia (B)* 61: 289-292
- ERBEN, R., 1987. Rotifer fauna in the periphyton of Karst rivers in Croatia, Yugoslavia. *Hydrobiologia* 147: 103-105
- ERBEN, R., 1991. Vertical distribution of the rotifer fauna and its seasonal changes in the Plitvice Lakes, Yugoslavia. *Verh. internat. Ver. Limnol.* 24: 1066-1068
- ERIKSEN, B. G., 1968. Marine Rotifers found in Norway, with description of two new and one little known species. *Sarsia* 33: 23-34

- ERIKSEN, B. G., 1969. Rotifers from two tarns in southern Finland, with a description of a new species, and a list of Rotifers, previously found in Finland. *Acta zool. Fenn.* 125: 1-36
- ESPARCIA, A., J. ARMENGOL, E. VICENTE & M. R. MIRACLE, 1991. Vertical distribution of *Anuraeopsis* species, as related to oxygen depletion in two stratified lakes. *Verh. internat. Ver. Limnol.* 24: 2745-2749
- ESPEEL, M., 1979. Limnologische studie van de kleiputten te Stuivekenskerke (W.-Vl.). *Lic. thesis*, RUG
- EVANS, W. A., 1984. Seasonal abundance of the psammic rotifers of a physically controlled stream. *Hydrobiologia* 108: 105-114
- EVERITT, D. A., 1981. An ecological study of an Antarctic freshwater pool with particular reference to Tardigrada and Rotifera. *Hydrobiologia* 83: 225-237
- FAAFENG, Bj. A. & J. P. NILSSEN, 1981. A twenty-year study of eutrophication in a deep soft-water lake. *Verh. internat. Ver. Limnol.* 21: 412-424
- FAIRCHILD, G. W., R. S. STEMBERGER, L. C. EPSKAMP & H. A. DEBANGH, 1977. Environmental variables affecting small-scale distributions of five Rotifer species in Lancaster Lake, Michigan. *Internat. Rev. ges. Hydrobiol.* 62: 511-521
- FERGG, I., 1964. Untersuchungen über die Variabilität der Rädertiere. IV. Vergleichend-biometrische Untersuchungen über die Variation von *Keratella cochlearis* und *Keratella quadrata*. *Zool. Anz. Suppl.* 27: 253-268
- FERGUSON, A. J. D., J. M. THOMPSON & C. S. REYNOLDS, 1982. Structure and dynamics of zooplankton communities maintained in closed systems, with special reference to the algal food supply. *J. Plankt. Res.* 4: 523-543
- FERNANDO, C. H., 1969. A guide to the freshwater fauna of Ceylon. Suppl. 3. *Bull. Fish. Res. Stn Ceylon* 20: 15-25
- FERNANDO, C. H., 1980a. The freshwater zooplankton of Sri Lanka, with a discussion of tropical freshwater zooplankton composition. *Internat. Rev. ges. Hydrobiol.* 65: 85-125
- FERNANDO, C. H., 1980b. The species and size composition of tropical freshwater zooplankton, with special reference to the Oriental Region (South-East Asia). *Internat. Rev. ges. Hydrobiol.* 65: 411-426
- FERNANDO, C. H., 1984. Reservoirs and lakes of Southwest Asia, in TAUB, F. B.: Lakes and reservoirs: 411-446. *Elsevier Sci. Publ.* A'dam, the Netherlands
- FERNANDO, C. H. & R. RAJAPAKSA, 1983. Some remarks on long-term and seasonal changes in the zooplankton of Parakrama Samudra, Sri Lanka (an ancient man-made lake). *Devel. Hydrobiol.* 12: 77-84
- FERNANDO, C. H. & N. P. ZANKAI, 1981. The Rotifera of Malaysia and Singapore, with remarks on some species. *Hydrobiologia* 78: 205-219
- FERRARA, D. & F. G. MARGARITORA, 1977. Popolamento zooplanctonico in ALFINITO, S. et al.: Indagini ecologiche sul Lago de Giulianello (Latium). *Ann. bot.* 35-36: 299-313
- FERRARI, I., 1972. Structure and dynamics of pelagic zooplankton in lakes Bolsena, Bracciano and Vico (Central Italy). *Mem. Ist. Ital. Idrobiol.* 29: 209-227
- FERRARI, I., 1976. Winter limnology of a mountain lake: Lago Santo Parmense (Northern Apennines, Italy). *Hydrobiologia* 51: 245-257
- FERRARI, I., C. BELLAVERA & L. CAMURRI, 1973. Richerche ivernali al Lago Santo Parmense (Apennino Settentrionale). Atenes Parmense - *Acta nat.* IX, fasc. 3: 1-20
- FERRARI, I., A. CALCO, M. S. MAZZOCCHI & G. MATTEUCCHI, 1984. Dynamics of planctonic rotifer populations in an experimental canal. *Att. 4° Simp. Dinam. Pop.* (Parma, 22-24 Ottobre 1981): 51-57
- FERRARI, I., M. T. CANTARELLI, M. G. MAZZOCCHI & R. TOSI, 1985. Analysis of a 24-hour cycle of zooplankton sampling in a lagoon of the Po River Delta. *J. Plankt. Res.* 7: 849-865
- FERRARI, I., A. DE MARCHI, P. MENOZZI, F. MINZONI & F. PICCOLI, 1984. Heleoplankton seasonal succession in an experimental ricefield in Northern Italy. *Verh. internat. Ver. Limnol.* 22: 1711-1716
- FERRARI, I., A. FARABEGOLI & R. MAZZONI, 1989. Abundance and diversity of planktonic rotifers in the Po River. *Hydrobiologia* 166/167: 201-208
- FERRARI, I., R. MAZZONI & A. SOLAZZI, 1986. Il popolamento zooplanctonico del fiume Po nell'estate 1985. *Att. 7° Congr. A.T.O.L.* (Trieste, 11-14 Giugno 1986): 261-266
- FESCI, S., 1969. Preliminary study concerning the ecosystems of the Doftana and Brabu lakes. *Hidro-biol. bucur.* 10: 221-229
- FILIMONOVA, Z. I., 1976. Freshwaters Rotifers (Rotatoria) of Sovjet Karelia. *Hydrobiol. Jnl.* 12 (1): 15-19

- FINDLEY, D. L., D. I. FINDLEY & J. R. STEIN, 1973. Surface nitrogen and plankton in Skaha Lake, British Columbia, Can. *Freshw. Biol.* 3: 111-122
- FOCKE, E., 1961. Die Rotatoriengattung *Notholca* und ihr Verhalten im Salzwasser. *Kieler Meeresf.* 17: 190-205
- FOLT, C. L., 1985. Predator efficiencies and prey risks at high and low prey densities. *Verh. internat. Ver. Limnol.* 22: 3210-3214
- FORAN, J. A. & R. H. KING, 1982. A regression analysis of the summer population of *Polyarthra vulgaris* in a northern Michigan bog lake. *Hydrobiologia* 94: 237-246
- FORRO, L. & L. RONKAY, 1983. Contributions to the aquatic microfauna (Rotatoria and Crustacea) of the Hortobagy National Park. *Fauna Hortobagy Nat. Park*: 57-65
- FORSYTH, D. J. & M. R. JAMES, 1991. Seasonal abundance and diel vertical dynamics of zooplankton in Lake Okaro. *Arch. Hydrobiol.* 121: 419-429
- FOTT, J., 1960. Zwei *Encentrum*-Arten (Rot.) aus der Hohen Tatra mit Beschreibung von *Encentrum martoides* n. sp. *Acta Soc. zool. bohemoslav.* 24: 175-182
- FRANCEZ, A. J., 1981. Rotifères de quelques tourbières d'Auvergne. *Ann. Stat. biol. Besse-en-Chandesse* 15: 276-287
- FRANCEZ, A. J., 1982. Etude botanique et zoologique du site de Chambedaze. Université de Clermont II. Ministère de l'environnement. *Parc des Volcans d'Auvergne*: 66-73
- FRANCEZ, A. J., 1984a. Rotifères sessiles observés en Auvergne. *Bull. Nat. parisiens*, NS 40: 73-80
- FRANCEZ, A. J., 1984b. Ecologie des peuplements des Rotifères sessiles de lacs-tourbières d'Auvergne (France). *Bull. Ecol.* 15: 231-237
- FRANCEZ, A. J., 1984c. Microfaune des tourbières. Aspects généraux et étude des peuplements de Cladocères et de Rotifères. Université Pierre et Marie Curie (Paris VI). *D.E.A. d'Ecologie* 108 pp. + XVII pp. Annexes
- FRANCEZ, A. J., 1987. Successions écologiques dans les tourbières: le peuplement en Rotifères du lac-tourbière de Chambedaze (Puy-de-Dôme, France). *Bull. Ecol.* 18: 31-38
- FRANCEZ, A. J., 1988. Le peuplement de rotifères libres de deux lacs-tourbières du Puy-de-Dôme (France). *Vie et Milieu* 38: 281-292
- FRANCEZ, A. J. & J. DEVAUX, 1985. Répartition des Rotifères dans deux lacs-tourbières du Massif Central (France). *Hydrobiologia* 128: 265-276
- FRANCEZ, A. J. & R. POURRIOT, 1984. Remarques taxinomiques sur quelques Rotifères des tourbières avec la description d'une espèce et d'une sous-espèce nouvelles. *Hydrobiologia* 109: 125-130
- FROST, Th. & P. K. MONTZ, 1988. Early zooplankton response to experimental acidification in Little Rock Lake, Wisconsin, USA. *Verh. internat. Ver. Limnol.* 23: 2279-2285
- FRY, D. L. & J. A. OSBORNE, 1980. Zooplankton abundance and diversity in Central Florida grass carp ponds. *Hydrobiologia* 68: 145-155
- FU, Y., K. HIRAYAMA & Y. NATSUKARI, 1991. Morphological differences between 2 types of the rotifer *Brachionus plicatilis* O. F. Müller. *J. experim. mar. biol. & ecol.* 151: 29-41
- FULTON III, R. S., 1988. Grazing on filamentous algae by herbivorous zooplankton. *Freshw. Biol.* 20: 263-271
- FULTON III, R. S. & H. W. PEARL, 1987. Effects of colonial morphology on zooplankton utilization of algal resources during blue-green algal (*Microcystis aeruginosa*) bloom. *Limnol. Oceanogr.* 32: 634-644
- GALAT, D. L., E. L. LIDER, SVIGG & S. R. ROBERTSON, 1981. Limnology of a large, deep, North American terminal lake, Pyramid Lake, Nevada, USA. *Hydrobiologia* 82: 281-317
- GALKOVSKAYA, G. A., J. EJSMONT-KARABIN & V. N. EVDOKIMOW, 1987. Relative protein metabolism in Rotifer *Brachionus calyciflorus* Pallas, in relation to temperature. *Int. Rev. ges. Hydrobiol.* 72: 59-70
- GALTOVSKAYA, G. A. & I. F. MITYANINA, 1989. Morphological structure and functional patterns of *Keratella cochlearis* (Gosse) populations in stratified lakes. *Hydrobiologia* 186/187: 119-128
- GALLAGHER, J. J., 1965. Winter Rotifers of Bayou de Siard, Monroe, Louisiana. *Proc. Louisiana Acad. Sci.* 28: 31-40
- GALLAGHER, J. J., 1966. Rotifers new to Louisiana. *Proc. Louisiana Acad. Sci.* 29: 59-63
- GALLIFORD, A. L., 1960a. The Rotifera and Entomostraca of the Ellesmere District. *Caradoc & Severn Valley Trans.* 14: 221-228
- GALLIFORD, A. L., 1960b. Notes on the microscopic Fauna and Flora of the Leeds-Liverpool Canal in South-West Lancashire. *Proc. Liverpool Nat. Field Club for 1959*: 23-28

- GALLIFORD, A. L., 1964. Further notes on the micro-fauna and -flora of Knowsley Park. *Proc. Liver-pool Nat. Field Club* 1964 - 69: 4 pp.
- GALLIFORD, A. L., 1967. Notes on the fauna of Dozmary Pool, Bodmin Moor, Cornwall. *J. Quek. micr. Club* 30: 277-280
- GALLUP, D. N. & M. HICKMAN, 1975. Effects of the Discharge of Thermal effluent from a power station on Lake Wabamun, Alberta, Canada. Limnological features. *Hydrobiologia* 46: 45-69
- GALLUP, D. N., P. VAN DER GIESSEN & H. BOERGER, 1971. A survey of plankton and bottom invertebrates of the Peace-Athabasca Delta region. *Can. Wildlife Service*, Edmonton. 36 pp.
- GANNON, J. E., 1981. Zooplankton of the North American Great Lakes. *Verh. internat. Ver. Limnol.* 21: 1725-1733
- GARNIER, J. & F. LESCHER-MOUTOUÉ, 1983. Successions saisonnières phytozooplanctoniques dans un écosystème lacustre peu profond (lac de Crêteil, France). *Abstr. 22nd. Congress SIL*: 111
- GARNIER, J. & F. LESCHER-MOUTOUÉ, 1984. Successions saisonnières phytozooplanktoniques dans un écosystème lacustre peu profond (lac de Crêteil, France). *Verh. internat. Ver. limnol.* 22: 996-1002
- GARREAU, F., C. ROUGIER & R. POURRIOT, 1988. Exploitation des ressources alimentaires par le prédateur planctonique *Asplanchna girodi* de Guerne, 1888 (Rot.) dans un lac de sablière. *Arch. Hydrobiol.* 112: 91-106
- GASITH, A. & A. S. PERRY, 1985. Use of limnocralls for pesticide toxicity studies: effect on zooplankton composition and dynamics. *Verh. internat. Ver. limnol.* 22: 2432-2436
- GAWLER, M., G. BALVAY, J.-Cl. DRUART & J.-P. PELLETIER, 1988. Plankton ecology of Lake Geneva: A test of the PEG-model. *Arch. Hydrobiol.* 114: 161-174
- GAY, C., 1979. Quelques aspects de l'écologie du plancton d'un lac de montagne eutrophe: le lac de Pétichet (Isère). *Annls Limnol.* 15: 275-290
- GEELEN, J. F. M., 1969. Vergelijkend planktononderzoek in twee Hatertse vennen. *Doctor's Dissertation Un. Nijmegen*, NL. 112 pp.
- GEORGE, M. G., 1961a. Observations on the rotifers from shallow ponds in Delhi. *Curr. Sci.* 30: 268-269
- GEORGE, M. G., 1961b. Diurnal migrations in two shallow ponds in Delhi, India. *Hydrobiologia* 18: 265-273
- GEORGE, M. G., 1966. Comparative plankton ecology of five fish tanks in Delhi, India. *Hydrobiologia* 27: 81-108
- GEORGE, M. G. & C. H. FERNANDO, 1969a. On the Rotifer *Keratella canadensis* Berzinš. *Can. J. Zool.* 47: 155-156
- GEORGE, M. G. & C. H. FERNANDO, 1969b. Seasonal distribution and vertical migration of planktonic rotifers in two lakes in Eastern Canada. *Verh. internat. Ver. Limnol.* 17: 817-829
- GEORGE, M. G. & C. H. FERNANDO, 1970. Diurnal migration in three species of Rotifers in Sunfish Lake, Ontario, Can. *Limnol. Oceanogr.* 15: 218-223
- GHILAROV, A. M., 1965. Vertical distribution of plankton rotifers (Rotatoria) in the Bolshoe Eremenskoje Lake (the Island Velikiy, Kandalaksha Bay, the White Sea). *Zool. Zh.* 44: 688-692
- GHILAROV, A. M., 1967. The zooplankton of arctic rock pools. *Oikos* 18: 82-95
- GHILAROV, A. M., 1972. A classification of Northern lakes, based on zooplankton. *Hydrobiol. Jnl.* 8 (2): 1-8
- GHILAROV, A. M., 1976. Feeding of *Cyclops strenuus* (Copepoda, Crustacea) in the Glukoboje Lake (Moscow District) in summer time. *Zool. Zh.* 55: 294-296
- GIBSON, C. E., R. H. FOY & A. G. FITZSIMONS, 1980. A limnological reconnaissance of the Lough Erne system, Ireland. *Internat. Rev. ges. Limnol.* 65: 49-84
- GIERE, O., 1968. Die Fluktuationen des marinen Zooplanktons im Elbe-Aestuar. *Arch. Hydro-biol./Suppl.* 31: 379-546
- GIJSELS, H., J. VAN FLETEREN, P. LIPPENS & D. MAERTENS, 1976. Tien jaar zoologische stages in en om de Ardennen. *Natuurwet. Tijdschr.* 58: 3-68
- GILBERT, J. J., 1963a. Mictic female production in the Rotifer *Brachionus calyciflorus*. *J. exp. Zool.* 153: 113-124
- GILBERT, J. J., 1963b. Contact chemoreception, mating behaviour and sexual isolation in the Rotifer genus *Brachionus*. *J. exp. biol.* 40: 625-641
- GILBERT, J. J., 1966. Rotifer ecology and embryological induction. *Science* 151: 1234-1237
- GILBERT, J. J., 1967. *Asplanchna* and postero-lateral spine-production in *Brachionus calyciflorus*. *Arch. Hydrobiol.* 64: 1-62
- GILBERT, J. J., 1968. Dietary control of sexuality in the Rotifer *Asplanchna brightwelli* Gosse. *Physiol. Zool.* 41: 14-43

- GILBERT, J. J., 1974. Effects of tocopherol on the growth and development of Rotifers. *Am. J. Clin. Nutr.* 27: 1005-1016
- GILBERT, J. J., 1975. Polymorphism and sexuality in the rotifer *Asplanchna*, with special reference to the effects of prey-type and clonal variation. *Arch. Hydrobiol.* 75: 442-483
- GILBERT, J. J., 1976. Selective cannibalism in the rotifer *Asplanchna sieboldi*: contact recognition of morphotype and clone. *Proc. Nat. Acad. Sci. US* 73: 3233-3237
- GILBERT, J. J., 1977. Defenses of males against cannibalism in the rotifer *Asplanchna*: size, shape and failure to elicit tactile feeding responses. *Ecol.* 58: 1128-1135
- GILBERT, J. J., 1980. Observations on susceptibility of some protists and rotifers to predation, by *Asplanchna girodi*. *Hydrobiologia* 73: 87-91
- GILBERT, J. J., 1983. Control of sexuality in *Asplanchna brightwelli*: threshold levels of dietary tocopherol and modifications of tocopherol responses by exogenous and endogenous factors. *Hydrobiologia* 104: 167-173
- GILBERT, J. J., 1985a. Escape response of the rotifer *Polyarthra*: a high-speed cinematographic analysis. *Oecologia (B)* 66: 322-331
- GILBERT, J. J., 1985b. Competition between rotifers and *Daphnia*. *Ecol.* 66: 1943-1950
- GILBERT, J. J., 1987. The *Polyarthra* escape response: defense against interference from *Daphnia*. *Hydrobiologia* 147: 235-238
- GILBERT, J. J., 1988. Susceptibility of ten rotifer species to interference from *Daphnia pulex*. *Ecol.* 69: 1826-1838
- GILBERT, J. J., 1989a. Competitive interactions between the rotifer *Synchaeta oblonga* and the Cladoceran *Scapholeberis kingi* Sars. *Hydrobiologia* 186/187: 75-80
- GILBERT, J. J., 1989b. The effect of *Daphnia* interference on a natural rotifer and ciliate community: short-term bottle experiment. *Limnol. Oceanogr.* 34: 606-617
- GILBERT, J. J., 1990. Differential effects of *Anabaena affinis* on Cladocerans and Rotifers. -Mechanisms and implications. *Ecol.* 71: 1727-1740
- GILBERT, J. J., & K. G. BOGDAN, 1981. Selectivity of *Polyarthra* and *Keratella* for flagellate and aflagellate cells. *Verh. internat. Ver. Limnol.* 21: 1515-1521
- GILBERT, J. J. & J. L. CONFER, 1986. Gigantism and the potential for interference competition in the rotifer genus *Asplanchna*. *Oecologia (B)* 70: 549-554
- GILBERT, J. J. & M. W. DURAND, 1990. Effect of *Anabaena flos-aquae* on the abilities of *Daphnia* and *Keratella* to feed and reproduce on unicellular algae. *Freshw. Biol.* 24: 577-596
- GILBERT, J. J. & K. L. KIRK, 1988. Escape response of the rotifer *Keratella*: description, stimulation, fluid dynamics and ecological significance. *Limnol. Oceanogr.* 33: 1440-1450
- GILBERT, J. J., & J. R. LITTON Jr., 1975. Dietary tocopherol and sexual reproduction in the rotifers *Brachionus calyciflorus* and *Asplanchna sieboldi*. *J. exp. Zool.* 194: 485-494
- GILBERT, J. J. & J. R. LITTON Jr., 1978. Sexual reproduction in the Rotifer *Asplanchna girodi*: effects of tocopherol and population density. *J. exp. Zool.* 204: 113-121
- GILBERT, J. J. & P. L. STARKWEATHER, 1977. Feeding in the rotifer *Brachionus calyciflorus*. I. Regulatory mechanisms. *Oecologia (B)* 28: 125-131
- GILBERT, J. J. & R. S. STEMBERGER, 1984. *Asplanchna*-induced polymorphism in the rotifer *Keratella slacki*. *Limnol. Oceanogr.* 29: 1309-1316
- GILBERT, J. J. & R. S. STEMBERGER, 1985a. Control of *Keratella* populations by interference competition from *Daphnia*. *Limnol. Oceanogr.* 30: 160-188
- GILBERT, J. J. & R. S. STEMBERGER, 1985b. Prey capture in the rotifer *Asplanchna girodi*. *Verh. internat. Ver. Limnol.* 22: 2997-3000
- GILBERT, J. J. & R. S. STEMBERGER, 1985c. The costs and benefits of gigantism in polymorphic species of the rotifer *Asplanchna*. *Arch. Hydrobiol. Beih.* 21: 185-192
- GILBERT, J. J. & J. K. WAAGE, 1967. *Asplanchna*, *Asplanchna*-substance and postero-lateral spine-length variation of the rotifer *Brachionus calyciflorus* in a natural environment. *Ecol.* 48: 1027-1031
- GILBERT, J. J. & E. S. WURDAK, 1978. Species-specific morphology of resting eggs in the Rotifer *Asplanchna*. *Trans. Amer. micr. Soc.* 97: 330-339
- GILLARD, A., 1967. Rotifères de l'Amazonie. *Bull. Inst. roy. Sc. nat. Belg.* 43, n° 30: 20 pp.
- GLIWICZ, Z. M., 1967. Zooplankton and temperature-oxygen conditions in two alpine lakes of the Tatra mountains. *Pol. Arch. Hydrobiol.* 14: 53-72
- GLIWICZ, Z. M., A. HILLBRICHT-ILKOWSKA & T. WEGLENSKA, 1978. Contribution of fish and invertebrate predation to the elimination of zooplankton biomass in two Polish lakes. *Verh. internat. Ver. Limnol.* 20: 1007-1011

- GLIWICZ, Z. M. & M. G. ROWAN, 1984. Survival of *Cyclops abyssorum tataricus* (Copepoda, Crustacea) in alpine lakes stocked with planktovorous fish. *Limnol. Oceanogr.* 29: 1290-1299
- GODEANU, St., 1961a. Contribution à la connaissance des Rotifères (Rotatoria Ehrenberg) de la R. P. Roumaine. *Comm. acad. R. P. R.* XI, 3: 337-344
- GODEANU, St., 1961b. Contribution à la connaissance des Rotifères (Rotatoria Ehrb., 1838) de la République Populaire Roumaine (II). *Comm. acad. R. P. R.* XI, 10: 1203-1212
- GODEANU, St. 1963a. Neue und bemerkenswerte Rädertiere aus dem Bucegi-Gebirge. (Südkarpathen) Rumäniens. *Zool. Anz.* 170: 374-380
- GODEANU, St., 1963b. Contributions à l'étude des rotifères de certains cours d'eau des monts Bucegi. *Stud. cerc. Biol. Ser. Zool. anim.* 15: 365-389
- GODEANU, St., 1969a. Contributii la studiul rotiferelor din ubele ape ale Muntibor Bucegi (II). *Stud. cerc. Biol. Ser. Zool. anim.* 21: 119-124
- GODEANU, St., 1969b. Biology and ecology of the plankton rotatoria populations from the lakes Herastrau, Floreasca and Tei. *Stud. cerc. Biol. Ser. Zool. anim.* 21: 179-288
- GODEANU, St., 1970a. Rotiferan fauna of the Laptici peat-bog. *Stud. cerc. Biol. Ser. Zool. anim.* 22: 157-165
- GODEANU, St. 1970b. Rotifers of some lakes of the Black Sea shore. *Soc. Stin. biol. R. S. Rom. Com. Zool.*: 51-59
- GODEANU, St., 1971. *Proalides subtilis* Rodewald, 1940 = *Liliferotrocha subtilis* (Rodewald); *Liliferotrocha urawensis* Sudzuki, 1959 synonymous with *Liliferotrocha subtilis* (Rodewald, 1940) (Rotatoria, Lecanidae, Liliferotrochinae). *Zool. Anz.* 186: 122-126
- GODEANU, St., 1976. Prezente Rotiferului *Trochosphaera solstitialis* Thorpe, 1853 in Romania. *Stud. cerc. Biol. Ser. Zool. anim.* 28: 83-84
- GODEANU, S. P., 1978. Specificity of the zooplankton in several lakes from northern Germany with different degree of eutrophication. *Verh. internat. Ver. Limnol.* 20: 963-968
- GODEANU, S., M. S. GODEANU & V. IONESCU, 1973. Annual plankton variation in the Obretin embanked basin (Danube delta). *Pence* 3: 211-260
- GODEANU, S. & V. ZINEVICI, 1983. Composition, dynamics and production of Rotatoria in the plankton of some lakes of the Danube Delta. *Hydrobiologia* 104: 247-257
- GOLDMAN, Ch. R., M. GERLETTI, P. JAVORNICKY, D. MELCHIORRI-SANTOLINI & E. DE AMEZAGA, 1968. Primary productivity, Bacteria, Phyto- and Zooplankton in Lake Maggiore: correlations and relationships with ecological factors. *Mem. Ist. Ital. Idrobiol.* 23: 49-127
- GONZALEZ, M. J. & T. M. FROST, 1992. Food limitations and seasonal population decline of rotifers. *Oecologia (B)* 89: 560
- GONZALEZ, M., T. M. FROST & P. MONTZ, 1990. Effects of experimental acidification on Rotifer population dynamics in Little Rock Lake, Wisconsin, USA. *Verh. internat. Ver. Limnol.* 24: 449-456
- GOPAL, B., P. K. GOEL, K. P. SHARMA & R. K. TRIVEDY, 1981. Limnological study of a fresh-water reservoir, Jamwa Ramgadh (Jaipur, Rajasthan, India). *Hydrobiologia* 83: 283-294
- GOPHEN, M., B. AZOULAY & M. N. BRUTON, 1988. Selective predation of Lake Kinneret zooplankton by fingerlings of *Clarias gariepinus*. *Verh. internat. Ver. Limnol.* 23: 1763-1765
- GOPHEN, M., S. SERRUYA & P. SPATARN, 1990. Zooplankton community changes in Lake Kinneret (Israel) in 1969-1985. *Hydrobiologia* 191: 39-46
- GREEN, J., 1972. Freshwater ecology in the Mato Grosso, Central Brazil. III. Association of Rotifera in meander lakes of the Rio Snia Missu. *J. nat. Hist.* 6: 229-241
- GREEN, J., 1974. The limnology of a New Zealand reservoir. *Internat. Rev. ges. Hydrobiol.* 59: 441-487
- GREEN, J., 1980. Asymmetry and variation in *Keratella tropica*. *Hydrobiologia* 73: 241-248
- GREEN, J., 1981a. Associations of Rotifers in Australian crater lakes. *J. Zool. Lon.* 193: 469-486
- GREEN, J., 1981b. Altitude and seasonal polymorphism of *Keratella cochlearis* (Rotifera) in lakes of the Auvergne, Central France. *Biol. Jnl. Linn. Soc.* 16: 55-61
- GREEN, J., 1986. Associations of zooplankton in six crater lakes in Arizona, Mexico and New Mexico. *J. Zool. Lond.* (A), 208: 135-159
- GREEN, J., 1992. Island biogeography, diversity and dominance of zooplankton in crater lakes of the Azores. *Biol. Jnl. Linn. Soc.* 46: 189-205
- GREEN, J., S. A. CORBET, E. WALTS & OEH BIAUW LAU, 1976. Ecological studies on Indonesian lakes. Overturn and restratification of Ranu Lamongan. *J. Zool. Lon.* 180: 314-354
- GREEN, J., S. A. CORBET, E. WALTZ & B. L. OEH, 1978. Ecological studies in Indonesian lakes. The montane lakes of Bali. *J. Zool. Lon.* 186: 15-18

- GREEN, J. & H. KRAMADIBRATA, 1988. A note on Lake Boang, an unusual acid lake in Flores, Indonesia. *Freshw. Biol.* 20: 195-198
- GREEN, J. & OEY BIAUW LAU, 1974. *Asplanchna* and the spines of *Brachionus calyciflorus* in two Javanese sewage ponds. *Freshw. Biol.* 4: 223-226
- GREGORY, R. S., G. S. BROWN & G. R. DABORN, 1983. Food habits of young anadromous Alewives, *Alosa pseudoharengus*, in Lake Ainslie, Nova Scotia. *Can. Field Nat.* 97: 423-426
- GRIMM, R., 1968. Biologie der gestauten Elbe. *Arch. Hydrobiol./Suppl.* 31: 281-378
- GRÖNBERG, B., 1975. Zooplankton in Lake Mälaren, 1965-1966 und 1970-1971. *SNV PM 655 - NLV Rapport* 82: 43 pp.
- GRUNDSTRÖM, R., 1987. Changes in population dynamics of *Keratella cochlearis* (Gosse), *Kellicottia longispina* (Kell.) and *Polyarthra vulgaris* Carlin in a fertilized enclosure. *Hydrobiologia* 147: 215-219
- GUISANDE, C. & N. MAZUELOS, 1991. Reproductive patterns of *Brachionus calyciflorus* Pallas at different food concentrations. *J. Plankt. Res.* 13: 279-286
- GUISANDE, C. & J. TOJA, 1988. The dynamics of various species of *Brachionus* (Rotatoria) in the Guadalquivir River. *Arch. Hydrobiol.* 112: 579-595
- GUISET, A., 1977. General distribution of planktonic rotifera in Spanish reservoirs. *Arch. Hydrobiol. Beih.* 8: 222-225
- GULATI, R. D., 1975. A study on the role of herbivorous zooplankton community as primary consumers of phytoplankton in Dutch lakes. *Verh. internat. Ver. Limnol.* 19: 1202-1210
- GULATI, R. D., 1984. The zooplankton and its grazing as measures of trophy in the Loosdrecht lakes. *Verh. internat. Ver. Limnol.* 22: 863-867
- GULATI, R. D., 1990. Zooplankton structure in the Loosdrecht lakes in relation to trophic status and recent restoration measures. *Hydrobiologia* 191: 173-188
- GULATI, R. D., J. EJSMONT-KARABIN, J. ROOTH & K. SIEVERTSEN, 1989. A laboratory study of phosphorus and nitrogen excretion of *Euchlanis dilatata lucksiana*. *Hydrobiologia* 186/187: 347-354
- GULATI, R. D., A. L. OOMS-WILMS, O. F. R. VAN TONGEREN, G. POSTEMA & K. SIEVERTSEN, 1992. The dynamics and role of limnetic zooplankton in Loosdrecht lakes (The Netherlands). *Hydrobiologia* 233: 69-86
- GULATI, R. D., G. POSTEMA & K. SIEVERTSEN, 1988. Composition, abundance and biomass, in Zooplankton studies in the Loosdrecht lakes. Limnological Institute - *Progress report* 1988: 49
- GULATI, R. D., J. ROOTH & J. EJSMONT-KARABIN, 1987. A laboratory study of feeding and assimilation in *Euchlanis dilatata lucksiana*. *Hydrobiologia* 147: 289-296
- GULATI, R. D., K. SIEVERTSEN & G. POSTEMA, 1985. Zooplankton structure and grazing activities, in relation to food quality and concentration in Dutch lakes. *Arch. Hydrobiol. Beih.* 21: 91-102
- GULYAS, F., 1973. Das Vorkommen von *Keratella tropica tropica* Apstein (Rotatoria) im Velencer See. Különlenyomat az Allatani Küzlemekek LX, 1-4. Szamaböl: 173-174
- HABERMAN, J., 1978. Seasonal dynamics of pelagic Rotifers of Lakes Peipsi-Pihkva and Vörtsjärv. *Eesti NSV Teaduste Akadeemia Toimetised* 27. *Koide Biologia*: 16-30
- HABERMAN, J., 1982. On the summer phyto- and zooplankton of Lake Palaeostomi. *Eesti NSV Teaduste Akadeemia Toimetised* 31. *Koide Biologia* 1982, n° 4: 283-291
- HABERMAN, J., 1983. Comparative analysis of plankton rotifer biomass in large Estonian lakes. *Hydrobiologia* 104: 293-296
- HABERMAN, J. & R. LAUGASTE, 1982. On the summer phyto- and zooplankton of Lake Palaeostomi. I. Species composition of phyto- and zooplankton. Brief characterization of the lake. *Eesti NSV Teaduste Akadeemia Toimetised* 31. *Koide Biologia* 1982, n° 3: 226-232
- HABERMAN, J. & A. MÄEMETS, 1987. On the zooplankton of some Evo forest lakes. *Lammi Notes* 14: 1-5
- HADA, Y., 1972. Studies on Plankton organisms in Akkeshi Bay, Eastern District of Hokkaido, Japan, in A. Y. TAKENOUTI et al.: *Biological Oceanography of the Northern North Pacific Ocean*: 173-188. *Idemetsu Shoten*, Tokyo, 1972
- HAGIWARA, A., C.-S. LEE, G. MIYAMOTO & A. HINO, 1989. Resting egg formation and hatching of the S-type rotifer *Brachionus plicatilis* at varying salinities. *Mar. Biol.* 103: 327
- HALBACH, U., 1973a. Quantitative Untersuchungen zur Association von planktischen Rotatorien in Teichen. *Arch. Hydrobiol.* 71: 233-254
- HALBACH, U., 1973b. Life Table Data and Population Dynamics of the rotifer *Brachionus calyciflorus* Pallas as influenced by periodically oscillating temperature, in W. WIESER (ed.): *Effects of temperature on ectothermic organisms*: 217-228. *Springer Verlag*, Wien.

- HALBACH, U., 1984. Population dynamics of rotifers and its consequences for ecotoxicology. *Hydrobiologia* 109: 79-96
- HALBACH, U. & H. J. BURKHARDT, 1972. Sind einfache Zeitverzögerungen die Ursache für periodische Populationsschwankungen? Vergleich experimenteller Untersuchungen an *Brachionus calyciflorus* Pallas (Rotatoria) mit Computersimulationen. *Oecologia* (B) 9: 215-222
- HALBACH, U. & G. HALBACH-KEUP, 1974. Quantitative Beziehungen zwischen Phytoplankton und der Populationsdynamik des Rotators *Brachionus calyciflorus* Pallas. Befunde aus Laboratoriumsexperimenten und Freilanduntersuchungen. *Arch. Hydrobiol.* 73: 273-309
- HAMILTON, S. K., S. J. SIPPEL, W. H. LEWIS Jr. & J. F. SAUNDERS III, 1990. Zooplankton abundance and evidence for its reduction by macrophyte mats in two Orinoco floodplain lakes. *J. Plankton Res.* 12: 345-364
- HAMMER, U. T., R. C. HAYNES, J. M. HESSELTINE & S. M. SWANSON, 1975. The saline lakes of Saskatchewan. *Verh. internat. Ver. Limnol.* 19: 589-598
- HANAZATO, T., 1990. A comparison between predation effects on zooplankton communities by *Neomysis* and *Chaoborus*. *Hydrobiologia* 198: 33-40
- HANAZATO, T., H. HAYASHI & T. TSCHIKAWA, 1989. Dynamics of zooplankton communities in enclosures of different types in a shallow eutrophic lake. *Jap. Jnl. Limnol.* 50: 25-37
- HANAZATO, T., T. TWAKUMA, M. YASUMO & M. SAKAMOTO, 1989. Effects of *Temephos* on zooplankton communities in enclosures in a shallow eutrophic lake. *Environm. Poll.* 59: 305-314
- HANAZATO, T. & M. YASUMO, 1988. Impact of predation of *Neomysis intermedia* on a zooplankton community in Lake Kasumigaura. *Verh. internat. Ver. Limnol.* 23: 2092-2098
- HANAZATO, T. & M. YASUMO, 1990. Influence of *Chaoborus* density on the effects of an insecticide on zooplankton communities in ponds. *Hydrobiologia* 194: 183-197
- HARDY, E. R., 1980. Composição do zooplâncton em cinco lagos da Amazônia Central. *Acta Amazoniana* 10: 577-609
- HARMSWORTH, R. V., 1975. The ecology of an artificial cooling lake (Lake Sangris, Ill., USA). *Verh. internat. Ver. Limnol.* 19: 2189-2200
- HAUER, J., 1961. Zur Kenntnis südamerikanischer Rotatorien. *Beitr. naturk. Forsch. SW-Deutschl.* XX: 67-69
- HAUER, J., 1964. *Lecane remanei*, n. sp., ein neues Rädertert aus dem Amazonegebiet. *Zool. Anz.* 172: 27-30
- HAUER, J., 1965a. Ueber einige im Stromgebiet des Amazonas neu entdeckte Räderterre. *Beitr. naturk. Forsch. SW-Deutschl.* XXIV: 41-46
- HAUER, J., 1965b. Zur Rotatorienfauna des Amazonegebietes. *Internat. Rev. ges. Hydrobiol.* 50: 341-389
- HAUER, J., 1966. *Brachionus gilliardi*, n. sp., *Beitr. naturk. Forsch. SW-Deutschl.* XXV: 73-74
- HAVENS, K. E. III, 1989. Seasonal succession in the plankton of a naturally acidic, highly humic lake in Northeastern Ohio, USA. *J. Plankt. Res.* 11: 1321-1327
- HAVENS, K. E. III, 1990. *Chaoborus* predation and zooplankton community structure in a rotifer-dominated lake. *Hydrobiol.* 198: 215-226
- HAVENS, K. E. III, 1991a. The importance of rotiferan and crustacean zooplankton as grazers of algal productivity in a freshwater estuary. *Arch. Hydrobiol.* 122: 1-22
- HAVENS, K. E. III, 1991b. Zooplankton dynamics in a freshwater estuary. *Arch. Hydrobiol.* 123: 69-97
- HAVENS, K. E. III, 1991c. Summer plankton dynamics in the limnetic and littoral zones of a humic acid lake. *Hydrobiologia* 215: 21-29
- HAVENS, K. E. III & J. DA COSTA, 1987. Freshwater plankton community succession during experimental acidification. *Arch. Hydrobiol.* 111: 37-65
- HAVENS, K. E. III & J. DA COSTA, 1988. An experimental analysis of the acid sensitivity of the planktonic rotifer *Keratella cochlearis*. *Internat. Rev. ges. Hydrobiol.* 73: 407-416
- HAYWARD, R. S. & J. M. VAN DEN AVYLE, 1986. The nature of zooplankton heterogeneity in a nonriverine impoundment. *Hydrobiologia* 131: 261-271
- HECKMAN, Ch. W., 1974. The seasonal succession of species in a Rice Paddy in Vientiane, Laos. *Internat. Rev. ges. Hydrobiol.* 59: 489-507
- HECKMAN, Ch. W., 1979. Rice field ecology in Northeastern Thailand. The effects of wet and dry seasons on a cultivated aquatic system, in J. ILLIES: Rice field ecology. *Monographiae biologicae* 34: 113-120. Ed. Junk, The Hague, NL

- HECKMAN, Ch. W., 1982. Ecophysiological and phylogenetic characterization of a wintertime biotic community in shallow freshwater habitats near Hamburg. *Internat. Rev. ges. Hydrobiol.* 67: 361-386
- HECKMAN, Ch. W., 1984. Effects of dike construction on the wetland ecosystem along the freshwater section of the Elbe estuary. *Arch. Hydrobiol./Suppl.* 61: 397-508
- HECKMAN, Ch. W., 1986. Tidal influence on the wetland community structure behind the dike along the Elbe estuary. *Arch. Hydrobiol./Suppl.* 75: 1-117
- HEERKLOSS, R., H. ARNDT, J. HELLWIG, V. VIETINGHOFF, F. GEORGI, B. WESSEL & W. SCHNESE, 1984. Consumption and assimilation by zooplankton related to primary production in the Baltic coastal inlet, Barther Bodden. *Limnologica* 15: 387-394
- HEERKLOSS, R., W. SCHNESE & B. ADAMKIEWICZ-CHOJNACKA, 1991. Seasonal variation in the Biomass of zooplankton in two shallow coastal water inlets differing in their stage of eutrophication. *Internat. Rev. ges. Hydrobiol.* 76: 397-404
- HEIMBOKEL, J. F., D. W. COATS, K. W. HENDERSON & M. A. TYLER, 1988. Reproduction rates and secundary production of three species of the rotifer genus *Synchaeta* in the estuarine Potomac River. *J. Plankt. Res.* 10 :
- HENDELBERG, M., G. MARLING & B. PEJLER, 1979. The ultrastructure of the lorica of the rotifer *Keratella serrulata* (Ehrb.). *Zoon* 7: 49-54
- HENRIKSON, L., H. G. NYMAN, H. G. OSCASSON & J. A. E. STENSON, 1985. Changes in the zooplankton community after lime treatment of an acidified lake. *Verh. internat. ver. Limnol.* 22: 3008-3013
- HERMANN, SCOTT, J., 1977. Analysis and structure of four lentic communities on the Colorado (USA) shortgrass prairie. *Internat. Rev. ges. Hydrobiol.* 61: 729-764
- HERZIG, A., 1980. Ten years quantitative data on a population of *Rhinoglena fertoensis* (Brachionidae, Monogononta). *Hydrobiologia* 73: 161-167
- HERZIG, A., 1983. Comparative studies on the relationship between temperature and duration of embryonic development in rotifers. *Hydrobiologia* 104: 237-246
- HERZIG, A. & W. KOSTE, 1989. The development of *Hexarthra* spp. in a shallow alkaline lake. *Hydrobiologia* 186/187: 129-136
- HERZIG, A. & O. MOOG, 1976. Die vertikale Verteilung des Zooplanktons im Goggausee (Der Einflus von Algen und *Chaoborus flavicans*). *Corinthia II* (166./66J.): 373-385
- HESSEN, D. O. & J. P. NILSSEN, 1985. Factors controlling rotifer abundances in a Norwegian eutrophic lake: an experimental study. *Ann. Limnol.* 21: 97-105
- HEWITT, D. P. & D. G. GEORGE, 1987. The population dynamics of *Keratella cochlearis* in a hyper-eutrophic tarn and the possible impact of predation by young roach. *Hydrobiologia* 147: 221-227
- HEYNIG, H., 1968. Das neuerrichtete Helme-Staubecken bei Kelbra (Kyffhäuser) im ersten Jahre seines Anstaus. *Limnologica* 6: 117-133
- HEYNIG, H., 1974. Das Helme-Staubecken bei Kelbra (Kyffhäuser). II. Limnologische und hygienische Verhältnisse im Zeitraum 1968-1969. *Limnologica* 9: 63-79
- HEYNIG, H., 1980. Das Helme-Staubecken bei Kelbra (Kyffhäuser). IV. Limnologische und hygienische Verhältnisse in den Jahren 1970-1975. *Limnologica* 12: 85-107
- HEYWOOD, R. B., 1977. A limnological survey of the Ablation Point Area, Alexander Island, Antarctica. *Phil. Trans. r. Soc. , Lond. B*, 279: 39-54
- HIGLER, L. W. G. & R. VAN HALEWIJN, 1991. Limnology of a tropical sewage pond. *Verh. intern. Ver. Limnol.* 24: 2163-2167
- HILLBRICHT-ILKOWSKA, A., 1964. The influence of the fish population on the biocenosis of a pond, using Rotifera fauna as an illustration. *Ekol. pol. A*, 12 (28): 453-503
- HILLBRICHT-ILKOWSKA, A., 1965. Character of the horizontal distribution of plankton rotifers in ponds with various densities of fish populations. *Bull. Acad. pol. Sc., Cl. II*, 13: 151-156
- HILLBRICHT-ILKOWSKA, A., 1967. Attempt of evaluation of the production and turnover of plankton rotifers on the example of *Keratella cochlearis* (Gosse). *Bull. Acad. pol. Sc. Cl. II*, 15: 35-40
- HILLBRICHT-ILKOWSKA, A., 1983. Morphological variation of *Keratella cochlearis* (Gosse) in Lake Biwa, Japan. *Hydrobiologia* 104: 297-305
- HILLBRICHT-ILKOWSKA, A., 1983a. Response of planktonic rotifers to the eutrophication process and to the autumnal shift of blooms in Lake Biwa, Japan. I. Changes in abundance and composition of rotifers. *Jap. J. Limnol.* 44: 93-106

- HILLBRICHT-ILKOWSKA, A., 1983b. Response of planktonic rotifers to the eutrophication process, and to the autumnal shift of blooms in Lake Biwa, Japan. II. Changes in fecundity and turnover of the dominant species. *Jap. J. Limnol.* 44: 107-115
- HILLBRICHT-ILKOWSKA, A., Z. GLIWICZ & I. SPODNIEWSKA, 1966. Zooplankton production and some trophic dependences in the pelagic zone of two Masurian lakes. *Verh. internat. Ver. Limnol.* 16: 432-440
- HILLBRICHT-ILKOWSKA, A., Z. KAJAK, J. EJSMONT-KARABIN, A. KARABIN & J. RYBAK, 1975. Ecosystem of the Mikolajskie Lake. The utilization of the consumers production by invertebrate predators in pelagic and profundal zones. *Pol. Arch. Hydrobiol.* 22: 53-64
- HILLBRICHT-ILKOWSKA, A. & T. WEGLENSKA, 1970. Some relations between productions and zooplankton structure of two lakes of a varying trophy. *Pol. Arch. Hydrobiol.* 17: 233-240
- HILLMAN, T. J. & R. J. SHIEL, 1991. Macro- and microinvertebrates in Australian billabongs. *Verh. internat. Ver. Limnol.* 24: 1581-1587
- HIRATA, H. & S. YAMASAKI, 1987. The effect of feeding on the respiration rate of the rotifer *Brachionus plicatilis*. *Hydrobiologia* 147: 283-288
- HOFFMANN, U., 1966. Die Weidatalsperrre. Ein Beitrag zur Limnologie der Talsperren. *Limnologica* 4: 561-578
- HOFFMANN, W., 1972. Zur Populationsökologie des Zooplanktons im Pluszsee. *Verh. internat. Ver. Limnol.* 18: 410-418
- HOFFMANN, W., 1974. Zur Taxonomie und Verbreitung von *Filinia*-Arten (Rotatoria) in holsteinischen Gewässern. *Faun.-ökol. Mitt.* 4: 437-444
- HOFFMANN, W., 1975. The influence of spring circulation on zooplankton dynamics in the Pluszsee. *Verh. internat. Ver. Limnol.* 19: 1241-1250
- HOFFMANN, W., 1980. On morphological variation in *Keratella cochlearis* populations from Holstein lakes (Northern Germany). *Hydrobiologia* 73: 255-258
- HOFFMANN, W., 1981. On temporal variation in the rotifer *Keratella cochlearis* (Gosse): the problem of the "Lauterborn-cycles". *Verh. internat. Ver. Limnol.* 21: 1522
- HOFFMANN, W., 1982. On the coexistence of two pelagic *Filinia* species (Rotatoria) in Lake Pluszsee. I. Dynamics of abundance and dispersion. *Arch. Hydrobiol.* 95: 125-137
- HOFFMANN, W., 1983. Interactions between *Asplanchna* and *Keratella cochlearis* in the Pluszsee (North Germany). *Hydrobiologia* 104: 363-365
- HOFFMANN, W., 1985. Dynamics of vertical zooplankton community structure in the Pluszsee: cluster analysis. *Verh. internat. Ver. Limnol.* 22: 2983-2986
- HOFFMANN, W., 1987. Population dynamics of hypolimnetic rotifers in the Pluszsee (Northern Germany). *Hydrobiologia* 147: 197-201
- HOFFMANN, W. & H.-J. KRAMBECK, 1991. Manipulation of the thermocline in a eutrophic lake: the response of the zooplankton. *Verh. internat. Ver. Limnol.* 24: 786-790
- HOLLAND, L., C. F. BRYAN & P. NEWMAN Jr., 1983. Water quality and the rotifer populations in the Atchafalaya River Basin, Louisiana. *Hydrobiologia* 98: 55-69
- HOLLOWDAY, E. D., 1979. The capture and ingestion of the plankton rotifer *Asplanchna priodonta*, Gosse, by the holotrichous ciliate *Trachelius*. *Microscopy* 33: 335-338
- HOLLOWDAY, E. D., 1988. On the occurrence in the British Isles of the sessile Rotifer *Cupelopagis vorax* (Leidy, 1857). *Microscopy* 36: 163-173
- HOLLOWDAY, E. D. & Ch. G. HUSSEY, 1989. A re-appraisal of two members of the genus *Notholca* from the Andes, with a note on the fine structure of the lorica of *N. foliacea* (Ehrb.). *Hydrobiologia* 186/187: 319-324
- HOLMBERG, O. & B. PEJLER, 1972. On the terrestrial microfauna of Surtsey during the summer 1970. *Surtsey Res. Progress Rep.* 6: 69-72
- HOLMBERG, O. & B. PEJLER, 1974. On the terrestrial microfauna of Surtsey during the summer 1972. *Surtsey Res. Progress Rep.* 7: 17-19
- HOPKINS, Th. L., 1966. The plankton of the St. Andrew-Bay system, Florida. *Publ. Inst. Mar. Sci. Texas* 11: 12-64
- HORKAN, J. P. K., D. N. GALLUP & J. R. NURSALL, 1977. Effects of thermal effluent upon the planctonic Rotifera-survival and egg-production. *Arch. Hydrobiol. Beih.* 8: 84-87
- HORSFALL, L., A. JELINEK & B. V. TIMMS, 1988. The influence of recreation, mainly power boating, on the ecology of the Thirlmere Lakes, NSW, Australia. *Verh. internat. Ver. Limnol.* 23: 580-587

- HOWELL, B. R., 1973. Marine fish culture in Britain. VIII. A marine rotifer, *Brachionus plicatilis* Müller, and the larvae of the mussel, *Mytilus edulis* L., as food for larval flatfish. *J. Cons. perm. int. Explor. Mer* 35: 1-6
- HRBACEK, J., O. ALBERTOVA, B. DESORTOVA, V. GOTTWALDOWA & J. POPOVSKY, 1986. Relation of the zooplankton biomass and share of large cladocerans to the concentration of total phosphorus, chlorophyl and transparency in Hubenov and Vrehlice Reservoirs. *Limnologica* 17: 301-308
- HUDEC, I., 1978. Die Rädertiere (Rotatoria) von Jursky Sur. *Biol. Bratislava* 33: 639-649
- HURLBERT, S. H., M. S. MULLA & H. R. WILLSON, 1972. Effects of an organophosphorous insecticide on the phytoplankton, zooplankton and insect populations of freshwater ponds. *Ecol. Monogr.* 42: 269-299
- HUSSEY, Ch. G., 1983. Some historical specimens of rotifers examined by scanning electron microscopy. *Hydrobiologia* 104: 131-134
- ILTIS, A., F. RISSCHER & S. SERVANT-VILDARY, 1984. Contribution à l'étude hydrobiologique des lacs salés du sud de l'Altiplano bolivien. *Rev. Hydrobiol. trop.* 17: 259-273
- IRVINE, K., H. BALLS & B. MOSS, 1990. The Entomostracan and Rotifer communities, associated with submerged plants in the Norfolk Broadland. Effects of plant biomass and species composition. *Internat. Rev. ges. Hydrobiol.* 75: 121-142
- JACKSON, D. F. & B. B. JANA, 1972. Removal of phosphates in wastewater by the periphyton communities of rotating discs (in Fayetteville, Onondaga County, N. Y.). *Verh. internat. Ver. Limnol.* 18: 881-890
- JANA, B. B., 1973. Seasonal periodicity of plankton in a freshwater pond in West Bengal, India. *Internat. Rev. ges. Hydrobiol.* 58: 881-890
- JANA, B. B. & R. N. DAS, 1980. Environmental factors affecting the seasonal changes of net plankton in two tropical fish ponds in India. *Schweiz. Z. Hydrol.* 42: 225-246
- JANKOVIC, M., 1975. Entwicklung des Zooplanktons im Bratislava-See, einem Staubecken vom Gebirgstypus. *Verh. Internat. Ver. Limnol.* 19: 1921-1927
- JANSSON, A. M., 1967. The food web of the *Cladophora*-belt fauna. *Helgoländer wiss. Meeresunters.* 15: 574-588
- JÄRNEFELT, H., 1961. Die Entwicklung der Sulfitablagen auf das Planktonbild. *Verh. internat. Ver. Limnol.* 14: 1057-1062
- JÄRNEFELT, H., 1964. Ueber das Kammerplankton einer Brackwasserbucht. *Verh. internat. Ver. Limnol.* 15: 402-414
- JAYANGAUDAR, I., 1980. Hydrobiological studies on the Ajwa reservoir, the source of raw water supply to the Baroda water works. *Hydrobiologia* 72: 113-123
- JENKEN, P. M., 1982. Temperature, hydrochemistry and plankton in Wicken Brickpits, 1930-1931. *Hydrobiologia* 97: 37-61
- JOHANSSON, S., 1983. Annual dynamics and production of rotifers in an eutrophic gradient in the Baltic Sea. *Hydrobiologia* 104: 335-340
- JOHANSSON, S., 1987. Factors influencing the occurrence of males in natural populations of *Synchaeta* spp. *Hydrobiologia* 147: 323-327
- JOLY, R., 1960. Contribution à la connaissance du peuplement d'un lac de tourbière: le Lac Luitel (Isère). *Ann. Ec. nat. Eaux for.* 17: 499-507
- JONASSON, P. M., C. LINDEGAARD, P. C. DALL, K. HAMBURGER & H. ADALSTEINSSON, 1990. Ecosystem studies on temperate lake Esrom and the subarctic lakes Myvatn and Thingvallavatn. *Limnologica* 20: 259-266
- JØRGERSEN, N. O. G., M. SØNDERGAARD, H. J. HANSEN, S. BOSSELMAN & B. RIEMANN, 1983. Diel variation in concentration, assimilation and respiration of dissolved free amino acids in relation to planctonic primary and secundary production in two eutrophic lakes. *Hydrobiologia* 107: 107-122
- JOSÉ DE PAGGI, S. B., 1976a. Distribucion espacial y temporal del zooplancton de un cuerpo de agua eutrofio (lago del Parque gral. Belgrano, Santa Fé). *Physis B* 35, n° 91: 171-183
- JOSÉ DE PAGGI, S. B., 1976b. Primeras observaciones sobre el zooplancton del río Santa Fe con especial referencia a las zonas effectadas por aportes aloctonos. *Rev. Asoc. Cienc. Nat. Lit.* 7: 139-150
- JOSÉ DE PAGGI, S. B., 1978a. First observations on longitudinal succession of zooplankton in the main course of the Paraná River between Santa Fe and Buenos Aires harbour. *Stud. Neotrop. Fauna & Environment* 13: 143-156

- JOSÉ DE PAGGI, S. B., 1978b. Observaciones sobre algunas rotiferas nuevos para la fauna argentina. *Neotropica* 24: 99-104
- JOSÉ DE PAGGI, S. B., 1979. Contribucion al conocimiento de la fauna argentina de rotiferos. II. Algunas especies de los generos *Lecane* Nitzsch & *Lepadella* Bory de St. Vincent. *Neotropica* 25: 37-44
- JOSÉ DE PAGGI, S. B., 1982a. *Notholca walterkosteai* sp. nov. y otros rotiferos dulceacuicolas de la peninsula Potter, isla 25 de Mayo (Shetland del Sur, Antarctica). *Rev. Asoc. Cienc. Nat. Litoral* 13: 81-95
- JOSÉ DE PAGGI, S. B., 1982b. Contribucion al conocimiento de la fauna argentina de rotiferos. III. Nuevos registros especificos en ambientes del río Parana. *Neotropica* 28: 117-124
- JOSÉ DE PAGGI, S. B., 1985. Distribucion vertical del zooplancton en el río Parana medio (km 603). *Rev. Hydrobiol. trop.* 18: 13-21
- JOSÉ DE PAGGI, S. B., 1989. Rotiferos de algunas provincias del Noroeste argentino. *Rev. Hydrobiol. trop.* 22: 223-238
- JOSÉ DE PAGGI, S. B., 1990. Ecological and biogeographical remarks on the rotifer fauna of Argentina. *Rev. Hydrobiol. trop.* 23: 297-311
- JOSÉ DE PAGGI, S. B., & W KOSTE, 1984. Checklist of the rotifers from Antarctic and Subantarctic areas (Rotatoria). *Senckenb. biol.* 65: 169-178
- JOSÉ DE PAGGI, S. B. & W. KOSTE, 1988. Rotifera from Soladilla river basin (Santa Fe Province, Argentina). *Hydrobiologia* 157: 13-20
- JYOTI, M. K. & H. SEHGAL, 1979. Ecology of rotifers of Surimar, a subtropical, freshwater lake in Jammu (J. & K.) India. *Hydrobiologia* 65: 23-32
- JYOTI, M. K. & H. SEHGAL, 1980. Rotifer fauna of Jammu (J. & K.) India. Part I. Loricates. *Limnologica* 12: 121-126
- KAJAK, Z., K. DUSOGE, A. HILLBRICHT-ILKOWSKA, E. PIECZYNISKI, A. PREJS, I. SPOD-NIEWSKA & T. WEGLENSKA, 1972. Influence of the artificially increased fish stock on the lake biocenosis. *Verh. internat. Ver. Limnol.* 18: 228-235
- KALAFATIC, V., 1978. Composition and dynamics of zooplankton in the Sava Lake. *Verh. internat. Ver. Limnol.* 20: 1664-1666
- KALBE, L., R. KEIL, F. RANDOW & M.-A. ROSTOCK, 1965. Der Kummerower See (Mecklemburg) aus limnologischer und hygienischer Sicht. *Arch. Hydrobiol.* 61: 161-183
- KANETSUNA, Y., 1973. On the relation between the plankton and the salinity of the waters of Hakuriki district, Japan. *Jap. J. Limnol.* 34 (1): 24-34
- KANKASLA, P., 1987. Structure, dynamics and production of mesoplankton community in the Bothnian Bay, related to environmental factors. *Internat. Rev. ges. Hydrobiol.* 72: 121-146
- KARABIN, A., 1978. The pressure of pelagic predation of the genus *Mesocyclops* (Copepoda, Crustacea) on small zooplankton. *Ecol. pol.* 26: 241-257
- KARTSASHEVA, N. V. & L. I. LEBEDEVA, 1981. The structure and abundance of Rybinsk reservoir zooplankton as a function of the concentration of zinc and cadmium compounds. *Hydrobiol. Jnl.* 17 (2): 60-65
- KARUNAKARAN, L. & A. JOHNSON, 1978. A contribution to the rotifer fauna of Singapore and Malaysia. *Malayan Nat. Jnl.* 32: 173-208
- KASIMOW, A. H., 1966. To the genesis of the freshwater fauna of the Caucasus. *Verh. internat. Ver. Limnol.* 16: 282-286
- KASPRZAK, P. & D. RONNEBERGER, 1982. Vergleichende Untersuchungen zur Struktur und Dynamik des Zooplanktons im Stechlinsee, Nehmitzsee und Hausssee 1978/79. *Limnologica* 14: 263-296
- KASYMOV, A. G., 1981. Regularities in the distribution of the high-altitude fauna of the Caucasian reservoirs and its analysis. *Verh. internat. Ver. Limnol.* 21: 513-517
- KEEFER, V. M. & R. W. PENNAK, 1977. Plankton and seston of a Colorado (USA) alpine lake: the winter anomaly and the inlet-outlet budget. *Internat. Rev. ges. Hydrobiol.* 62: 255-278
- KEEN, R. & T. MILLER, 1977. Daily timing of hatching of a Rotifer, *Keratella cochlearis*. *Hydrobiologia* 56: 127-128
- KEIL, R., L. KALBE & F. RANDOW, 1963. Das Gütebild der Warnow. *Schweiz. Z. Hydrol.* 35: 351-379
- KERTESZ, G., 1960. Die Rotatorien des Péteri-Sees. *Ann. Univ. Scient. Bud. Sect. biol.* 3: 243-251

- KIEFER, F., 1968. Der Bodensee. *Der Landkreis Konstanz*, Bd. I. Amtliche Kreisbeschreibung, herausgegeben von der Staatlichen Archivverwaltung Baden-Württemberg in Verbindung mit dem Landkreis Konstanz 1968.
- KING, C. E., 1967. Food, age and the dynamics of a laboratory population of rotifers. *Ecol.* 48: 111-128
- KING, C. E., 1970. Comparative survivorship and fecundity of mictic and amictic females of Rotifers. *Physiol. Zool.* 43: 206-213
- KING, C. E., 1972. Adaptation of rotifers to seasonal variation. *Ecol.* 53: 408-418
- KING, C. E., 1980. The genetic structure of zooplankton populations, in W. Ch. KERFOOT: *Evolution and ecology of plankton communities*: 315-328. The University Press of New England, Hanover, New Hampshire.
- KING, C. E., H. B. BAYNE, T. K. CANNON & A. E. KING, 1983. Cryopreservation of monogonont rotifers. *Hydrobiologia* 104: 85-88
- KING, C. E. & M. R. MIRACLE, 1980. A perspective on aging in rotifers. *Hydrobiologia* 73: 13-19
- KING, C. E. & T. W. SNELL, 1977. Sexual recombination in rotifers. *Heredity* 39: 357-360
- KING, C. E. & T. W. SNELL, 1980. Density-dependant sexual reproduction in natural populations of the Rotifer *Asplanchna girodi*. *Hydrobiologia* 73: 149-152
- KING, C. E. & Y. ZHAO, 1987. Coexistence of rotifer (*Brachionus plicatilis*) clones in Soda Lake, Nevada (USA). *Hydrobiologia* 147: 57-64
- KIRILLOVA, S. G. & V. A. SOKOLOVA, 1972. Zooplankton of the Kolyma Indigirka Lakes. *Hydrobiol. Jnl.* 8 (4): 67-69
- KIRK, K. L. & J. J. GILBERT, 1990. Suspended clay and the population dynamics of planctonic rotifers and cladocerans. *Ecol.* 71: 1741-1755
- KLEMENT, V., 1960. Zur Rotatorienfauna ephemerer Gewässer. *Jh. Ver. Naturk. Württ.* 115: 338-349
- KLEMENT, V., 1962. Zur Kenntnis der Rädertiere (Rotatoria) des Favorite-Parkes. *Jh. Ver. Naturk. Württ.* 117: 325-326
- KLIMOWICZ, H., 1964. Rotifers from "astatic waters". Part I. The littoral of Lake Kisajno. *Pol. Arch. Hydrobiol.* 12 (XXV): 279-305
- KLIMOWICZ, H., 1967. Rotifers from astatic waters. Part II. Rotifers of small water bodies from the Mikolajki region. *Pol. Arch. Hydrobiol.* 14: 91-110
- KLIMOWICZ, H., 1968. Occurrence of Rotifers (Rotatoria) in sewage ponds. *Pol. Arch. Hydrobiol.* 15: 25-235
- KLIMOWICZ, H., 1972a. Rotifers of the near bottom zone of lakes Mikolajskie and Taltowisko. *Pol. Arch. Hydrobiol.* 19: 167-178
- KLIMOWICZ, H., 1972b. The microfauna of activated sludge. II. Assemblages of microfauna in block aeration tanks. *Acta Hydrobiol.* 14: 19-36
- KLIMOWICZ, H., 1975. Annual development in a river water intake weir and in a treatment pond and its removal in treatment plants. *Acta Hydrobiol.* 17: 299-308
- KLIMOWICZ, H., 1979. Plankton from the canal of the River Odra and its reduction during the water treatment for waterworks. *Acta Hydrobiol.* 21: 177-184
- KLIMOWICZ, H., 1981. The plankton of the River Vistula in the region of Warsaw in the years 1977-1979. *Acta Hydrobiol.* 23: 47-67
- KLOSE, H., 1968. Untersuchungen über den Indikationswert des Potamoplanktons. *Internat. Rev. ges. Hydrobiol.* 57: 781-805
- KNAUER, G. W. & A. L. BUIKEMA Jr., 1984. Rotifer production in a small impoundment. *Verh. internat. Ver. Limnol.* 22: 1475-1481
- KNOTNERUS, J., 1973. Voorkomen van cladoceren en rotiferen tussen waterplanten van het Naardermeer en de Keverdijk (uitwateringssloot van het Naardermeer). *Doct. Dissertation Zool. lab. Univ. Amsterdam*
- KOCH-ALTHAUS: see ALTHAUS
- KOKSVIK, J. I. & H. REINERTSEN, 1991. Effects of fish alimentation on the plankton community of a lake, used in fish farming. *Verh. internat. Ver. Limnol.* 24: 2387-2392
- KORN, H., 1972. Zur Kenntnis des Retrocerebralkomplexes der Rotatorien. *Z. Morph. Ökol. Tiere* 71: 150-159
- KORSTAD, J., Y. OLSEN & O. VADSTEIN, 1989. Life history characteristics of *Brachionus plicatilis* fed different algae. *Hydrobiologia* 186/187: 43-50
- KORSTAD, J., O. VADSTEIN & Y. OLSEN, 1989. Feeding kinetics of *Brachionus plicatilis* and *Isochrysis galbana*. *Hydrobiologia* 186/187: 51-57

- KOSEL, V., 1973. Some further data on *Balatro calvus* Clarapède 1867 (Rotatoria) found in Slovakia. *Biol. Bratislava* 28: 691-695
- KOSICKI, S., 1960. The vertical distribution of pelagic Rotatoria during the summer stagnation in lakes Mikolajskie and Sniardwy. *Pol. Arch. Hydrobiol.* 8: 9-61
- KOSSWIG, K., 1967. Der Sackwiesensee in den Ostalpen (Hochschwabgebiet). Zur Limnologie eines dystrophen Gipsgewässers. *Internat. Rev. ges. Hydrobiol.* 52: 321-359
- KOSTE, W., 1961. *Paradicranophorus wockei*, nov. spec., ein Rädertier aus dem Psammon eines norddeutschen Niederungsbaches. *Zool. Anz.* 167: 138-141
- KOSTE, W., 1962. Ueber die Rotatorienfauna des Darnsees in Epe bei Bramsche, Kreis Bersenbrück. *Veröff. Naturwiss. Ver. Osnabrück* 30: 73-173
- KOSTE, W., 1965. Die Rotatorien des Naturdenkmals "Engelbergs Moor" in Druchhorn, Kreis Bersenbrück. *Veröff. Naturwiss. Ver. Osnabrück* 31: 49-82
- KOSTE, W., 1968a. Ueber moosbewohnende Rädertiere von Strohdächern alter Gebäude in Hahlen und Kl. Bokern. *Mitt. n° 15 Kreisheimath. Bersenbrück*: 123-128
- KOSTE, W., 1968b. Ueber die Rotatorienfauna des Naturschutzgebietes "Achmer Grasmoor" in Achmer, Kreis Bersenbrück. *Veröff. Naturwiss. Ver. Osnabrück* 32: 107-192
- KOSTE, W., 1968c. Ueber *Proales sigmoidea* (Skorikov, 1896), eine für Mitteleuropa neue Rotatorienart, und *Proales daphnicola* (Thompson, 1892). *Arch. Hydrobiol.* 65: 240-245
- KOSTE, W., 1968d. Das Rädertier-Porträt. Das moosbewohnende Rädertier *Tetrasiphon*. *Mikrokosmos* 57: 334-337
- KOSTE, W., 1969a. Das Rädertier-Porträt. Das parasitische Rädertier *Albertia naidis*. *Mikrokosmos* 58: 212-215
- KOSTE, W., 1969b. Das Rädertier-Porträt. Das sessile Rädertier *Ptygura velata*. *Mikrokosmos* 58: 1-6
- KOSTE, W., 1969c. Das Rädertier-Porträt. Das schönste aller Rädertiere: *Stephanodiscus fimbriatus*. *Mikrokosmos* 58: 370-373
- KOSTE, W., 1969d. Das Rädertier-Porträt. *Notommata copeus* und einige verwandte Arten. *Mikrokosmos* 58: 212-216
- KOSTE, W., 1970a. Ueber eine parasitische Rädertierart, *Albertia reichelti* n. sp. *Zool. Anz.* 184: 428-434
- KOSTE, W., 1970b. Das Rädertier-Porträt. Das Putzer-Rädertier *Proales daphnicola*. *Mikrokosmos* 59: 49-51
- KOSTE, W., 1970c. Das Rädertier-Porträt. Die Rädertiergattung *Lindia*. *Mikrokosmos* 59: 134-138
- KOSTE, W., 1970d. Das Rädertier-Porträt. *Collotheca trilobata*, ein seltenes sessiles Rädertier. *Mikrokosmos* 59: 195-199
- KOSTE, W., 1970e. Das Rädertier-Porträt. *Macrotrachela quadricornifera*, ein moosbewohnendes, bdelloides Rädertier. *Mikrokosmos* 59: 328-332
- KOSTE, W., 1970f. Zur Rotatorienfauna Nordwestdeutschlands. *Festschr. Naturwiss. Ver. Osnabrück* 33: 139-163
- KOSTE, W., 1970g. Ueber die sessilen Rotatorien einer Moorblänke in Nordwestdeutschland. *Arch. Hydrobiol.* 68: 96-125
- KOSTE, W., 1971. Das Rädertier-Porträt. Die Rädertiergattung *Collotheca*-Mitteleuropäische Arten mit besonders auffallenden Koronalfortsätzen. *Mikrokosmos* 60: 161-167
- KOSTE, 1972a. Das Rädertier-Porträt. Ein seltener Auszenparasit an Süßwasser-Oligochaeten: *Cephalodella parasitica*. *Mikrokosmos* 61: 10-12
- KOSTE, W., 1972b. Das Rädertier-Porträt. *Collotheca campanulata longicaudata*. *Mikrokosmos* 61: 97-99
- KOSTE, W., 1972c. Das Rädertier-Porträt. *Rhinoglena frontalis*, ein lebendgebärendes Rädertier. *Mikrokosmos* 61: 358-360
- KOSTE, W., 1972d. Ueber zwei seltene parasitische Rotatorienarten: *Drilophaga bucephalus* Vejdovsky und *Proales gigantea* (Glascott). *Osnabr. Naturw. Mitt.* 1: 149-158
- KOSTE, W., 1972e. Ueber ein sessiles Rädertier aus Amazonien, *Floscularia nootii* sp. n; *Arch. Hydrobiol.* 70: 534-540
- KOSTE, W., 1972f. Rotatorien aus Gewässern Amazoniens. *Amazoniana* 3: 258-505
- KOSTE, W., 1972g. Das Rädertier-Porträt. Ein Rädertier des Hochmoors: *Monommata arndti*. *Mikrokosmos* 61: 269-273
- KOSTE, W., 1972f. Ueber ein sessiles Rädertier aus Amazonien, *Ptygura elsteri* n. sp., mit Bemerkungen zur Taxonomie des Artcomplexes *Ptygura melicerta* (Ehrb., 1832). *Internat. Rev. ges. Hydrobiol.* 57: 875-882

- KOSTE, W., 1973a. Das Rädertert-Porträt. Ein merkwürdiges festsitzendes Rädertert: *Cupelopagis vorax*. *Mikrokosmos* 62: 101-106
- KOSTE, W., 1973b. *Horaella thomassoni*, ein neues Rädertert aus Gewässern der Guiana-Brasilianischen Region der Neotropis. *Arch. Hydrobiol.* 72: 375-383
- KOSTE, W., 1974a. Rotatorien aus einem Ufersee des Unteren Rio Tapajos, dem Lago Paroni (Amazonien). *Gewäss. Abwäss.* 53/54: 43-68
- KOSTE, W., 1974b. Zur Kenntnis der Rotatorienfauna der "schwimmenden Wiese", einer Uferlagune in der Varzea Amazoniens, Brasilien. *Amazoniana* 5: 25-59
- KOSTE, W., 1974c. Das Rädertert-Porträt. Die Rädertertengattung *Notholca*. *Mikrokosmos* 63: 48-52
- KOSTE, W., 1974d. Das Rädertert-Porträt. *Ptygura pectinifera*, die "Kammträgerin", eine Einwanderin in Warmwasseraquarien. *Mikrokosmos* 63: 182-187
- KOSTE, W., 1975a. Ueber den Rotatorienbestand einer Mikrobiozönose in einem tropischen aquatischen Saumbiotop, der *Eichhornia crassipes*-zone im Litoral des Bung-Borapet, einem Stausee in Zentralthailand. *Gewäss. Abwäss.* 57/58: 43-58
- KOSTE, W., 1975b. Das Rädertert-Porträt. *Macrochaeten*, die "Igel" unter den Rädertern. *Mikrokosmos* 64: 143-147
- KOSTE, W., 1975c. Das Rädertert-Porträt. *Seison annulatus*, ein ektoparasit des marinens Krebses *Nebalia*. *Mikrokosmos* 64: 341-347
- KOSTE, W., 1976a. Das Rädertert-Porträt. *Trochospaera aequatorialis*, das "Kugelrädertert". *Mikrokosmos* 65: 265-271
- KOSTE, W., 1976b. Ueber die Rädertertbestände der oberen und mittleren Hase in den Jahren 1966-1969. *Osnabr. Naturw. Mitt.* 4: 191-263
- KOSTE, W., 1977. Ueber drei neue Formen des Genus *Hexarthra* SCHMARDA 1854: *H. jenkinae*, f. *nakuru*, n. f., *H. brandorffii* n. sp. und *H. polyodonta soaplakeiensis* n. ssp. *Gewäss. Abwäss.* 62/63: 7-16
- KOSTE, W., 1978a. Rotatoria. Die Räderterte Mitteleuropas, begründet von Max Voigt. Monogononta. 673 pp. + 234 pl. *Gebr. Borntraeger, Berlin & Stuttgart*
- KOSTE, W., 1978b. Ueber *Testudinella ohlei* KOSTE, 1972, ein Rädertert der U.-Ordnung Flosculariacea aus der Guiana-Brasilianischen Region der Neotropis. *Arch. Hydrobiol.* 82: 359-363
- KOSTE, W., 1978c. Das Rädertert-Porträt. *Synchaeta grandis*, ein in Mitteleuropa vom Aussterben betrohtes Planktonrädertert. *Mikrokosmos* 67: 331-336
- KOSTE, W., 1979a. Das Rädertert-Porträt. *Hexarthra mira*, ein sechsärmige Planktonrädertert. *Mikrokosmos* 68: 134-139
- KOSTE, 1979b. *Lindia deridderi*, n. sp., ein Rädertert der Familie Lindiidae (Ueberordnung Monogononta) aus SE-Australien. *Arch. Hydrobiol.* 87: 504-511
- KOSTE, W., 1979c. New rotifers from the River Murray, South-Eastern Australia, with a review of the Australian species of *Brachionus* und *Keratella*. *Aust. J. mar. freshw. Res.* 30: 237-253
- KOSTE, W., 1980. Ueber zwei Plankton-Rädertertaxa, *Filinia australiensis* n. sp. und *Filinia hof-manni* n. sp., mit Bemerkungen zur Taxonomie der *longiseta-terminalis*-Gruppe, Genus *Filinia* BORY DE ST. VINCENT, 1824, Familie Filiniidae BARTOS, 1959 (Ueberordnung Monogononta). *Arch. Hydrobiol.* 90: 230-256
- KOSTE, W., 1981a. Das Rädertert-Porträt. Einige auffallende *Synchaeta*-Arten aus Küstengewässern. *Mikrokosmos* 70: 169-175
- KOSTE, W., 1981b. Zur Morphologie, Systematik und Oekologie von neuen monogononten Rädertern (Rotatorien) aus dem Ueberschwemmungsgebiet des Magela Creek in der Alligator-River-Region Australiens, NT. Teil I. *Osnabr. Naturwiss. Mitt.* 8: 97-126
- KOSTE, W., 1982a. Ueber *Dicranophorus robustus* Harring & Myers 1928, ein selenes carnivores Rädertert aus der Familie Dicranophoridae, Ueberordnung Monogononta, klasse Rotatoria. *Osnabr. Naturwiss. Mitt.* 9: 65-84
- KOSTE, W., 1982b. Das Rädertert-Porträt. *Ploesoma truncatum*, ein räuberisches Planktonrädertert. *Mikrokosmos* 71: 167-173
- KOSTE, W., 1983. Das Rädertert-Porträt. *Lecane*, eine formen- und artenreiche Rädertertengattung. *Mikrokosmos* 72: 174-180
- KOSTE, W., 1984. Das Rädertert-Porträt. *Trichotria tetractis* und verwandte Arten. *Mikrokosmos* 73: 113-118
- KOSTE, W., 1985a. Das Rädertert-Porträt. *Cephalodella gigantea*, ein selenes Rädertert der Faulschlammzone. *Mikrokosmos* 74: 168-173

- KOSTE, W., 1985b. Zur Morphologie, Anatomie, Oekologie und Taxonomie von *Paradicranophorus wockei* Koste, 1961 (Aschelminthes: Rotatoria: Dicranophoridae). *Senckenb. Biol.* 66: 153-166
- KOSTE, W., 1986a. Das Rädertert-Porträt. *Collotheaca edentata*, ein sessiles Rädertert ohne "Zähne". *Mikrokosmos* 75: 302-305
- KOSTE, W., 1986b. Ueber die Rotatorienfauna in Gewässern südöstlich von Concepcion, Paraguay, Südamerika. *Osnabr. Naturwiss. Mitt.* 12: 129-155
- KOSTE, W., 1987. Das Rädertert-Porträt. Ein Rädertert, das Krebse verschlingt: *Asplanchnopus multiceps*. *Mikrokosmos* 76: 171-175
- KOSTE, W., 1988a. Rotatorien aus Gewässern am Mittleren Sungai Makaham, einem Uberschwimmungsgebiet in E Kalimantan, Indonesian Borneo. *Osnabr. Naturwiss. Mitt.* 14: 91-136
- KOSTE, W., 1988b. Ueber die Rotatorien einiger Stillgewässer in der Umgebung der Biologischen Station Panguana im tropischen Regenwald in Peru. *Amazoniana* 10: 303-325
- KOSTE, W., 1989a. Ueber Rädertere (Rotatoria) aus dem Lago do Macaco, einem Ufersee des mittleren Rio Trombetas, Amazonien. *Osnabr. Naturwiss. Mitt.* 15: 199-214
- KOSTE, W., 1989b. Das Rädertert-Porträt. Die sessile Art *Octotrocha speciosa*. *Mikrokosmos* 78: 115-121
- KOSTE, W. & K. BÖTTGER, 1989. Rotatorien aus Gewässern Ecuadors. *Amazoniana* 10: 407-438
- KOSTE, W. & K. BÖTTGER, 1992. Rotatorien aus Gewässern Ecuadors II. *Amazoniana* 12: 263-303
- KOSTE, W. & E. R. HARDY, 1984. Taxonomic studies and new distribution records of Rotifera (Phylum Aschelminthes) from Rio Jalapu and Uatamâ, Amazonas, Brazil. *Amazoniana* 9: 17-29
- KOSTE, W., W. J. JANETZKY & E. VARESCHI, 1991. Ueber die Rotatorienfauna in Bromelien-Phytotelmen in Jamaika. *Osnabr. Naturwiss. Mitt.* 17: 143-170
- KOSTE, W. & JOSÉ DE PAGGI, S., 1982. Rotifera of the Superorder Monogononta recorded from Neotropis. *Gewäss. Abwäss.* 68/69: 71-102
- KOSTE, W. & J. POLTZ, 1983. Ueber ein selsames, Schlammbewohnendes Rädertert: *Paradicranophorus robustus* (Glascott, 1893) aus dem Dümmmer, NW-Deutschland. *Osnabr. Naturwiss. Mitt.* 10: 27-41
- KOSTE, W. & J. POLTZ, 1984. Ueber die Rädertere (Rotatoria, Phylum Aschelminthes) des Dümmers, NW Deutschland. *Osnabr. Naturwiss. Mitt.* 11: 91-125
- KOSTE, W. & J. POLTZ, 1987. Ueber die Rädertere (Rotatoria, Phylum Aschelminthes) des Alfsees, eines Hochwasser-Rückhaltbeckens der Hase, NW Deutschland, FRG. *Osnabr. Naturwiss. Mitt.* 13: 186-220
- KOSTE, W. & B. ROBERTSON, 1983. Taxonomic studies of the Rotifera (Phylum Aschelminthes) from a Central varzea lake, Lago Camaleão (Ilha de Marchantaria, Rio Solimões, Amazonas, Brazil). *Amazoniana* 8: 225-254
- KOSTE, W. & B. ROBERTSON, 1990. Taxonomic studies of the Rotifera from shallow waters on the Island of Maraca, Roraima, Brazil. *Amazoniana* 11: 185-200
- KOSTE, W., B. ROBERTSON & E. HARDY, 1984. Further taxonomic studies on the Rotifera from Lago Camaleão, a Central Amazonian Varzea Lake (Ilha de Marchantaria, Rio Solimões, Amazonas, Brazil). *Amazoniana* 8: 555-576
- KOSTE, W. & R. J. SHIEL, 1980a. Preliminary remarks on the characteristics of the rotifer fauna of Australia (Notogaea). *Hydrobiologia* 73: 221-227
- KOSTE, W. & R. J. SHIEL, 1980b. On *Brachionus dichotomus* SHEPHARD, 1991 (Rotatoria, Brachionidae) from the Australian region, with a description of a new subspecies, *Brachionus dichotomus reductus*. *Proc. r. Soc. Vict.* 91: 127-134
- KOSTE, W. & R. J. SHIEL, 1980c. New Rotifera from Australia. *Trans. r. Soc. S Aust.* 104: 133-144
- KOSTE, W. & R. J. SHIEL, 1983. Morphology, systematics and ecology of new monogonont rotifers (Rotatoria) from the Alligators region, Northern Territory. *Trans. r. Soc. S Aust.* 107: 109-121
- KOSTE, W. & R. J. SHIEL, 1986a. Rotifers from Australian inland waters. I. Bdelloidea (Rotifera: Digononta). *Aust. J. mar. freshw. Res.* 37: 765-792
- KOSTE, W. & R. J. SHIEL, 1986b. New Rotifera (Aschelminthes) from Tasmania. *Trans. r. Soc. S Aust.* 110: 93-109
- KOSTE, W. & R. J. SHIEL, 1987a. Tasmanian rotifera: affinities with the Australian fauna. *Hydrobiologia* 147: 31-43
- KOSTE, W. & R. J. SHIEL, 1987b. Rotifera from Australian inland waters. II. Epiphaniidae and Brachionidae (Rotifera: Monogononta). *Invertebr. Taxon.* 7: 949-1021

- KOSTE, W. & R. J. SHIEL, 1989a. Rotifera from Australian inland waters. III. Euchlanidae, Mytilinidae and Trichotriidae (Rotifera: Monogononta). *Trans. r. Soc. S Aust.* 113: 85-114
- KOSTE, W. & R. J. SHIEL, 1989b. Rotifera from Australian inland waters. IV. Colurellidae (Rotifera: Monogononta). *Trans. r. Soc. Aust.* 113: 119-143
- KOSTE, W. & R. J. SHIEL, 1990a. Rotifera from Australian inland waters. V. Lecanidae (Rotifera: Monogononta). *Trans. r. Soc. S Aust.* 114: 1-36
- KOSTE, W. & R. J. SHIEL, 1990b. Rotifera from Australian inland waters. VI. Proalidae (Rotifera: Monogononta). *Trans. r. Soc. S Aust.* 114: 129-143
- KOSTE, W. & R. J. SHIEL, 1991. Rotifera from Australian inland waters. VII. Notommatidae (Rotifera: Monogononta). *Trans. r. Soc. S Aust.* 115: 111-159
- KOSTE, W., R. J. SHIEL & M. A. BROCK, 1983. Rotifera from Western Australian wetlands with description of two new species. *Hydrobiologia* 104: 9-17
- KOSTE, W., R. J. SHIEL & L. W. TAN, 1988. New rotifers (Rotifera) from Tasmania. *Trans. r. Soc. S Aust.* 112: 119-131
- KOSTE, W. & W. TOBIAS, 1990. Zur Kenntnis der Räderterfauna des Kinda-Stausees in Zentral-Birma (Aschelminthes: Rotatoria). *Osnabr. Naturwiss. Mitt.* 16: 83-110
- KOSTE, W. & K. WULFERT, 1964. Rotatorien aus der wüste Gobi. *Limnologica* 2: 483-490
- KOSTRICHKINA, Ye. M., R. L. LINE, V. BERZIN'SH, B. A. MODRYA & M. B. HAZMACH, 1992. Species-level prediction of zooplankton abundance in the Gulf of Riga. *Hydrobiol. Jnl.* 28 (6): 27-34
- KOVAL'CHUK, N. Ye., 1990. Microzoobenthos and near-bottom Zooplankton of the Dniester Reservoir during its development. *Hydrobiol. Jnl.* 23 (5): 24-30
- KOWALCZYK, C. & S. RADWAN, 1962. The results of investigations carried out on zooplankton (Rotatoria, Cladocera, Copepoda) in Duszatinskie lakes. *Ekol. pol., Sér. B:* 307-312
- KOWALCZYK, C. & S. RADWAN, 1982. Groups of pelagic zooplankton in three lakes of different trophy. *Acta Hydrobiol.* 24: 39-51
- KOWNACKI, A., J. WOJTUSIAK & R. ZUREK, 1976. New and rare species of Rotatoria, Cladocera and Chironomidae (Diptera) for the aquatic fauna of Afghanistan. *Acta Hydrobiol.* 18: 291-304
- KOZHOV, M. M., 1961. Ueber biologische Produktivität der offenen Regionen des Baikalsees. *Verh. internat. Ver. Limnol.* 14: 176-181
- KOZHOV, M. M., 1963. Lake Baikal and its life. Ed. Junk, The Hague. 344 pp.
- KOZLOVA, I. V., 1972. Plankton in the "Ozero Ayatskoye"-reservoir as a source of food for Whitefish. *Hydrobiol. Jnl.* 8 (5): 31-39
- KOZLOVA, I. V., 1976. Productivity of zooplankton in Lake Duvankul' (Southern Urals). *Hydrobiol. Jnl.* 12 (5)): 17-22
- KRAMBECK, H. J., 1988. Partial destratification of eutrophic lakes: a tool for "ecosystem modelling". *Verh. internat. Ver. Limnol.* 23: 801-806
- KRATZ, T. K., T. M. FROST & J. J. MAGNUSEN, 1987. Interferences from spatial and temporal variability in ecosystems: long-term zooplankton data from lakes. *Am. Nat.* 129: 830-846
- KRAU, L., 1962. *Brachionus variegatus*, nova especie de Rotatoria (Brachionidae). *Mem. Istit. Osw. Cruz*, Rio de Janeiro, 60: 389-391
- KRAZHAN, S. A., N. N. KHARITANOVA, K. I. BENKO, S. A. ISAYEVA, N. M. MIKULINA & Z. G. KOKRINA, 1976. Species composition of the zooplankton in Ukrainian fish-fattening ponds with different fish stocking densities. *Hydrobiol. Jnl.* 12 (5): 22-29
- KRUGLOVA, A. N., 1981. The importance of Lacustrine Zooplankton to the feeding capacity of the Rizhma Lake-River system (Basin of Lake Onega). *Hydrobiol. Jnl.* 17 (1): 15-20
- KRUGLOVA, A. N., 1983. Zooplankton of small rivers of the Kola Peninsula. *Hydrobiol. Jnl.* 19 (5): 22-25
- KRUGLOVA, A. N., 1991. Zooplankton of the Lake-and-River system of the Umba River (White Sea Basin). *Hydrobiol. Jnl.* 27 (6): 55-62
- KRYLOV, P. I., 1988. Predation of the freshwater cyclopoid copepod *Megacyclops gigas* on Lake zooplankton ; functional response and prey selection. *Arch. Hydrobiol.* 113: 231-250
- KRYLOVA, A. G., 1971. Gas exchange in Rotifera. *Hydrobiol. Jnl.* 7 (1): 92-96
- KRYNCHKOVA, N. M., 1990. Planktonic Rotifers of biological treatment ponds. *Hydrobiol. Jnl.* 25 (6): 41-48
- KRZANOWSKI, W., 1979. Zooplankton of heated water of the Rybrik dam reservoir in 1975-1976. *Acta Hydrobiol.* 21: 261-278
- KUBICEK, F., 1964. Zur Biologie des Stausees Kninicki. *Techn. Water* 8: 371-446

- KUBICEK, F., 1968a. Die organische Drift des Svratkaflusses im Abschnitt zwischen Zwei Talsperren. *Hydrobiologia* 31: 402-416
- KUBICEK, F., 1968b. Das Zooplankton der neu eingelassenen Talsperren (Vir, Orova). *Fol. Fac. Sci. nat. Univ. Purk. Brun. Biol.* 19: 1-99
- KUCKLEUTZ, V., 1991. Influence of NTA, EDTA, phosphate and copper on the ecosystem of ponds. Part 2: Zooplankton. *Verh. internat. Ver. limnol.* 24: 2145-2148
- KUCZYNSKI, D., 1987a. Rotiferos del genero *Brachionus* nuevos para la Antartida. *Inst. Ant. Argent. Contr.* 360: 11 pp.
- KUCZYNSKI, D., 1987b. The rotifer fauna of Argentine Patagonia as a potential limnological indicator. *Hydrobiologia* 150: 3-10
- KUCZYNSKI, D., 1991a. Rotifers from Reconquista River, Argentina: the genus *Brachionus*, with description of new species. *Hydrobiologia* 215: 135-152
- KUCZYNSKI, D., 1991b. Rotiferos del Rio Reconquista (Provincia Buenos Aires, Argentina): familia Brachionidae. *Anal. Soc. Cientif. Argent.* 221: 65-80
- KUNDIJEV, V. A. & T. V. KRYZHANOVSKIY, 1986. Zooplankton as factor in the environment of freshwater fish larvae. *Hydrobiol. Jnl.* 22 (1): 80-81
- KÜSTERS, E. R., G. FRIEDRICH & H. V. HERBST, 1975. Produktionsbiologische Aspekte der Entwicklung eines mit nährstoffreichem Wasser gespeisten künstlichen Sees. *Verh. internat. Ver. Limnol.* 19: 1880-1888
- KUTIKOVA, L. A., 1962a. Species of the genus *Polyarthra* Ehrb. (Rotatoria) in the River Luga (Leningrad region). *Tr. Zool. Inst. AN USSR* 31: 453-462
- KUTIKOVA, L. A., 1962b. List of Rotatoria of the Luga district of the Leningrad region. *Acad. Sci. USSR, Sect. Zool.* 31: 463-492
- KUTIKOVA, L. A., 1964a. A new rotifer from the Antarctic. *Soviet Ant. Exp.* 1: 88-89
- KUTIKOVA, L. A., 1964b. Rotifers from the coast of east Antarctica. *Soviet Ant. Exp.* 1: 162
- KUTIKOVA, L. A., 1965. Fauna of the Karelian Lakes. Invertebrates. *Acad. Sci. USSR, Sect. Zool.* 34: 52-70
- KUTIKOVA, L. A., 1970. Kolvratki fauni SSSR, in Fauna in the USSR, Vol 104. *Academy of Sciences of the USSR, Leningrad.* 744 pp. 1174 figs.
- KUTIKOVA, L. A., 1978. On the genesis of the rotatoria fauna of Baikal Lake. *Verh. internat. Ver. Limnol.* 20: 1108-1110
- KUTIKOVA, L. A. & J. HABERMAN, 1986. Rotifers (Rotatoria) of Lake Vörtsjärv. *Proc. Acad. Sci. Est. SSR. Biol.* 35: 113-121
- KUTIKOVA, L. A., A. A. KOSOVA & O. A. KHODOREOSKY, 1976. *Brachionus charini*, sp. n. (Rotatoria), a commensal of Phyllopods. *Zool. Jnl.* 55: 763-767
- KUZNETSOVA, V. I., 1982. Zooplankton of the Cooling Pond of the Cheropet' Power Plant. *Hydrobiol. Jnl.* 18 (5): 39-45
- LACROIX, G., Ph. BOET, J. GARNIER, F. LESCHER-MOUTOUÉ, R. POURRIOT & P. TESTAROL, 1989. Factors controlling the planktonic community in the shallow Lake de Créteil, France. *Internat. Rev. ges. Hydrobiol.* 74: 353-370
- LAIR, N., 1975. Sur la production des Copépodes dans deux lacs du Massif Central Français. *Verh. internat. Ver. Limnol.* 19: 3204-3211
- LAIR, N., 1977. Biomasse et production dans deux lacs du Massif Central Français. *Arch. Hydrobiol.* 79: 247-273
- LAIR, N., 1978. Répartition spatio-temporelle, biomasse et production des populations zooplanctoniques du lac d'Aydat en période estivale. *Hydrobiologia* 61: 237-256
- LAIR, N., 1980. The rotifer fauna of the River Loire (France) at the level of the nuclear power plant. *Hydrobiologia* 73: 153-160
- LAIR, N., 1990. Effects of invertebrate predation on the seasonal succession of a zooplankton community: a two year study of Lake Aydat, France. *Hydrobiologia* 198: 1-12
- LAIR, N., 1991. Comparative grazing activities of the rotifer and crustacean communities in a eutrophic lake. *Verh. internat. Ver. Limnol.* 24: 924-927
- LAIR, N., 1992. Daytime grazing and assimilation rates of planctic copepods *Acanthodiaptomus denticornis* and *Cyclops vicinus vicinus*. Comparison of spatial and resource utilization by rotifers and cladoceran communities in a eutrophic lake. *Hydrobiologia* 231: 107-117
- LAIR, N. & H. O. ALI, 1990. Grazing and assimilation rates of natural populations of planktonic rotifers *Keratella cochlearis*, *Keratella quadrata* and *Kellicottia longispina* in a eutrophic lake (Aydat, France). *Hydrobiologia* 194: 119-131

- LAIR, N. & H. AYADI, 1989. The seasonal succession of planktonic events in Lake Aydat, France. A comparison with the PEG model. *Arch. Hydrobiol.* 115: 589-602
- LAIR, N. & W. KOSTE, 1984. The rotifer fauna and population dynamics of Lake Studer 2 (Kerguelen Archipelago), with description of *Filinia terminalis kerguelensis* n. ssp. and a new record of *Kera-tella sancta* Russell 1944. *Hydrobiologia* 108: 57-64
- LAIR, N., G. LIMMÉRIOUX & F. RESTITUTO, 1978. Examen physiochimique et répartition du plancton de la Loire, en période d'étiage (été 1976) au niveau de la centrale nucléaire de Saint-Laurent-des-Eaux. *Cah. Lab. Hydrobiol. Montereau* 6: 53-88
- LAIR, N. & F. RESTITUTO, 1976. Projet Alpin O.C.D.E. pour la lutte contre l'eutrophisation. Lacs du Massif Central Français. II. Le Lac de Tazenat, interrelations entre paramètres. *Ann. Stat. biol. Besse-en-Chandesse* 10: 100-144
- LAIR, N. & D. SARGOS, 1981. Dynamique des populations planctoniques en Loire moyenne. *Ann. Stat. Biol. Besse-en-Chandesse* 15: 60-114
- LALANCETTE, L.-M., 1987. Productivité comparée de deux étang situés à proximité du lac Saint-Jean, au Québec. *Internat. Rev. ges. Hydrobiol.* 72: 205-223
- LAMAYE, J. C., 1975. Premier aperçu de l'hydrobiologie de la vallée de la Woluwe. *Bull. Soc. r. Bot. Belg.* 108: 296-314
- LAMPERT, W. & K. G. ROTHHAUPT, 1991. Alternating dynamics of rotifers and *Daphnia magna* in a shallow lake. *Arch. Hydrobiol.* 120: 447-456
- LANDAU, R., M. GOPHEN & P. WALLINS, 1988. Larval *Mirogrex terraesanctae* (Cyprinidae) of Lake Kinneret (Israel): growthrate, plankton selectivity, consumption rates and interaction with rotifers. *Hydrobiologia* 169: 91-106
- LANGELAND, A., 1974. Long-term changes in the plankton of Lake Tyrifjord, Norway. *Norw. J. Zool.* 22: 207-219
- LANGELAND, A., 1988. Decreased zooplankton density in a mountain lake resulting from predation by recently introduced *Mysis relicta*.
- LANNER, M. & B. PEJLER, 1973. The effect of cooling water discharges on zooplankton in a bay of Lake Mälaren. *Rep. Inst. Freshw. Res. Drott. 53:* 31-33
- LARKIN, P. A., 1964. Canadian Lakes. *Verh. internat. Ver. limnol.* 15: 76-90
- LARSON, D. W. & M. W. GLASS, 1987. Spirit Lake, Mount St. Helens, Washington: limnological investigation. *U.S. Army Corps of Engineers, Portland District, USA*
- LARSON, G. L., 1973. A limnology study of a high mountain lake in Mt. Rainier National Park, Washington State, USA. *Arch. Hydrobiol.* 72: 10-48
- LARSSON, P., 1971. Vertical distribution of Planktonic Rotifers in a meromictic lake, Blankvatn near Oslo. *Norw. J. Zool.* 19: 47-75
- LARSSON, P., 1972. Distribution and estimation of standing crop of zooplankton in a mountain lake with fast renewal. *Verh. internat. Ver. Limnol.* 18: 334-342
- LAXHUBER, R., 1987. Abundance and distribution of pelagic rotifers in a cold, deep oligotrophic alpine lake (Konigssee). *Hydrobiologia* 147: 189-196
- LAXHUBER, R. & U. HARTMANN, 1988. The influence of temperature on developmental stages of the cold-stenothermal rotifer *Notholca caudata* Carlin. *Verh. internat. Ver. Limnol.* 23: 2016-2018
- LEAH, R. T., Br. MOSS & D. E. FORREST, 1980. The role of predation in causing major changes in the limnology of a hyper-eutrophic lake. *Internat. Rev. ges. Hydrobiol.* 65: 223-247
- LECHNER, M., 1966. Untersuchungen zur Embryonalentwicklung des Rädertieres *Asplanchna girodi* de Guerne. *Arch. Entwicklungsmech. Org.* 157: 117-173
- LESCHER-MOUTOUÉ, F., J. GARNIER & R. POURRIOT, 1985. Interactions entre les peuplements planctoniques et piscicole du lac de Créteil: impact d'une reproduction exceptionnelle de Percidés. *Bull. Ecol.* 16: 9-17
- LEENTVAAR, R., 1960. Waarnemingen in het Biesbosch-complex. *Lev. natuur* 63: 276-283
- LEENTVAAR, P., 1961. Quelques rotateurs rares observés en Hollande. *Hydrobiologia* 18: 245-251
- LEENTVAAR, P., 1963a. Stiefvader Rijn, kunstmoeder Maas. *Natura* 60: 50-56
- LEENTVAAR, P., 1963b. Dune waters in the Netherlands. I. Quackjeswater, Breede water en Vogelmeer. *Acta bot. neerl.* 12: 498-520
- LEENTVAAR, P., 1963c. De hydrobiologische toestand van het Laarder Wasmeer in 1961. *Biol. Jb. Dodonaea* 31: 284-300
- LEENTVAAR, P., 1965. Hydrobiologische waarnemingen in het plassengebied van NW-Overijssel. I. *Biol. Jb. Dodonaea* 33: 243-266

- LEENTVAAR, P., 1967a. Duinmeren II: Zwanewater, Muy, Oerd en Van Hunenplak. *Biol. Jb. Dodonaea* 35: 228-266
- LEENTVAAR, P., 1967b. Observations in guanotropic environments. *Hydrobiologia* 29: 441-489
- LEENTVAAR, P., 1967c. The artificial Brokopondo Lake of the Suriname River. Its biological implications. *Atas simp. biota amaz.* 3: 127-140
- LEENTVAAR, P., 1969. De zuidelijke Vechtplassen, flora en fauna. Een overzicht van de hydrobiologische toestand in de Vechtplassen in 1957 en 1958. *RIVON-verh.* 7: 81-133
- LEENTVAAR, P., 1970. Opmerkingen bij de Erpewaaien. *Lev. natuur* 73: 129-135
- LEENTVAAR, P., 1975. Hydrobiological observations in Surinam with special reference to the man-made Brokopondo Lake. *Stud. Fauna Surin. & other Guyanas* 15, n° 56: 1-173
- LEENTVAAR, P., 1979. Aditions and corrections to the Brokopondo study. (Surinam). *Amazoniana* 6: 521-528
- LEENTVAAR, P., 1980. Note on some Brachionidae (Rotifera) from the Netherlands, in DUMONT, H. J. & J. GREEN (ed.): *Proc. 2nd intern. Rotifer Symp.*: 259-262
- LEIGHTON, G., 1981. Biomass relationships between phytoplankton, zooplankton and aerobic bacteria in a reservoir. *Verh. internat. Ver. Limnol.* 21: 962-966
- LEMLY, A. D. & J. F. DIMMICK, 1982. Structure and dynamics of zooplankton communities in the littoral zone of some North Carolina lakes. *Hydrobiologia* 88: 299-307
- LEWIS, W. M., 1977. Feeding selectivity of a tropical *Chaoborus* population. *Freshw. Biol.* 7: 311-325
- LEWKOWICZ, M., 1971. Biomass of zooplankton and production of some species of Rotatoria and *Daphnia longispina* in carp ponds. *Pol. Arch. Hydrobiol.* 18: 215-223
- LEWKOWICZ, M., 1972. Rare and new species of rotifera in the fauna of Poland. *Acta Hydrobiol.* 14: 143-148
- LEWKOWICZ, M., 1984. Effect of intensive carp culture on zooplankton community. *Verh. internat. Ver. Limnol.* 22: 1695-1698
- LI, J. L. & H. W. LI, 1979. Species-specific factors affecting predator-prey interactions of the copepod *Acanthocyclops vernalis* with its natural prey. *Limnol. Oceanogr.* 24: 613-626
- LICHARDOVA, A. N., 1964. Contribution à l'étude des rotifères des eaux du bras Pecenské du Danube près de Bratislava. *Techn. Water* 8: 519-532
- LIKENS, G. E. & J. J. GILBERT, 1970. Notes on quantitative sampling of natural populations of planktonic Rotifers. *Limnol. Oceanogr.* 15: 816-820
- LINDSTRÖM, K., 1983. Changes in growth and size of *Keratella cochlearis* (Gosse) in relation to some environmental factors in cultures. *Hydrobiologia* 104: 325-328
- LINDSTRÖM, K. & B. PEJLER, 1975. Experimental studies on the seasonal variation of the Rotifer *Keratella cochlearis* (Gosse). *Hydrobiologia* 46: 191-197
- LIPSEY, L. L. & S. MALCOLM, 1981. Summer zooplankton communities of selected borrow-pit ponds in Northern Illinois. *Hydrobiologia* 77: 81-85
- LÖFFLER, H., 1962. Zur Systematik und Oekologie der chilenischen Süßwasserentomostraken. *Beitr. neotrop. Fauna* 2: 143-222
- LÖFFLER, H., 1969. High Altitude Lakes in Mt. Everest Region. *Verh. internat. Ver. Limnol.* 17: 373-385
- LÖFFLER, H., 1981. The winter condition of Lake Niriz in Southern Iran. *Verh. internat. Ver. Limnol.* 21: 528-534
- LOPEZ, T., J. TOJA & N. A. GABELLONE, 1991. Limnological comparison of two peridunal ponds in the Doñana National Park (Spain). *Arch. Hydrobiol.* 120: 357-378
- LOZHUCKOVA, A. N., G. L. VASSIL'JEVA & D. I. STORM, 1981. Toxicity of Polyphenols for Rotatoria and lower Crustacea. *Hydrobiol. Jnl.* 17 (6): 67-71
- LUBZENS, E., R. FISHLER & V. BERDUGO-WHITE, 1980. Induction of sexual reproduction and resting egg productions in *Brachionus plicatilis* reared in sea water. *Hydrobiologia* 73: 55-58
- LUBZENS, E., G. KOLODNY, S. PERRY, N. GALAI, R. SHESHINSKI & Y. WAX; 1990. Factors affecting survival of Rotifers (*Brachionus plicatilis* Müller O. F.) at 4 degrees C. *Aquacult.* 91: 23-47
- LUCIANI, A., J. L. CHASSÉ & P. CLEMENT, 1983. Aging in *Brachionus plicatilis*: the evolution of swimming as a function of age at two different calcium concentrations. *Hydrobiologia* 104: 141-146
- LYNCH, M., B. MONSON, M. SANDHEINRICH & L. WEIDER, 1981. Size-specific mortality rates in zooplankton populations. *Verh. internat. Ver. Limnol.* 21: 363-368

- LYUBIMOVA, T. S., 1974. Effects of fertilization on zooplankton growth in carp nursery ponds in the Southern Urals. *Hydrobiol. Jnl.* 10 (1): 24-29
- Mc GULLOUGH, J. D. & R. D. LEE, 1980. An ecological study of the rare rotifer species *Trochosphaera solstitialis* (THORPE, 1893) and the first report of the male. *Hydrobiologia* 71: 7-18
- MAC ISAAC, H. & J. J. GILBERT, 1990. Letter to the editor: Does exploitative or interference competition from *Daphnia* limit the abundance of *Keratella* in Loch Leven? A reassessment of MAY & JONES (1980). *J. Plankton Res.* 12: 1315-1322
- MC LAREN I. A., 1964. Zooplankton of Lake Haze, Ellesmere Island, and a nearby pond, with special reference to the copepod *Cyclops scutifer* Sars. *Can. J. Zool.* 42: 613-629
- MADALINSKI, K., 1961. Moss dwelling Rotifers of Tatra streams. *Polsk. Arch. Hydrobiol.* IX (XXII): 243-263
- MÄEMETS, A. & A., (1963?). The Neeruti Lakes. *Eesti NCV Teaduste Akadeemia juures asuva Eesti Looduseuurijate seltsi Aastaraamat*. 64: 92-136
- MÄEMETS, A., 1983. Rotifers as indicators of lake types in Estonia. *Hydrobiologia* 104: 357-361
- MÄEMETS, A. & L. KUTIKOVA, 1979. A new rotifer *Ploesoma peipsiense* sp. nov. in Lake Peipsi. *Eeste NSV teaduste Akadeemia toimetised* 28. *Koide Biologia* n° 2: 98-101
- MAES, L., 1985. The rotifer fauna of a polluted lowland brook-pond system during an aeration experiment. *Biol. Jb. Dodonaea* 53: 210-232
- MAGNIEN, R. E. & J. J. GILBERT, 1983. Diel cycles of reproduction and vertical migration in the rotifer *Keratella crassa* and their influence on the estimation of population dynamics. *Limnol. Oceanogr.* 28: 957-969
- MAKAREWICZ, J. & G. E. LIKENS, 1975. Niche analysis of a zooplankton community. *Science* 190: 1000-1003
- MAKAREWICZ, J. & G. E. LIKENS, 1979. Structure and function of the zooplankton community of Mirror Lake, New Hampshire, USA. *Ecol. Monogr.* 49: 109-127
- MALLEY, D. F. & P. S. S. CHANG, 1991. Early observations in the zooplankton community of a Precambrium Shield lake, receiving experimental additions of cadmium. *Verh. internat. Ver. Limnol.* 24: 2248-2253
- MALLEY, D., P. S. S. CHANG, D. L. FINDLAY & G. A. LINSEY, 1988. Extreme perturbation of the zooplankton community of a small Precambrium Shield lake by the addition of nutrients. *Verh. internat. Ver. Limnol.* 23: 2237-2247
- MALONE, B. J. & D. J. Mc QUEEN, 1983. Horizontal patchiness in zooplankton populations in the Ontario kettle lakes. *Hydrobiologia* 99: 101-124
- MAMARIL, A. C. Sr. & C. H. FERNANDO, 1978. Freshwater zooplankton of the Philippines (Rotifera, Cladocera and Copepoda). *Nat. and Appl. Sci. Bull.* 30: 109-221
- MANDAL, B. K., 1976. The fluctuation and distribution of plankton in a freshwater fishpond at Burdwan, West Bengal, India. *Limnologica* 11: 9-16
- MARGARITORA, F. G., D. CROSETTI, M. GIGLI & F. LOMBARDI, 1981. Prime osservazioni sulla struttura e distribuzione della biocenosi di un laghetto carsico: il lago superiore di Percile (Lazio). *Riv. Idrobiol.* XX: 673-687
- MARINONE, M. C. & M.-E. ZAGARESE, 1991. A field and laboratory study on factors affecting polymorphism in the rotifer *Keratella tropica*. *Oecologica* 86: 372-377
- MARKIC, M., 1983. The Rotatoria-Monogononta of the River Drava in Slovenia, Yugoslavia. *Hydrobiologia* 104: 229-230
- MARLIER, G. & C. WATTIEZ, 1977. Les étangs de Mirwart, in DUVIGNEAUD, P. & P. KESTEMONT: Productivité biologique en Belgique. SCOPE: *Travaux de la Section Belge du Programme Biologique International*: 419-452
- MARSH, Cl. E., J. Mc. GRAW Jr. & R. C. HARREL, 1978. Rotifer population in a southeast Texas oxbow lake with emphasis on cyclomorphosis of *Keratella cochlearis*. *Southwest Nat.* 23: 633-640
- MARTENS, A., 1985. Seizoionale en verticale distributie van zooplankton in een eutroof reservoir in Oost-Vlaanderen. *Lic. thesis, RUG*
- MARTIN, P., 1961. Le plancton du Lac Léman (Inventaire). *Schw. Zeitschr. Hydrol.* 23: 462-493
- MARTINEZ, Cl. C. & S. JOSÉ DE PAGGI, 1988. Especies de *Lecane* Nitzsch (Rotifera, Monogononta) en ambientes acuaticos del Chaco Oriental y del velle aluvial del rio Parana (Argentina). *Rev. Hydrobiol. trop.* 21: 279-295
- MARX, M., E. PANN, C. SORESCU & C. DRAGU, 1967. Contributii la studiul hidrobiologic al lacului Victoria (Reg. Oltina). *Hidrobiol. Bucur.* 8: 99-117

- MATSUMARA-TUNDISI, T., A. C. RIETZLER & J. G. TUNDISI, 1989. Biomass (dry weight and carbon content) of plankton crustacea from Broa-reservoir (Sao Carlos-S. P.) and its fluctuations across one year. *Hydrobiologia* 179: 229-236
- MATSUMARA-TUNDISI, T. & J. G. TUNDISI, 1976. PLankton studies in a lacustrine environment. I. Preliminary data on zooplankton ecology of Broa Reservoir. *Oecologia* 25: 265-270
- MATSUMARA-TUNDISI, T., J. G. TUNDISI, A. SAGGIO, A. OLIVEIRA NETO & E. G. ESPINDOLA, 1991. Limnology of Samuel Reservoir (Brazil, Rondônia) in the filling phase. *Verh. internat. Ver. Limnol.* 24: 1482-1488
- MATVEEVA, L. K., 1983. Community structure of planktonic rotifers in a mesotrophic lake. *Hydrobiologia* 104: 353-356
- MATVEEVA, L. K., 1986. Pelagic Rotifers of Lake Glubokoje from 1897 to 1984. *Hydrobiologia* 184: 45-54
- MATVEEVA, L. K., 1991. Pelagic rotifers as indicators of lake trophic state? *Verh. internat. Ver. Limnol.* 24: 2761-2763
- MAY L., 1983. Rotifer occurrence in relation to water temperature in Loch Leven, Scotland. *Hydrobiologia* 104: 311-315
- MAY, L., 1985. The use of procaine hydrochloride in the preparation of rotifer samples for counting. *Verh. internat. Ver. Limnol.* 22: 2987-2990
- MAY, L., 1986. Rotifer sampling - a complete list of species from one visit? *Hydrobiologia* 134: 117-120
- MAY, L., 1987. Effects of incubation temperature on the Hatching of rotifer resting eggs collected from sediments. *Hydrobiologia* 147: 335-338
- MARTIN, J. H., 1965. Phytoplankton - zooplankton relationships in Narragansett Bay. *Limnol. Oceanogr.* 10: 185-191
- MEINEL, W., U. MATTHIAS & B. SCHADE, 1983. Zooplankton- und Zoobenthosuntersuchungen im Litoral einer neu entstandenen Talsperre (Twistetalsperre, Nordhessen). *Arch. Hydrobiol.* 97: 145-162
- MENDIS, A. S. & C. H. FERNANDO, 1962. Guide to the freshwater fauna of Ceylon. *Bull. Fish. Stn. Ceylon* 12: 32-39
- MENZEL, R., 1968. Zur ökologie eines Kolkes während der Sommerstagnation. *Arch. Hydrobiol.* 65: 100-123
- MENZEL, R. & F. ROTH, 1972. Spektrale Phototaxis von Planktonrotatorien. *Experientia* 28: 356-357
- MESHKOVA, T. M. & A. A. NIKOGOSYAN, 1978. Changes in the zooplankton in connection with eutrophication of Lake Seven. *Hydrobiol. Jnl.* 14 (6): 16-19
- MICHAEL, R. G., 1968a. Studies on the Zooplankton of a tropical fish pond. *Hydrobiologia* 32: 47-68
- MICHAEL, R. G., 1968b. A new rotifer, *Conochilus madurai*, sp. n., from an astatic pool in Madurai, South India. *Zool. Anz.* 177: 439-441
- MICHAEL, R. G., 1969. Seasonal trend in physiochemical factors and plankton of a freshwater fishpond and their role in fish culture. *Hydrobiologia* 33: 144-160
- MICHELANGELLI, F., E. ZOPPI DE ROA & R. POURRIOT, 1979-1980. Rotifères en savanes inondables de Mantecal (Venezuela). *Cah. O. R. S. T. O. M. - Sér. hydrobiol.* 13: 47-59
- MILLER, G. W. & F. W. RABE, 1969. A limnological comparison of two small Idaho Reservoirs. *Hydrobiologia* 33: 523-546
- MIKSCHI, E., 1989. Rotifer distribution in relation to temperature and oxygen content. *Hydrobiologia* 186/187: 209-214
- MIKULSKI, J. St., 1964. Some biological features of perchpike lakes. *Verh. internat. Ver. Limnol.* 15: 151-157
- MINKOFF, G., E. LUBZENS & D. KAHAN, 1983. Environmental factors affecting hatching of rotifer (*Brachionus plicatilis*) resting eggs. *Hydrobiologia* 114: 61-69
- MIRABDULLAEV, I. M., 1992. On species of the genus *Lophocharis* Ehrenberg, 1838 (Rotifera: Monogononta) from Uzbekistan. *Hydrobiologia* 245: 163-168
- MIRACLE, M. R., 1974. Niche structure in freshwater zooplankton: a principal components approach. *Biology* 55: 1306-1316
- MIRACLE, M. R., 1975. Segregation of zooplankton populations in several depressions within one lake basin. *Verh. internat. Ver. Limnol.* 19: 1133-1139
- MIRACLE, M. R., 1977a. Migration patchiness and distribution in time and space of planktonic rotifers. *Arch. Hydrobiol. Beih.* 8: 19-37
- MIRACLE, M., 1977b. Epidemiology in rotifers. *Arch. Hydrobiol. Beih.* 8: 138-141

- MIRACLE, M. R., 1978. Composicion especifica de las comunidades zooplanktonicas de 153 lagos de los Pireneos y su interes biogeografico. *Oecol. aquat.* 3: 167-191
- MIRACLE, M. R., 1981. Biogeografia del zooplankton dels llacs del Pireneus. *Trab. Inst. Cat. Hist. Nat.* 9: 147-154
- MIRACLE, M. R., 1982. Biogeography of the freshwater zooplanktonic communities of Spain. *Jnl. Biogeogr.* 9: 455-467
- MIRACLE, M. R. & A. GUISET, 1977. Some effects of enclosure on congeneric species of rotifers. *Arch. Hydrobiol. Beih.* 8: 94-97
- MIRACLE, M. R., M. SERRA, R. OLTRA & E. VICENTE, 1988. Differential distribution of *Brachionus* species in three coastal lagoons. *Verh. internat. Ver. Limnol.* 23: 2006-2015
- MIRACLE, M. R., M. SERRA, E. VICENTE & C. BLANCO, 1987. Distribution of *Brachionus* species in Spanish mediterranean wetland. *Hydrobiologia* 147: 75-81
- MIRACLE, M. R. & E. VICENTE, 1983. Vertical distribution and rotifer concentrations in the chemocline of meromictic lakes. *Hydrobiologia* 104: 258-267
- MIRON, I., 1960. Contributions à l'étude des invertébrés aquatiques de la Bistritze et de ses affluents dans la zone du lac de barrage de Bicaz. *An. Sti. Univ. Jasi, ser. nova, sect. II, St. Nat.* 6: 719-728
- MITISKA, J. et collab., 1962. An investigation og the Jordan reservoir, Bohemia. *Technol. Water (Prague)* 6: 437-485
- MITRAKHOVICH, P. A., V. P. LYAKNOVICH & S. A. BOYKOVA, 1983. Zooplankton of Lake Lakoul' under the influence of thermal pollution. *Hydrobiol. Jnl.* 19 (4): 54-58
- MIZUNO, T. & S. MORI, 1970. Preliminary hydrobiological survey of some southeast inland waters. *Biol. J. Limnol. Soc.* 2: 77-117
- MODENUTTI, B. E., 1987. Caracterizacion y variacion espacial del zooplancton del Avroyo Rodriguez (Provincia de Buenos Aires, Argentina). *An. Inst. cienc. del Mar y Limnol. Univ. Auron. Mexico* 14: 21-28
- MODENUTTI, B. E. & M. C. CLAPS, 1988. Monogononta Rotifers from Plankton and Peryphyton of Pampasic Lotic Environments (Argentina). *Limnologica* 19: 167-175
- MOGIL'CHENKO, V. I., 1986. Some aspects of the feeding of larval commercial fishes in Kanev Reservoir. *Hydrobiol. Jnl.* 22 (4): 34-39
- MOMMAERTS, J. P., 1966. Caractérisation écologique de trois écosystèmes lacustres de Haute-Belgique. *Bull. Soc. r. Bot. Belg.* 99: 227-261
- MOORE, J. W., 1978a. Composition and structure of zooplankton communities in eighteen arctic and subarctic lakes (Canada). *Intern. Rev. ges. Hydrobiol.* 63: 545-565
- MOORE, J. W., 1978b. Some factors influencing the density and birth rate of three subarctic rotifer populations. *Arch. Hydrobiol.* 83: 251-271
- MOORE, J. W., 1980. Seasonal cycle of zooplankton and related phytoplankton development in three shallow, mesotrophic lakes in Northern Canada. *Internat. Rev. ges. Hydrobiol.* 65: 357-378
- MORALES-BAQUERO, R., 1987. Rotifer fauna of lakes and ponds over 2500 m above sea level in the Sierra Nevada, Spain, with description of a new subspecies. *Hydrobiologia* 147: 97-101
- MORALES-BAQUERO, R., 1988. Body size and variability of *Euchlanis dilatata* Ehrb. in high mountain lakes of Sierra Nevada (Spain). *Arch. Hydrobiol.* 112: 597-610
- MORALES-BAQUERO, R., L. CRUZ-PIZZARO & P. CARRILLO, 1988. Biomass and production of *Echlanis dilatata* in high mountain lakes and ponds of Sierra Nevada (Spain). *J. Plankt. Res.* 10: 1065-1075
- MORALES-BAQUERO, R., L. CRUZ-PIZARRO & P. CARRILLO, 1989. Patterns in the composition of the rotifer communities from high mountain lakes and ponds in Sierra Nevada (Spain). *Hydrobiologia* 186/187: 215-221
- MORDUKHAI-BOLTOVSKOI, Ph. D., 1979a. Composition and distribution of Caspian Fauna in the light of modern data. *Internat. Rev. ges. Hydrobiol.* 64: 1-38
- MORDUKHAI-BOLTOVSKOI, Ph. D., 1979b. The River Volga and its Life: 416-418. Vol. 33 of the *Monographiae biologicae*. W. Junk, The Hague, NL
- MORENO, L., H. GARCIA, L. PACHECO, H. SEGERS & A. INFANTE, 1992. Rotifers from Nicaragua. *Acta Cient. Venezol.* 43: 243-247
- MORLING, G. & B. PEJLER, 1990. Acidification and zooplankton development in some West-Swedish lakes 1966-1983. *Limnologica* 20: 307-318

- MOUREAU, Z., J. P. SCOHY, B. GODDEERIS, C. WATTIEZ, S. WARTEL & V. METENS, 1988. L'étang de Virelles et son bassin versant: qualité des eaux. *Studiedocumenten K. B. I. N.* n° 50: 59 pp.
- MOURELATOS, S., R. POURRIOT & C. ROUGIER, 1990. Taux de filtration du Rotifère *Brachionus calyciflorus*: comparaison des méthodes de mesure; influence de l'âge. *Vie et Milieu* 40: 39-43
- NAEF, J. & P. MARTIN, 1985. Plancton du Léman (X). Année 1984. *Arch. Sc. Genève* 38: 177-200
- NAIDENOW, W., 1963. Das Zooplankton der Donau vor dem Bulgarischen Ufer. *Bull. Inst. Zool. Mus. Bulg.* 13: 147-156
- NAIDENOW, W. & F. WAWRIK, 1984. Zur Kenntnis der Rotatorien der Teiche des niederösterreichischen Waldviertels. *Limnologica* 15: 157-177
- NAIDU, K. V., 1967. A contribution to the rotatorian fauna of South India. *J. Bombay nat. Hist. Soc.* 64: 384-388
- NARITA, T. & S. MORI, s.d. Secundary production of zooplankton. JIBP-PF Research group of Lake Biwa. Productivity of Freshwwater communities in Lake Biwa. *Contribution from the Otsu Hydrobiological Station*, University of Kyoto, n° 251
- NAUWERCK, A., 1966. Beobachtungen über das Phytoplankton klarer Hochgebirgsseen. *Schweiz. Zeitschr. Hydrol.* 28: 4-28
- NAUWERCK, A., 1978. Notes on the Planktonic Rotifers of Lake Ontario. *Arch. Hydrobiol.* 84: 269-301
- NAYAR, C. K. G., 1964. Morphometric studies on the rotifer *Brachionus calyciflorus* Pallas. *Curr. Sci.* 33: 469-470
- NAYAR, C. K. G., 1965a. Taxonomic notes on the Indian species of *Keratella* (Rotifera). *Hydrobiologia* 26: 457-462
- NAYAR, C. K. G., 1965b. Cyclomorphosis of *Brachionus calyciflorus*. *Hydrobiologia* 25: 538-544
- NAYAR, C. K. G., 1968. Rotifer fauna of Rajasthan, India. *Hydrobiologia* 31: 168-185
- NAYAR, C. K. G., 1970. Studies on the Rotifer population of two ponds at Pilani, Rajasthan. *J. Zool. Soc. India* 22: 21-34
- NAYAR, C. K. G. & K. K. N. NAIR, 1969. A collection of Brachionid rotifers from Kerala. *Proc. Ind. Acad. Sci., Sect. B*, 69: 223-233
- NEAGU-GODEANU, M., G. A. VASILIU & St. GODEANU, 1968. Contributions to the knowledge of the plankton from Obretin-Lake. *Trav. Mus. Hist. nat. "Grigore Antipa"* 8: 251-263
- NEILL, W. E., 1984. Regulation of rotifer densities by crustacean zooplankton in an oligotrophic montane lake in British Columbia. *Oecologia* 61: 175-181
- NEUMANN-LEITAO, S., M. N. PARANAGUA & J. L. VALENTIN, 1992. The planctonic rotifers of the estuarine lagunar complex of Suape (Pernambuco, Brazil). *Hydrobiologia* 232: 133-143
- NIKOLAYEV, I. I., 1972. Seasonal biological structure of Lake Onega. *Verh. intern. Ver. Limnol.* 18: 542-547
- NIKOLAYEV, I. I. & N. A. PETROVA, 1978. Indication of anthropogenic eutrophication of Lake Ladoga. *Hydrobiol. Jnl.* 14 (5): 1-8
- NIPKOW, F., 1961. Die Rädertiere im Plankton des Zürichsees und ihre Entwicklungsphasen. *Schweiz. Hydr. Zeitschr.* 23: 398-461
- NOGRADI, T. F., 1962. On the Rotifers and Limnology of an Hungarian bog pond. *Can. J. Zool.* 40: 677-684
- NOGRADI, Th., 1976. Canadian Rotifers. I. Lac Echo, Québec. *Nat. Can.* 103: 425-436
- NOGRADI, T., 1980. Canadian Rotifers. II. Parc Mont Tremblant, Québec. *Hydrobiologia* 71: 35-46
- NOGRADY, T., 1982. Correlation of rotifer associations in a chain of lakes fed by reclaimed sewage. *Hydrobiologia* 89: 277-284
- NOGRADY, T., 1983. Some new and rare warmwater rotifers. *Hydrobiologia* 106: 107-114
- NOGRADY, T., 1988. The littoral rotifer plankton of the Bay of Quinte (Ontario) and its horizontal distribution as indicators of Trophy. I. A full season study. *Arch. Hydrobiol. Suppl.* 79: 145-165
- NOGRADY, T., 1989. Rotifer associations of some wetlands in Ontario, Canada. *Hydrobiologia* 186/187: 223-228
- NOGRADY, T. & J. P. SMOLL, 1989. Rotifers from five high arctic ponds (Cape Herschel, Ellesmere Island, NWT). *Hydrobiologia* 173: 231-242
- NOLL, M., 1972. Besiedlungsfolgen in fluszwasser-beschickten Grundwasser-Anreichungsbecken. *Arch. Hydrobiol.* 70: 355-378
- NOVOTNA-DVORAKOVA, M., 1963. See DVORAKOVA, A. M., 1963

- O'CONNELL, M. & C. W. ANDREWS, 1977. Plankton ecology in Long Pond, St. John's, Newfoundland: a polluted pond characterized by a high flushing rate. *Internat. Rev. ges. Hydrobiol.* 62: 133-152
- ODEN, B. J., 1979. The freshwater littoral meiofauna in a South Carolina reservoir receiving thermal effluents. *Freshw. Biol.* 9: 191-304
- ODERMATT, J., 1970. Limnologie des Lauerzersees (Plankton). *Schweiz. Zeitschr. Hydrol.* 32: 1-75
- OEHMS, M. & A. SEITZ, 1992. Population dynamics and vertical distribution of pelagic rotifers in oligotrophic maar lakes. *Arch. Hydrobiol. Beih.* 38: 193-208
- OHLE, W., 1981. Photosynthesis and chemistry of an extremely acidic bathing pond in Germany. *Verh. internat. Ver. Limnol.* 21: 1172-1177
- OKINO, T. & Y. SATOH, 1986. Morphology, physics, chemistry and biology of Lake Rara in West Nepal. *Hydrobiologia* 140: 125-133
- OLIVEIRA, R., T. MONTEIRO, C. CABECADAS, C. VALE & M. J. BROGEIRA, 1985. A mine waste discharge rich in copper - an example of planktonic communities. *Verh. internat. Ver. Limnol.* 22: 2395-2404
- OLIVIER, S. R., 1965. Rotiferos planctonicos de Argentina. *Revista Mus. Plata (NS), T. 8. Zoologia* n° 63: 177-260
- OLRIK, K., S. LUNDER & K. RASMUSSEN, 1984. Interactions between phytoplankton, zooplankton and fish in the nutrient rich shallow Lake Hjarbaek Fjord, Denmark. *Internat. Rev. ges. Hydrobiol.* 69: 389-405
- OLSSEN, H., P. BLOMQUIST & H. OLOFSSON, 1992. Phytoplankton and zooplankton community structure after nutrient additions to the oligotrophic lake Hecklan, Sweden. *Hydrobiologia* 243/244: 147-155
- OLTRA, R. & M. R. MIRACLE, 1984. Comunidades zooplanctonicas de la Albufera de Valencia. *Limnetica* 1: 51-61
- OMALY, N., 1968. Le Lac Pavin. Historique et hydrobiologie. *Rev. Sc. nat. Auvergne* 34: 7-31
- OMALY, N., 1969. Etude biologique du Lac Tazenat. Dynamique des populations planctoniques au cours de l'année 1968. *Ann. Stat. biol. Besse-en-Chandesse* 4: 103-152
- OMALY, N., 1970. Première contribution à l'étude des bassins de la Haute-Tarentaine. Le Lac de Crégut et ses annexes: évolution, microendémisme chez les Calanides. *Ann. Stat. biol. Besse-en-Chandesse* 5: 303-347
- OMALY, N. & J. DÉVAUX, 1971. Observations écologiques sur trois petits étangs de la région de Besse-en-Chandesse (Puy-de-Dôme). *Rev. Sci. nat. Auvergne* 37: 23-45
- OOSTEN, T. D. R., 1972. Diverse poldersloten in het Spaarnewouderprojekt 1971-1972. *Doct. dissertation Zool. Lab. univ. Amsterdam*
- OPALINSKI, K. W., 1972. Freshwater fauna and flora in Haswell Island (QueenMary Land, Eastern-Antarctica). *Pol. Arch. Hydrobiol.* 19: 377-381
- ORCUTT, J. D. Jr. & M. L. PACE, 1984. Seasonal dynamics of rotifer and crustacean zooplankton populations in a eutrophic monomictic lake, with a note on rotifer sampling techniques. *Hydrobiologia* 119: 73-80
- OULAD ALI H., N. LAIR & H. AYADI, 1990. The seasonal succession of rotifers in the eutrophic Aydat lake. Spatio-temporal organisation. *Ann. Sci. nat. Zoologie, Paris*, 13e série, Vol. 11: 113-121
- PAASIVIRTA, L. & J. SÄRKÄÄ, 1978. Effects of pulp mill and municipal effluents and humus load on the macro- and microzoobenthos of some Finnish lakes. *Verh. internat. Ver. Limnol.* 20: 1779-1788
- PACE, M. L. & J. D. ORCUTT, 1981. The relative importance of protozoans, rotifers and crustaceans in a freshwater zooplankton community. *Limnol. Oceanogr.* 26: 822-830
- PAGGI, J. C., 1973. Contribucion al conocimiento de los Rotiferos dulceacuicolas planctonicos de la Republica Argentina. *Physis Sect. B., Aguas Cont. org.* 32 (85): 321-330
- PAGGI, J. C., 1978. Sobre la presencia de *Trochosphaera aequatorialis* Semper y *Horaëlla thomassoni* Koste (Rotatoria, Testudinellidae) en lagunas del valle de inundacion del Rio Parana. *Rev. Asoc. Cienc. Nat. Litoral* 9: 77-82
- PAGGI, J. C., 1980. Campaña limnologica "Keratella I" en el Rio Parana media (Argentina): zooplancton de ambientes lenticos. *Ecología* 4: 77-88
- PAGGI, J. C., 1981. Observaciones sobre el zooplancton de lagunas lagos de la Patagonia extraandina. I. Rotiferos. *Studies neotrop. Fauna and Environments* 16: 23-33

- PAGGI, J. C., 1986. Limnological studies in the Potter Peninsula, 25 de May Island, South Shetland Islands: biomass and spatial distribution of plankton. *Biomass Scientific Series* 7: 175-191
- PAGGI, J. C. & S. J. DE PAGGI, 1974. Primeros estudios sobre el zooplancton de las aguas loticas del Parana medio. *Physis Sect. B., Aguas cont. org.* 33 (86): 91-114
- PANT, M. C., A. JOSHI & P. C. SHARMA, 1985. Species composition, temporal abundance and community structure of zooplankton in Lake Sat. Tal (U. P.), India. *Arch. Hydrobiol.* 102: 519-535
- PARCHUK, G. V., 1985. Zooplankton of the Soviet Section of the Danube. *Hydrobiol. Jnl.* 21 (2): 5-10
- PARISE, A., 1961. Sur les genres *Keratella*, *Synchaeta*, *Polyarthra* et *Filinia* d'un lac italien. *Hydrobiologia* 18: 121-135
- PARISE, A., 1964. Prove di allevamento di rotiferi. *Ric. Sci.* 34 (II8): 465-480
- PARISE, A., 1966. The genus *Euchlanis* (Rotatoria) in the Marsh of Fucecetio (Central Italy), with description of a new species. *Hydrobiologia* 27: 328-336
- PASHA, S. M. K., 1961. On a collection of Fresh Water Rotifers from Madras. *J. Zool. Soc. India* 13: 50-55
- PASHKOVA, B. V., 1992. Diurnal variations of near-bottom zooplankton in the shallows of Kakhova Reservoir. *Hydrobiol. Jnl.* 28 (5): 50-59
- PATIL, C. S. & B. Y. M. GONDER, 1985. Ecological study of freshwater zooplankton of a subtropical pond (Karnataka Pond, India). *Internat. Rev. ges. Hydrobiol.* 70: 259-267
- PAULI, H.-R., 1989. A new method to estimate individual dry weight of rotifers. *Hydrobiologia* 186/187: 355-361
- PAWLOWSKI, L. K., 1960. Rotifères nouveaux et rares parmi la faune de la Pologne. *Bull. Soc. sci. lett. Lodz.* , 11: 1-15
- PAWLOWSKI, L. K., 1968. Nouvelles observations sur les Rotifères de la rivière Grabia. *Soc. Scient. Lodz. Nauk Math.-Przyrodn. TN* 103. 52 pp.
- PAWLOWSKI, L. K., 1970. Les Rotifères de la rivière Grabia. Variations observées au cours d'un cycle de vingt-quatre heures dans la composition de la faune des Rotifères. *Soc. Scient. Lodz. Nauk Math.-Przyrodn. TN* 110. 127 pp.
- PAWLOWSKI, L. K., 1973. Les Rotifères de la Rivière Grabia dans le profil longitudinal. *Pr. Wydz. math.-przyr. Lodz TN* 116. 43 pp.
- PEDROS-ALIO, C. & T. D. BROCK, 1983. The impact of zooplankton feeding on the epilimnetic bacteria of a eutrophic lake. *Freshw. Biol.* 13: 227-239
- PEJLER, B., 1961. The zooplankton of Ösbysjön, Djursholm. I. Seasonal and vertical distribution of the species. *Oikos* 12: 225-248
- PEJLER, B., 1962a. On the taxonomy and ecology of benthic and periphytic Rotatoria (Investigations in Northern Swedish Lapland). *Zool. Bidr. Uppsala* 33: 327-422
- PEJLER, B., 1962b. Morphological studies on the genera *Notholca*, *Kellicottia* and *Keratella* (Rotatoria). *Zool. Bidr. Uppsala* 33: 295-309
- PEJLER, B., 1962c. On the variation of the Rotifer *Keratella cochlearis* (Gosse). *Zool. Bidr. Uppsala* 35: 1-17
- PEJLER, B., 1962d. Taxonomic notes on some planktic freshwater rotifers. *Zool. Bidr. Uppsala* 35: 307-319
- PEJLER, B., 1962e. *Notholca caudata* Carlin (Rotatoria), a new presumed glacial relict. *Zool. Bidr. Uppsala* 33: 453-457
- PEJLER, B., 1964. Regional-ecological studies of swedish freshwater zooplankton. *Zool. Bidr. Uppsala* 36: 407-515
- PEJLER, B., 1965. Regional-ecological studies of Swedish freshwater plankton. *Zool. Bidr. Uppsala* 36: 407-515
- PEJLER, B., 1972. Rotifer plankton in brackish and freshwater localities in Central Sweden. *Oikos* 23: 416-419
- PEJLER, B., 1974. PLanktonic rotatoria from Björnöya (Bear Island, Arctic Ocean). *Nor. Polarinst. Årbok* 1972: 69
- PEJLER, B., 1977. On the global distribution of the family Brachionidae (Rotatoria). *Arch. Hydrobiol. Suppl.* 53: 255-306
- PEJLER, B., 1980. Variation in the genus *Keratella*. *Hydrobiologia* 73: 207-213
- PEJLER, B. & B. BĚRZINŠ, 1989. On choice of substrate and habitat in brachionid rotifers. *Hydrobiologia* 186/187: 137-144

- PENNAK, R. W., 1966. Structure of zooplankton populations in the littoral macrophyte zone of some Colorado lakes. *Trans. Amer. microsc. Soc.* 85: 329-349
- PENNAK, R. W., 1978. Anomalous primary production conditions in some Colorado alpine lakes, USA. *Verh. internat. Ver. Limnol.* 20: 434-437
- PERSOONE, G., 1964. The importance of fouling in the harbour of Ostend in 1964. *Helgol. wiss. Meeresuntersuch.* 12: 444-448
- PERSOONE, G. & N. DE PAUW, 1968. Pollution in the harbour of Ostend (Belgium). Biological and hydrographical consequences. *Helgol. wiss. Meeresuntersuch.* 17: 302-320
- PERSOONE, G., P. SORGELOOS, C. GROLIÈRE & N. OMALY, 1968. Sur la limnologie de quelques lacs du Puy-de-Dôme: physico-chimie et plancton de surface. *Ann. Stat. biol. Besse-en-Chandesse* 3: 43-114
- PERSSON, G., 1978. Experimental lake fertilization in the Kuokkel area, northern Sweden: the response by the planktonic rotifer community. *Verh. internat. Ver. Limnol.* 20: 875-880
- PESCHECK, E., 1961. Beiträge zur Biologie der Salzbecken im Neusiedler Seegebiet. *Verh. internat. Ver. Limnol.* 14: 1124-1131
- PETROVA, T. A., 1976. Zooplankton from the lower reaches of the Irgiz and Turgay Rivers in Western Kazakhstan. *Hydrobiol. Jnl.* 12 (3): 51-54
- PFANNKUCHE, O., H. JELINEK & E. HARTWIG, 1975. Zur Fauna eines Süßwasserwattes im Elbe-Aestuar. *Arch. Hydrobiol.* 76: 475-498
- PILARSKA, J., 1972. The dynamics of growth of experimental populations of the Rotifer *Brachionus rubens* Ehrb. *Pol. Arch. Hydrobiol.* 19: 265-277
- PILARSKA, J., 1977. Eco-physiological studies on *Brachionus rubens* Ehrb. (Rotatoria). I. Food selectivity and feeding rate. *Pol. Arch. Hydrobiol.* 24: 319-328
- PINEL-ALLOUL, B., P. LEGENDRE & E. MAGNIN, 1979. Zooplancton limnétique de 46 lacs et 17 rivières du territoire de la Baie de James. *Can. J. Zool.* 57: 1693-1709
- PINEL-ALLOUL, B., E. MAGNIN & G. CODIN-BLUMER, 1982. Effects de la mise en eau du réservoir Desaulniers (Territoire de la Baie de James) sur le zooplancton d'une rivière et d'une tourbière réticulée. *Hydrobiologia* 86: 271-296
- PINTO, J., 1992. Zooplankton distribution in the Bolivian part of the lake, in DEJOUX, C. & A. ILTIS (ed.): Lake Titicaca: 268-275. *Monographiae biologicae* 68. Kluwer Academic Publishers, Dordrecht, NL
- PIVODA, B., 1977. Migration of planktonic rotifera in Lunzer Obersee (Austria). *Arch. Hydrobiol. Beih.* 8: 50-52
- PIYASIRI, S. & J. K. U. JAYAKODY, 1991. Ecology of zooplankton in Victoria Reservoir, Sri Lanka. I. Composition and population structure of the zooplankton. *Verh. internat. Ver. Limnol.* 24: 1430-1435
- POLISHCHUK, L. N., 1976. Zooplankton of Dniester Estuary and the surrounding coastal area, exposed to anthropogenic influences. *Hydrobiol. Jnl.* 12 (6): 28-35
- POLISHCHUK, V. V., 1987. Zooplankton of the River Mouth Areas of Liev Reservoir. *Hydrobiol. Jnl.* 23 (5): 71-81
- POLISHCHUK, V. V., 1991. New information on the zooplankton of Lake Nohel. *Hydrobiol. Jnl.* 27 (3): 11-18
- POMAZKOVA, G. I. & O. M. KOZHOVA, 1976. Zoocenoses of artificial ponds in the purification plant at the Baikal Cellulose Plant. *Hydrobiol. Jnl.* 12 (6): 36-42
- PONTIN, R. M., 1989. Opportunist Rotifers: colonising species of young ponds in Surrey, England. *Hydrobiologia* 186/187: 229-234
- PONTIN, A. J. & M. PONTIN, 1980. Bullhousen Pond, Bisley, 1979. *Newsletter 55* (sept. 1980) of the Surrey Trust for Nature Conservation: 8-9
- POPCHENKO, V. I., L. V. LOMAKINA & I. I. POPCHENKO, 1981. Phytophilous assemblages of organisms in Saratov Reservoir. *Hydrobiol. Jnl.* 17 (2): 10-14
- POPESCU-GORJ, A. & E. COSTEA, 1961. Recherches hydrobiologiques et piscicoles dans les étangs d'Olténa (vallée inondable du Danube). *Hidrobiol. Bucur.* 2: 29-125
- POPESCU-MARINESCU, V., 1974. Das Zooplankton in den Donaumündungen und im Vordelta während des Jahres 1970. *Arch. Hydrobiol. Suppl. Bd.* 44: 372-382
- POPESCU-MARINESCU, V., N. NICOLESCU, D. NICOLESCU, V. ZINEVICI, E. PRUNESCU-ARION & L. TEODORESCU, 1990. L'évolution des biocénoses planctoniques et benthoniques des eaux du Danube, aux km 80 et 62, 900, pendant les années 1981-1985. *Rev. roum. biol.-Biol. anim.* 35: 49-67

- POPESCU-MARINESCU, V. & V. ZINEVICI, 1967. Forschungen über die Planktonbiozönose der Röhrichte mit beständigem Wasser (Donau-delta). *Hidrobiol. Bucur.* 8: 65-79
- POPOVSKA-STANKOVIC, O., 1990. Zooplankton of the Doiran Lake. *Poseb. izd. Prirodonauc. muz. Mak.* 9: 7-124
- POURRIOT, R., 1960. Recherches sur l'écologie du Rotifère *Eosphora najas* Ehrb. *Hydrobiologia* 16: 309-322
- POURRIOT, R., 1964. Etude expérimentale de variations morphologiques chez certaines espèces de Rotifères. *Bull. Soc. zool. France* 89: 555-561
- POURRIOT, R., 1965. Recherches sur l'écologie des Rotifères. Thèse de Doctorat. *Suppl. n° 21 à "Vie et Milieu".* 210 pp.
- POURRIOT, R., 1966. Quelques espèces de Rotifères nouvelles pour l'Europe. *Arch. Hydrobiol.* 61: 476-481
- POURRIOT, R., 1970. Quelques *Trichocerca* (Rotifères) et leurs régimes alimentaires. *Ann. Hydrobiol.* 1: 155-171
- POURRIOT, R., 1971. Deux formes nouvelles du genre *Squatinnella* (Rotifères) observées en Sologne. *Cah. Natural. Bull. N. P.*, n. s. 27: 97-101
- POURRIOT, R., 1972. Etude hydrobiologique de deux petits étangs de prairie. Observations sur la distribution de la température et du plancton et l'influence d'un couvert végétal de *Lemna minor*. *Ann. Hydrobiol.* 3: 33-46
- POURRIOT, R., 1973. Rapports entre la température, la taille des adultes, la longueur des oeufs et le taux de développement embryonnaire chez *Brachionus calyciflorus* Pallas (Rotifère). *Ann. Hydrobiol.* 4: 103-115
- POURRIOT, R., 1974. Relations prédateur-proie chez les Rotifères: influence du prédateur (*Asplanchna brightwelli*) sur la morphologie de la proie (*Brachionus bidentata*). *Ann. hydrobiol.* 5: 43-55
- POURRIOT, R., 1975a. Rotifères des Antilles. *Cah. ORSTOM, sér. hydrobiol.* 9, n° 2: 81-90
- POURRIOT, R., 1975b. Relations prédateur-proie: réactions adaptives et fluctuations des populations du zooplancton sous l'influence d'une prédation sélective. *Ann. biol.* 14: 69-86
- POURRIOT, R., 1976. Réflexions sur les Rotifères en tant qu'indicateurs biologiques. *Bull. franç. piscicult.* 48, n° 260: 148-152
- POURRIOT, R., D. BENEST, P. CHAMP & C. ROUGIER, 1982. Influence de quelques facteurs du milieu sur la composition et la dynamique saisonnière du zooplancton de la Loire. *Acta Oecol. Génér.* 3: 351-371
- POURRIOT, R., D. BENEST, P. CLEMENT & C. ROUGIER, 1984. Morphologie comparée d'oeufs de durée de Brachonidés (Rotifères). *Bull. Soc. zool. France* 109: 231-239
- POURRIOT, R. & P. CLEMENT, 1974. Photopériodisme et cycle hétérogamique chez *Notommata copeus* (Rotifère). III. Recherche du seuil minimal d'éclairement. *Arch. zool. exp. gén.* 115: 641-650
- POURRIOT, R. & P. CLEMENT, 1975. Influences du groupement et de la densité de population sur le cycle de reproduction de *Notommata copeus* (Rotifère). I. Mise en évidence et essai d'interprétation. *Arch. zool. exp. & gén.* 116: 375-422
- POURRIOT, R. & M. DELUZARCHES, 1971. Recherches sur la biologie des rotifères. II. Influence de la température sur la durée de développement embryonnaire et post-embryonnaire. *Annls. limnol.* 7: 25-52
- POURRIOT, R. & L. LEBORGNE, 1970. Teneurs en protéines, liptides et glucides de zooplancton d'eau douce. *Ann. hydrobiol. INRA* 1: 171-178
- POURRIOT, R. & C. ROUGIER, 1976. Influence de l'âge des parents sur la production de femelles mictiques chez *Brachionus calyciflorus* Pallas et *Brachionus rubens* Ehrb. (Rotifères). *C. R. Acad. Sci.* 283 D: 1497-1500
- POURRIOT, R. & C. ROUGIER, 1979. Influences conjuguées du groupement et de la qualité de la nourriture sur la reproduction de *Brachionus plicatilis* (Rotifère). *Neth. J. Zool.* 29: 242-264
- POURRIOT, R. & C. ROUGIER, 1986. Rythmes de reproduction de femelles sexuées chez le Rotifère *Brachionus calyciflorus* en élevage à température constante. *Bull. Soc. zool. Fr.* 111: 203-207
- POURRIOT, R. & C. ROUGIER, 1991. Importance volumétrique des oeufs chez les Rotifères planctoniques. *Ann. Limnol.* 27: 15-24
- POURRIOT, R., C. ROUGIER & D. BENEST, 1980. Hatching of *Brachionus rubens* O. F. Müller resting eggs (Rotifera). *Hydrobiologia* 73: 51-54
- POURRIOT, R., C. ROUGIER & D. BENEST, 1981. Rôle de la lumière et de la température dans l'éclosion des oeufs de durée de *Brachionus rubens* Ehrb. (Rotifère). *Neth. J. Zool.* 31: 637-649

- POURRIOT, R., C. ROUGIER & D. BENEST, 1986. Qualité de la nourriture et contrôle de la mixis chez le Rotifère *Brachionus rubens* Ehrb. *Bull. Soc. zool. France* 111: 105-111 et 208-209
- POURRIOT, R., G. ROUYER & M. PELTIER, 1972. Prolifération de Rotifères épiphytes et pollution thermique dans la Loire. *Bull. fr. Piscicult.* 44, n° 244: 111-118
- POURRIOT, R. & T. W. SNELL, 1983. Resting eggs in Rotifers. *Hydrobiologia* 104: 213-224
- POURRIOT, R., M. TASSIGNY & J. JUNQUA, 1969. Etude des variations saisonnières de trois biotopes aquatiques et de leurs biocénoses aquatiques. *Rech. hydrobiol. cont.* 1: 61-95
- POURRIOT, R. & E. ZOPPI DE ROA, 1981. Dicranophoridae (Rotifères Monogonontes) du Venezuela. *Bull. Soc. zool. France* 106: 195-199
- PRAY, F. A., 1965. Studies on the early development of the rotifer *Monostyla cornuta* Müller. *Trans. Amer. micr. Soc.* 84: 210-216
- PREISZLER, K., 1977a. Horizontal distribution and "avoidance of shore" by rotifers. *Arch. Hydrobiol. Beih.* 8: 43-46
- PREISZLER, K., 1977b. Do rotifers show "avoidance of shore"? *Oecologia* 27: 253-260
- PREISZLER, K., 1980. Field experiments on the optical orientation of pelagic rotifers. *Hydrobiologia* 73: 199-203
- PRIDDLE, J. & H. J. G. DARTNALL, 1978. The biology of an antarctic aquatic moss community. *Freshw. Biol.* 8: 469-480
- PRINS, R. & J. DAVIS, 1966. The fate of planktonic rotifers in a polluted stream. *Occasional papers of the C. C. Adams Center for Ecological Studies*, n° 15: 14 pp.
- PRUNESCU-ARION, E., 1967. Date preliminare asupra planctonului riului Moldava. *Hidrobiol. Bucur.* 9: 75-81
- PRUNESCU-ARION, E., 1989. Strukturelle Aspekte betreffend die Rotiferen aus der Donau bei ihrem Eintritt ins Delta (Rumänischer Abschnitt) in der Zeitspanne 1981-1985. *Rev. roum. Biol.-Biol. anim.* 34: 49-52
- RACHYUNAS, L. A., 1971. Distribution of invertebrates in the cooling pond of a Lithuanian regional power station. *Hydrobiol. Jnl.* 7 (1): 68-72
- RADWAN, S., 1966. Espèces des Rotifères (Rot.) nouvelles pour la faune de la Pologne, leur distribution et écologie. *Ann. Univ. M. C. SKLODOWSKA, Sect. C*, 21: 121-130
- RADWAN, S., 1967. Apparition des Rotifères (Rotatoria) du genre *Testudinella* Bory de St. Vincent, 1826 dans la faune de Pologne. *Ann. Univ. M. C. SKLODOWSKA, Sect. C*, 22: 41-56
- RADWAN, S., 1968. Espèces rares et intéressantes de Rotifères des Lacs de Sosnowickie. *Pol. Arch. Hydrobiol.* 15: 237-249
- RADWAN, S., 1969. Rotifères (Rotatoria) de trois lacs à trophisme différent. *Pol. Arch. Hydrobiol.* 16: 51-65
- RADWAN, S., 1971. On some new for European fauna and rare species of Rotifers. *Ann. Univ. M. C. SKLODOWSKA, Sect. C*, 26: 169-175
- RADWAN, S., 1974. The rotifer (Rotatoria) species of the Parczew Region Peat Bogs. *Ann. Univ. M. C. SKLODOWSKA, Sect. C*, 29: 215-230
- RADWAN, S., 1975. Rare rotifer (Rotatoria) species in the pelagic zone of the Leczna and Włodawa Lakes. *Ann. Univ. M. C. SKLODOWSKA, Sect. C*, 30: 243-255
- RADWAN, S., 1976. Planctonic rotifers as indicators of lake trophy. *Ann. Univ. M. C. SKLODOWSKA, Sect. C*, 227-235
- RADWAN, S., 1978. Dynamics of production of the pelagic rotifers in three lakes of different trophism. *Verh. internat. Ver. Limnol.* 20: 1017-1021
- RADWAN, S., 1980. The effect of some biotic and abiotic factors on the fertility of planktonic rotifer species. *Hydrobiologia* 73: 59-63
- RADWAN, S., 1984. The influence of some abiotic factors on the occurrence of rotifers of Leczna and Włodawa Lake District. *Hydrobiologia* 112: 117-124
- RADWAN, S., C. KOWALCZYK, W. KOWALIK & W. ZWOLSKI, 1984. Littoral fauna in two lakes of different trophy (L. Bialskie nd L. Piaseczno, Eastern Poland). *Verh. internat. Ver. Limnol.* 22: 991-995
- RADWAN, S., C. KOWALCZYK, W. ZWOLSKI, W. KOWALIK & B. STEPIEN, 1985. Factors affecting nutrient budget in lakes of the R. Jorka watershed (Masurian Lakeland, Poland). VIII. Zoosyrthon in the tributaries and outlet of lake Jorzeć. *Ecol. Pol.* 33: 311-328
- RADWAN, S. & A. PALEOLOG, 1983. Notes on the rotifers of coal mine waters in Eastern Poland. *Hydrobiologia* 104: 307-309

- RADWAN, S. & B. POPIOLEK, 1977. Comparison of biomass and production of some species of Rotifers (Rotatoria) in spring and summer in the pelagic zone of three lakes in the Leczna-Wlodawa region. *Ann. Univ. M. C. SKLODOWSKA*, Sect. C, 22: 277-291
- RADWAN, S. & B. POPIOLEK, 1989. Percentage of rotifers in spring zooplankton in lakes of different trophy. *Hydrobiologia* 186/187: 235-238
- RADWAN, S., W. ZWOLSKI, C. KOWALCZYK, W. KOWALIK & A. PALEOLOG, 1991. Ecological characteristics of coal mine waters in Lublin Coal Basin (Eastern Poland) and their influence on the river zoocenosis. *Verh. internat. Ver. Limnol.* 24: 2156-2158
- RAI, H., 1974. Limnological studies on the River Yamuna at Delhi, India. Part II. The dynamics of potamoplankton populations in the River Yamuna. *Arch. Hydrobiol.* 73: 492-517
- RAO, G. Ch. & P. N. GANAPATI, 1968. The interstitial fauna inhabiting the beach sand of Waltair coast. *Proc. Nat. Inst. Sci. Ind.* 34: 82-125
- RAUH, F., 1963. Untersuchungen über die Variabilität der Rädertiere. III. Die experimentelle Beeinfluszung der Variation von *Brachionus calyciflorus* und *Brachionus capsuliflorus*. *Z. Morph. Oekol. Tiere* 53: 61-106
- RAVERA, O., 1969. Seasonal variation of the biomass and biocenotic structure of plankton in the Bay of Ispra (Lago Maggiore). *Verh. internat. Ver. Limnol.* 17: 237-254
- RAVICHANDRAU, S. & P. S. RAMANIBAE, 1988. Plankton and related parameters of Buckingham canal at Madras, India. - A canonical correlation analysis. *Arch. Hydrobiol.* 114: 117-132
- RAWSON, D. S., 1960. A limnological comparison of twelve large lakes in northern Saskatchewan. *Limnol Oceanogr.* 5: 195-211
- RECK, E., 1981. Faunistisch-ökologische Untersuchungen an Rotatorien eines eutrophen Teiches. *Decheniana* 134: 227-233
- REID, J. W. & P. N. TURNER, 1988. Planktonic rotifera, copepoda and cladocera from Lagos Açu and Viana, State of Maranhao, Brazil. *Rev. Brasil. Biol.* 48: 485-495
- RENAUD-DEBYSER, J. & B. SALVAT, 1963. Eléments de prospérité des biotopes des sédiments meubles intertidaux et écologie de leurs populations en microfaune et microflore. *Vie et Milieu* 14: 464-550
- RESTITUTO, F. & N. LAIR, 1976. Projet alpin O. C. D. E. pour la lutte contre l'eutrophisation. Lacs du massif central français. III. Le Lac Pavin, interrelations entre paramètres. *Ann. Stat. biol. Besse-en-Chandesse* 10: 146-191
- REY, J., 1984. Export de matériel zooplanctonique dans l'exutoire d'une retenue des Pyrénées. *Verh. internat. Ver. Limnol.* 22: 1493-1497
- REY, J., 1988a. Etude comparée de la dynamique du zooplankton de trois réservoirs d'altitude et d'un lac naturel dans les Pyrénées. *Annls. Limnol.* 24: 139-160
- REY, J., 1988b. Impact de la vidange sur le développement des populations zooplanctoniques dans une retenue des Pyrénées. *Annls. Limnol.* 24: 235-241
- REY, J. & J. CAPBLANC, 1975. Dynamique des populations et production du zooplankton du lac de Port-Bielh (Pyrénées centrales). *Annls. Limnol.* 11: 1-45
- RICHMAN, S., M. D. BAILIFF, L. J. MACKEY & D. W. BOLGRIEN, 1984. Zooplankton standing stock, species composition and size distribution along a trophic gradient in Green Bay, Lake Michigan. *Verh. internat. Ver. Limnol.* 22: 475-487
- RIEDL, R., 1970. Rotatoria in *Fauna und Flora der Adria* 220-222. Ed. P. Parey, Hamburg.
- RIEMANN, F., 1966. Die interstitielle Fauna im Elbe-Aestuar. Verbreitung und Systematik. *Arch. Hydrobiol./Suppl. Bd.* 31: 1-279
- RIGLER, F. H., 1978. Limnology in the high Arctic: a case study of Char Lake. *Verh. internat. Ver. Limnol.* 20: 127-140
- RIGLER, F. H., M. E. Mc CALLUM & J. C. ROFF, 1974. Production of zooplankton in Char Lake. *J. Fish Res. Bd. Can.* 31: 637-646
- RINGELBERG, J. & J. STEENVOORDEN, 1986. Diel variation in the egg ratio of rotifers throughout the season (preliminary report). *Hydrobiol. Bull.* 19: 153-158
- RITTERBUSCH, B., 1974. Zur Ökologie des tierischen Psammoms in einem Sickerbecken der Berliner Wasserwerke und seine Bedeutung für die Reinigung von infiltriertem Oberflächenwasser. *Z. angew. Zool.* 61: 301-346
- RIV'YER, I. K., S. S. BAKASTOV & A. I. SARALOV, 1981. Vertical distribution of plankton in the Mologa River in winter. *Hydrobiol. Jnl.* 17 (3): 11-21
- ROBOTTI, C., 1975a. Cyclomorphosis and amphigony in *Brachionus calyciflorus* Pallas (Rotatoria). *Experientia* 31: 524-525

- ROBOTTI, C., 1975b. Chromosome complement and male haploidy of *Asplanchna priodonta* Gosse, 1850 (Rotatoria). *Experientia* 31: 1270-1272
- ROCHE, K. F., 1987. Post-encounter vulnerability of some rotifer prey types to predation by the copepod *Acanthocyclops robustus*. *Hydrobiologia* 147: 229-233
- ROCHE, K. F., 1989. Predator-prey interactions between *Acanthocyclops robustus* (Copepoda: Cyclopoida) and some zooplankton prey. *Doctor's dissertation, RUG*
- RODEWALD-RUDESCU, L. & S. GODEANU, 1961. Die Rädertierfauna Rumäniens. VII. Neue und bemerkenswerte Rädertiere aus Rumänien. *Zool. Anz.* 167: 341-359
- ROFF, J. & R. KWIATKOWSKI, 1977. Zooplankton and zoobenthos communities of selected northern Ontario lakes of different acidities. *Can. J. Zool.* 55: 899-911
- ROGOZ, I., 1972. Recherches sur le plankton de la rivière d'Oltet et de ses principaux affluents. *Hidrobiol. Bucur.* 13: 115-123
- ROMO, S., 1990. Seasonal zooplankton patterns in a shallow oligotrophic lake: Loch Rusky (Scotland). *Annls. Limnol.* 26: 11-17
- RONNEBERGER, D., 1975. Zur Kenntnis der Grundwasserfauna des Saale-Einzugsgebiet (Thüringen). *Limnologica* 9: 323-419
- RONNEBERGER, D. & W. SCHÖNBORN, 1985. Rotatoria, in CASPERS, S. J.: *Lake Stechlin, a temperate, oligotrophic lake*: 505-508. Junk Publishers, Dordrecht, NL. 553 pp.
- ROSTAN, J. C., 1976. Dynamique du plancton rotiférien dans un étang piscicole de la Dombes. *Mém. doc. Ain* 1: 48-55
- ROTHHAUPT, K. O., 1985. A model approach to the population dynamics of the rotifer *Brachionus rubens*, in two-stage chemostat culture. *Oecologia* 65: 252-259
- ROTHHAUPT, K. O., 1990a. Differences in particle-size-dependant feeding efficiencies of closely related rotifer species. *Limnol. Oceanogr.* 35: 16-23
- ROTHHAUPT, K. O., 1990b. Changes of the functional responses of the rotifers *Brachionus rubens* and *Brachionus calyciflorus* with particle size. *Limnol. Oceanogr.* 35: 24-32
- ROTHHAUPT, K. O., 1991. The influence of toxic and filamentous blue-green algae on feeding and population growth of the rotifer *Brachionus rubens*. *Internat. Rev. ges. Hydrobiol.* 76: 67-72
- ROUGIER, C., 1969. Etat du peuplement rotatori du Léman pour l'anée 1967. *Ann. Stat. biol. Besse-en-Chandesse* 4: 26-29
- ROUGIER, C. & R. POURRIOT, 1977. Aging and the control of the reproduction in *Brachionus calyciflorus* Pallas (Rotatoria). *Exp. geront.* 9: 137-151
- RUDESCU, L. 1960. Fauna Republicii populare Romîne. Trochelminthes. Vol. II, fasc. II. Rotatoria. *Acad. R. P. R.* 1192 pp. 899figs.
- RUDESCU, L., 1961. Rotiferii din Marea Neagra. *Hidrobiol. Bucur.* 3: 283-329
- RUDESCU, L., 1966. Die Hochmoore der Poiana Stampei (Region Suceava) und ihre wissenschaftliche Bedeutung. *Hidrobiol. Bucur.* 7: 163-178
- RUDESCU, L., 1968. Die Rotatorien, Gastrotrichen und Tardigraden der Schilfrohrgebiete des Donaudeltas. *Hidrobiol. Bucur.* 9: 195-202
- RUDESCU, L. & A. C. BANU, 1973. Inundationsgebiete und Delta in ihrer Beziehung zum Donau-regime. *Arch. Hydrobiol./Suppl.* 44: 206-222
- RUDESCU, L. & V. MARINESCU-POPESCU, 1969. Beitrag zur Kenntnis des gegenseitigen Einflusses der Donau und des Schwarzen Meeres. *Verh. internat. Ver. Limnol.* 17: 1112-1121
- RUDOLF, K., 1966. Untersuchungen über die Ernährungsbeziehungen in der Freiwasserregion der Säidenbachsperrre. *Wiss. Z. K. K. -M. Univ. Leipzig, Math. Nat. R.* 15: 229-245
- RÜHMANN, D., 1961a. Beobachtungen über die Verbreitung der planktischen Rotatorien im groszhamburgischen Raum. *Abhandl. Verhandl. naturw. Ver. Hamburg N. F.* 5: 131-148
- RÜHMANN, D., 1961b. Das Rädertier *Brachionus diversicornis*. *Mikrokosmos* 50: 301-303
- RÜHMANN, D., 1962. Rädertiere im Plankton des Hamburger Hafens. *Mikrokosmos* 51: 294-297
- RÜHMANN, D., 1964. Rädertiere in Hamburger Gewässern. *Mikrokosmos* 53: 366-367
- RÜHMANN, D., 1965a. Ueber die Verbreitung der planktonischen Rädertiere im groszhamburgischen Raum. *Faun. Mitteil.* Bd. 2, H. 9: 221-222
- RÜHMANN, D., 1965b. Das Rädertier *Brachionus budapestinensis*. *Mikrokosmos* 54: 205-206
- RÜHMANN, D., 1967. Alster-Plankton unter dem Mikroskop. *Mikrokosmos* 56: 314-316
- RUMENGAU, I. F. M., H. KAYANO & K. HIRAYAMA, 1991. Karyotypes of S and L type rotifers *Brachionus plicatilis* O. F. Müller. *J. exp. Mar. Biol. Ecol.* 154: 171-176
- RUSSELL, C. R., 1960. An Index of the Rotatoria of New Zealand and Outlying Islands from 1859 to 1959. *Trans. roy. Soc. N. Zeal.* 88: 443-461

- RUTTNER-KOLISKO, A., 1964. Kleingewässern am Ostrand der persischen Salzwüste. Ein Beitrag zur Limnologie arider Gebiete. *Verh. internat. Ver. Limnol.* 15: 201-208
- RUTTNER-KOLISKO, A., 1966a. The influence of climatic and edaphic factors on small astatic waters in the East Persia salt desert. *Verh. internat. Ver. Limnol.* 16: 524-531
- RUTTNER-KOLISKO, A., 1966b. *Notholca lapponica*, nov. spec., eine psammophile Rotatorienart aus Nord-Skandinavian. *Anz. Oesterr. Akad. Wiss. Math. Naturw. Kl.* 9: 1-4
- RUTTNER-KOLISKO, A., 1970. *Synchaeta calva*, nov. spec., a new Rotifer from the English Lake District. *Internat. Rev. ges. Hydrobiol.* 55: 387-390
- RUTTNER-KOLISKO, A., 1971. Rotatorien als Indikatoren für den Chemismus von Binnensalzgewässern. *Ber. Oesterr. Akad. Wiss. Math. Naturw. Kl., Abt. I*, 179: 283-298
- RUTTNER-KOLISKO, A., 1972. Der Einfluss der Temperatur und Salzgehalt des Mediums auf Stoffwechsel- und Vermehrungsintensität von *Brachionus plicatilis* (Rotatoria). *Verh. deutsch. zool. Ges.* 65: 88-95
- RUTTNER-KOLISKO, A., 1975a. The influence of fluctuating temperature on plankton Rotifers. A graphical model based on life data of *Hexarthra fennica* from Neusiedlersee, Austria. *Symp. Biol. Hung.* 15: 197-204
- RUTTNER-KOLISKO, A., 1975b. The vertical distribution of plankton rotifers in a small alpine lake with sharp oxygen depletion (Lunzer Obersee). *Verh. internat. Ver. limnol.* 19: 1286-1294
- RUTTNER-KOLISKO, A., 1977a. Population dynamics of rotifers as related to climatic conditions in Lunzer Obersee and Untersee. *Arch. Hydrobiol. Beih.* 8: 88-93
- RUTTNER-KOLISKO, A., 1977b. Amphoteric reproduction in a population of *Asplanchna priodonta*. *Arch. Hydrobiol. Beih.* 8: 178-181
- RUTTNER-KOLISKO, A., 1977c. The effect of the microsporid *Plistophora asperospora* on *Conochilus unicornis* im Lunzer Untersee (LUS). *Arch. Hydrobiol. Beih.* 8: 135-137
- RUTTNER-KOLISKO, A., 1977d. Comparison of various sampling techniques, and results of repeated sampling of planktonic rotifers. *Arch. Hydrobiol. Beih.* 8: 13-18
- RUTTNER-KOLISKO, A., 1978. Influence of fluctuating temperature on plankton rotifers. II. Laboratory experiments. *Verh. internat. Ver. Limnol.* 20: 2400-2405
- RUTTNER-KOLISKO, A., 1979. Brackish water rotifers from the Mickletown Lagoons in the Lower Aire Valley. *Nat.* 104: 113
- RUTTNER-KOLISKO, A., 1980. The abundance and distribution of *Filinia terminalis* in various types of lakes as related to temperature, oxygen, and food. *Hydrobiologia* 73: 169-175
- RUTTNER-KOLISKO, I. & G. MALICKY, 1988. Succession and abundance of Rotifers in Lunzer Untersee as determinated by biological interactions and abiotic events. *Verh. internat. Ver. Limnol.* 23: 510-515
- SAETER, O. A., 1971. Phytoplankton and zooplankton of some lakes in Northeastern Norway. *Schweiz. Z. Hydrol.* 33: 200-220
- SAKSENA, D. N. & N. KULKARNI, 1986. On the Rotifer fauna of two sewage channels of Gwalior (India). *Limnologica* 17: 139-148
- SAKSENA, D. N. & S. P. SHARMA, 1986. Morphological form variation in a Loricate Rotifer, *Keratella tropica* APSTEIN from a perennial pond, Janaktal, Gwalior, India. *Internat. Rev. ges. Hydrobiol.* 71: 283-288
- SAKSENA, S. B. & A. D. ADONI, 1973. Diurnal variation in Sagar Lake, Sagar (India). I. Studies in the deep water area. *Hydrobiologia* 43: 535-543
- SALAZKIN, A. A., 1971. Zooplankton in oligotrophic lakes of the Numid zone in the Northwestern USSR. *Hydrobiol. Jnl.* 7 (3): 23-28
- SALNIKOV, L. P. & N. G. BADNINA, 1979. Zooplankton in the ponds of the Kochubey Fishfarm (Dagestanian ASSR). *Hydrobiol. Jnl.* 16 (1): 72
- SALT, G. W., 1977. An analysis of the diets and reproductive performances of five sympatric species of *Asplanchna*. *Arch. Hydrobiol. Beih.* 8: 123-125
- SALT, G. W., 1989. Comparisons of the diets and reproductive performances of two sympatric rotifers, *Asplanchna girodi* and *Asplanchna priodonta*. *Freshw. Biol.* 22: 417-430
- SALT, G. W., G. F. SAKBADINI & M. L. COMMINS, 1978. Trophi morphology relative to food habits in six species of rotifers (Asplanchnidae). *Trans. Am. Micr. Soc.* 97: 469-485
- SANCHEZ, L., E. VASQUEZ & L. BLANCO, 1985. Limnological investigations of the rivers Ura-coa, Yato, Morichal Largo and Claro in the Eastern Plains of Venezuela. *Verh. internat. Ver. Limnol.* 22: 2153-2160

- SANDERS, R. W., K. G. PORTER, S. J. BENNETT & A. E. DE BIASE, 1989. Seasonal pattern of bacterivory by flagellates, ciliates, rotifers and cladocerans in a freshwater planktonic community. *Limnol. Oceanogr.* 34: 673-687
- SANDGREN, C. D. & J. T. LEHMAN, 1990. Response of chlorophyl, phytoplankton and microzooplankton to the invasion of Lake Michigan by *Bythotrephes*. *Verh. internat. Ver. Limnol.* 24: 386-392
- SANDSTRÖM, O., 1979. The horizontal distribution of Baltic epi- and metalimnic zooplankton. *Internat. Rev. ges. Hydrobiol.* 64: 251-262
- SARKAR, M. L. & H. RAI, 1964. The limnology of Suray Fund, India. I. Physico-chemical conditions and plankton. *Trans. Am. micr. Soc.* 83: 260-265
- SARMA, S. S. S., 1988. New records of freshwater rotifers (Rotifera) from Indian waters. *Hydrobiologia* 160: 263-269
- SARMA, S. S. S. & T. R. RAO, 1987. Effect of food level on body size and egg size in a growing population of the rotifer *Brachionus patulus* Müller. *Arch. Hydrobiol.* 111: 245-253
- SARMA, S. S. S. & T. R. RAO, 1991. The combined effects of food and temperature on the life history parameters of *Brachionus patulus* Müller (Rotifera). *Internat. Rev. ges. Hydrobiol.* 76: 225-239
- SARTORY, P. K. & E. D. HOLLOWDAY, 1983. On the occurrence of a rarely observed mutilation in the sessile rotifer *Stephanoceros fimbriatus* (Goldfuss) and its possible bearing on the identity of *Collotheca monoceros* (Zacharias). *Microscopy* 34: 507-510
- SAUNDERS, J. F., 1980. Diel patterns of reproduction in rotifer populations from a tropical lake. *Freshw. Biol.* 10: 35-39
- SAUNDERS, J. F., III & W. M. LEWIS Jr., 1988a. Zooplankton abundance and transport in a tropical white-water-river. *Hydrobiologia* 162: 147-155
- SAUNDERS, J. F. III & W. M. LEWIS Jr., 1988b. Composition and seasonality of the zooplankton of Lake Valencia, Venezuela. *J. Plankt. Res.* 10: 957-985
- SAUNDERS, J. F. III & W. M. LEWIS Jr., 1989. Zooplankton abundance in the lower Orinoco River, Venezuela. *Limnol. Oceanogr.* 34: 397-409
- SAUNDERS-DAVIES, A. P. & R. M. PONTIN, 1987. A centrifugation method for measuring the relative density (specific gravity) of planktonic Rotifers (Rotifera), with values for the relative density of *Polyarthra major* Burckhardt and *Keratella cochlearis* (Gosse). *Hydrobiologia* 147: 379-381
- SCHABER, P. & A. SCHRIMPFF, 1984. On morphology and ecology of the *Filinia-longiseta-terminalis* group (Rotatoria) in Bavarean and Tyrolean lakes. *Arch. Hydrobiol.* 101: 247-257
- SCHADEN, R., 1977a. Zur Zyklomorphose einiger Brachioniden (Rotatoria) Zentralamazoniens. *Amazoniana* 6: 257-266
- SCHADEN, R., 1977b. Jahreszeitliche Veränderungen in der Zusammensetzung des Rotatorienbestandes des Lago do Castanho, in Zentralamazonien. *Amazoniana* 6: 267-273
- SCHADEN, R., 1977c. Zur Diversität und Identität amazonischer Rotatorienzooome. *Amazoniana* 6: 347-371
- SCHEDA, S. M. & B. C. COWELL, 1988. Rotifer grazers and phytoplankton: seasonal experiments on natural communities. *Arch. Hydrobiol.* 114: 31-44
- SCHEPENS, D., 1960. Bijdrage tot de kennis van de Rotatoria van België. *Biol. Jb. Dodonea* 28: 88-95
- SCHINDLER, D. W., 1988. Experimental studies of chemical stressors on whole lake ecosystems. *Verh. internat. Ver. Limnol.* 23: 11-41
- SCHINDLER, D. W. & B. NOVEN, 1971. Vertical distribution and seasonal abundance of zooplankton in two shallow lakes of the experimental lake area, northwestern Ontario. *J. Fish. Res. Bd. Can.* 28: 245-256
- SCHLÜTER, A., 1984. Oekologische Untersuchungen an einem Stillgewässer im tropischen Regenwald von Peru, unter besonderer Berücksichtigung der Amphibien. *Doctor's dissertation*, Univ. Hamburg.
- SCHMID-ARAYA, J. M., 1991a. Distributional aspects of Rotifera in Central and South Chile. *Arch. Hydrobiol.* 120: 481-493
- SCHMID-ARAYA, J. M., 1991b. The effects of food concentration on the life histories of *Brachionus plicatilis* (O. F. M.) and *Encentrum linnhei* Scott. *Arch. Hydrobiol.* 131: 87-102
- SCHNEIDER, D. W., 1990. Direct aspects of the independent effects of exploitative and interference competition between *Daphnia* and rotifers. *Limnol. Oceanogr.* 35: 916-922

- SCHNITZLER, H., 1971. Oekologische Untersuchungen am PLankton der Riveristalsperre. *Arch. Hydrobiol.* 69: 60-94
- SCHOENBERG, St. A., 1990. Short-term productivity response of algae and bacteria to zooplankton grazing in two freshwater lakes. *Freshw. Biol.* 23: 395-410
- SCHROEVERS, P. J., 1962. Eutrofierung in een Drentse heiplas: het Mekelermeer. *Lev. Natuur* 65: 221-228
- SCHROEVERS, P. J., 1965. Hydrobiologische waarnemingen in NW Overijssel. II. Het bezinkingsplankton van het Kippenest bij Wanneperveen. *Biol. Jb. Dodonaea* 33: 267-342
- SCHULZ, H. 1961. Qualitative und quantitative Untersuchungen im Elbe-Aestuar. *Arch. Hydrobiol./Suppl.* 26: 5-105
- SCHWARZ, S., 1963. Produktionsbiologische Untersuchungen am Zooplankton der Rügenschen, Hiddenseer und Darszer Boddengewässer (1953-1955). II. Die Rotatorien. *Z. Fischerei, N. F.* 11: 641-672
- SCOTT, A. P. & S. M. BAYNES, 1978. Effect of algal diet and temperature on the biochemical composition of the rotifer *Brachionus plicatilis*. *Aquaculture* 14: 247-260
- SCOTT, J. M., 1974. A new marine rotifer of the genus *Encentrum*. Its morphology and cultivation. *Zool. J. Linn. Soc.* 54: 247-251
- SEBESTYEN, O., 1974. Present and past occurrence of members of the genus *Filinia* Bory de St. Vincent 1826 (Testudinellidae: Rotatoria) in Lake Balaton. *Annal. Biol. Tihany* 41: 289-315
- SEELBACH, P. W. & W. F. Mc DUFFET, 1983. Distribution and abundance of zooplankton in an alkaline marsh in Northumberland County, Pennsylvania. *Internat. Rev. ges. Hydrobiol.* 68: 379-395
- SEGERS, H., 1990. Contribution to the knowledge of the rotifer fauna of the Galapagos Islands. *Biol. Jb. Dodonaea* 58: 113-119
- SEGERS, H., 1991. Nomenclatural note on a primary homonym in the genus *Lecane* (Lecanidae, Rotifera). *Rev. Hydrobiol. trop.* 24: 77
- SEGERS, H., A. O. AJAJI, G. Y. CHIAMBENG, H. P. CHUAH, M. DEL CASTILLO, M. G. DIRECTO, M. LUZURIAGA DE CRUZ, L. MORENO, A. L. OLIVEIRA-NETO & Y. RETUANING-WIDYASTUTI, 1991. Fourteen rotifer species new to the Belgian fauna, with some nomenclatural and taxonomical remarks on some *Squatinnella*-species. *Belg. J. Zool.* 121: 193-201
- SEGERS, H., N. EMIR & J. MERTENS, 1992. Rotifera from north and northeast Anatolia (Turkey). *Hydrobiologia* 245: 179-189
- SELIN, P. & K. KOSKINEN, 1988. The sedimentation ponds as the water treatment system in the peat production areas and their effect on the water quality and plankton communities. *Verh. internat. Ver. Limnol.* 23: 1564-1571
- SENDACZ, S., 1984. A study of the zooplankton community of Billings Reservoir, São Paulo. *Hydrobiologia* 113: 121-127
- SERAFOVA-HADZISCE, J., 1974. Zooplankton in some lakes of the Aegean Lake Zone. *Fragmenta Balcanica* 9: 165-170
- SERGEYEVA, O. A., R. A. KALINICHENKO, L. G. LEUCHINA & Ye. V. MEDIANIK, 1989. Effect on the cooling system of a fossil power plant station on plankton. *Hydrobiol. Jnl.* 25 (6): 36-40
- SERRA, M. & M. R. MIRACLE, 1983a. Biometric analysis of *Brachionus plicatilis* ecotypes from Spanish lagoons. *Hydrobiologia* 104: 279-291
- SERRA, M. & M. R. MIRACLE, 1983b. Enzyme polymorphism of *Brachionus plicatilis* populations from several Spanish lagoons. *Abstracts 22nd Congress SIL*: 319
- SERRA, M. & M. R. MIRACLE, 1985. Enzyme polymorphism in *Brachionus plicatilis* populations from several lagoons. *Verh. internat. Ver. Limnol.* 22: 2991-2996
- SERRA, M. & M. R. MIRACLE, 1987. Biometric variation in three strains of *Brachionus plicatilis* as a direct response to abiotic variables. *Hydrobiologia* 147: 83-89
- SERRUYA, C & U. POLLINGHER, 1984. Lakes of the Warm Belt. *Cambr. Univ. Press.* 589 pp.
- SHARMA, B. K., 1976. Rotifers collected from North-West India. *Newsl. zool. Serv. India* 2: 255-258
- SHARMA, B. K., 1978a. Contribution to the rotifer fauna of West Bengal. Part I. Family Lecanidae. *Hydrobiologia* 57: 143-153
- SHARMA, B. K., 1978b. Two new Lecanid Rotifers from India. *Hydrobiologia* 60: 191-192
- SHARMA, B. K., 1978c. Contribution to the rotifer fauna of West Bengal. II. Genus *Lepadella* Bory de St. Vincent 1826. *Hydrobiologia* 58: 83-88

- SHARMA, B. K., 1979a. Rotifers from West Bengal. III. Further studies on the Eurotatoria. *Hydrobiologia* 64: 239-250
- SHARMA, B. K., 1979b. Rotifers from West Bengal. IV. Further contributions to the Eurotatoria. *Hydrobiologia* 65: 39-47
- SHARMA, B. K., 1979c. Further contributions to the Lecanid fauna (Rotifera: Lecanidae) of West-Bengal. *Acta Hydrobiol.* 21: 53-59
- SHARMA, B. K., 1980a. Contribution to the Rotifer fauna of Panjab State, India. I. Family Brachionidae. *Hydrobiologia* 76: 249-254
- SHARMA, B. K., 1980b. Contribution to the Rotifer fauna of Orissa, India. *Hydrobiologia* 70: 225-233
- SHARMA, B. K., 1983. The Indian species of the genus *Brachionus* (Eurotatoria: Monogononta: Brachionidae). *Hydrobiologia* 104: 31-39
- SHARMA, B. K., 1987a. Indian Brachionidae (Eurotatoria: Monogononta) and their distribution. *Hydrobiologia* 144: 269-275
- SHARMA, B. K., 1987b. The distribution of the Lecanid rotifers (Rotifera: Monogononta: Lecanidae) in North-Eastern India. *Rev. Hydrobiol. trop.* 20: 101-105
- SHARMA, B. K., 1990. The genus *Testudinella* (Eurotatoria: Gnesiotrocha: Testudinellidae) in North-Eastern India. *Hydrobiologia* 199: 29-33
- SHARMA, B. K. & R. G. MICHAEL, 1980. Synopsis of taxonomic studies on Indian Rotatoria. *Hydrobiologia* 73: 229-236
- SHARMA, B. K. & S. SHARMA, 1984. A note on some Eurotatoria from Punjab State, India. *Hydrobiologia* 109: 279-282
- SHARMA, B. K. & S. SHARMA, 1987. On species of genus *Lepadella* (Eurotatoria: Monogononta: Colurellidae) from North-Eastern India, with remarks on Indian taxa. *Hydrobiologia* 147: 15-22
- SHARMA, P. C. & M. C. PANT, 1984. Structure of a littoral zooplankton community of two Kumaun lakes (U. P.), India. *Limnologica* 16: 51-65
- SHARMA, P. C. & M. C. PANT, 1985. Species composition of zooplankton in two Kumaun Himalayan Lakes (U. P.), India. *Arch. Hydrobiol.* 102: 387-403
- SHARMA, P. C. & M. C. PANT, 1987. Limnology of a subtropical lake, Bhimtal (U. P.), India. *Limnologica* 18: 15-27
- SHARMA, S. P. & D. N. SAKSENA, 1984. Form variation in the Rotifer *Brachionus calyciflorus* Pallas from a perennial impoundment in India. *Internat. Rev. ges. Hydrobiol.* 69: 747-752
- SHIEL, R. J., 1979. Synecology of the Rotifera of the River Murray, South Australia. *Aust. J. Mar. Freshwater Res.* 30: 255-263
- SHIEL, R. J., 1981. Planktonic Rotifera of the Murray-Darling river system, Australia: endemism and polymorphism. *Verh. internat. Ver. Limnol.* 21: 1523-1530
- SHIEL, R. J., 1983. The genus *Brachionus* (Rotifera: Brachionidae) in Australia, with a description of a new species. *Proc. r. Soc. Vict.* 95: 33-37
- SHIEL, R., 1985. Zooplankton of the Darling River system, Australia. *Verh. internat. Ver. Limnol.* 22: 2136-2140
- SHIEL, R. J., 1986. Common zooplankters (excluding Protozoa) of the Murray-Darling System, in DAVIES, B. R. & K. F. WALKER: *The ecology of River Systems*: 661-677. Junk Publishers, Dordrecht, NL
- SHIEL, R. J. & W. KOSTE, 1979. Rotifera recorded from Australia. *Trans. r. Soc. S. Aust.* 103: 57-68
- SHIEL, R. J. & W. KOSTE, 1983. Rotifer communities of billabongs in northern and south-eastern Australia. *Hydrobiologia* 104: 41-47
- SHIEL, R. J. & W. KOSTE, 1985a. New species and new records of Rotifera (Aschelminthes) from Australian waters. *Trans. r. Soc. S. Austr.* 109: 1-15
- SHIEL, R. J. & W. KOSTE, 1985b. Records of Rotifers epizoic on cladocerans from S Australia. *Trans. r. Soc. S. Austr.* 109: 179-180
- SHIEL, R. J. & W. KOSTE, 1986. Australian Rotifera: Ecology and Biogeography, in DE DECKER, P. & W. D. WILLIAMS, (eds.): *Limnology in Australia*. C. S. I. R. O. Junk, Dordrecht, NL: 141-150
- SHIEL, R. J. & W. KOSTE, 1992. Rotifera from Australian inland waters. VIII. Trichocercidae (Monogononta). *Trans. r. Soc. S. Aust.* 116: 1-27
- SHIEL, R. J., W. KOSTE & W. TAN, 1989. Tasmania revisited: rotifer communities and habitat heterogeneity. *Hydrobiologia* 186/187: 239-245
- SHIEL, R. J., C. J. MERRICK & G. G. GAUF, 1987. The Rotifera of impoundments in Southeastern Australia. *Hydrobiologia* 147: 23-29

- SHIEL, R. J., K. F. WALKER & W. D. WILLIAMS, 1982. Plankton of the Lower River Murray, South Australia. *Aust. J. Mar. Freshw. Res.* 33: 301-327
- SHIEL, R. J. & K. F. WALKER, s.d. Zooplankton of regulated and unregulated rivers: the Murray-Darling River system, Australia, in A. LILLEHAMMER & S. J. SALTVETT: *Regulated rivers*: 263-270. Universitetsforlaget A. S. Oslo-Bergen-Stavanger-Tromso.
- SHOEMAKER, J., S. H. DAVIS & R. K. WILLIAMS, 1987. *Notommata allantois* in Northeastern Texas lakes: a rotifer previously only known from Europe. *Texas J. Sci.* 39: 194-196
- SIBBING, W., 1962. Studienwochen in der Biologischen Station "Heiliges Meer". *Praxis Naturwiss.* 11: 9-15
- SIEGFRIED, C. A., 1991. The pelagic rotifer community of an acidic clearwater lake in the Adirondack mountains of New York State. *Arch. Hydrobiol.* 122: 441-462
- SIEGFRIED, C. A., J. A. BLOOMFIELD & J. W. SUTHERLAND, 1989. Planktonic rotifer community structure in Adirondack, New York, USA lakes in relation to acidity, trophic state and related water quality characteristics. *Hydrobiologia* 175: 33-48
- SIEGFRIED, C. A. & M. E. KOPACHE, 1980. Feeding of *Neomysis mercedis* (Holmes). *Biol. Bull.* 159: 193-205
- SIEGFRIED, C. A. & J. W. SUTHERLAND, 1983. Lake acidification and the biology of Adirondack lakes. I. Rotifer communities. *Abstracts 22nd Congress SIL*: 321
- SIEGFRIED, C. A., J. W. SUTHERLAND, Sc. O. QUIN & J. A. BLOOMFIELD, 1984. Lake acidification and the biology of Adirondack lakes. I. Rotifer communities. *Verh. internat. Ver. Limnol.* 22: 549-558
- SLADECEK, V., 1961a. Contribution to the knowledge of the Rotifer *Ascomorpha minima* Hofsten. *Hydrobiologia* 18: 274-276
- SLADECEK, V., 1961b. Eine bemerkenswerte *Encentrum*-Art (Rot.). *Zool. Anz.* 167: 360-362
- SLADECEK, 1962a. Ueber die Excretophoren bei den festsitzenden Räderieren der Ordnung Collothecacea. *Zool. Anz.* 169: 488-491
- SLADECEK, V., 1962b. To the knowledge of the bioseston of the reservoir Vrané on the river Vltava. *Cas. nar. muz. Praha* 131: 76-80
- SLADECEK, V., 1963. Notes on the ecology of two sessile rotifera from acid waters. *Techn. Water* 7: 563-569
- SLADECEK, V., 1969. *Collotheca undulata* n. sp. *Zool. Anz.* 182: 417-420
- SLADECKOVA, A., 1960a. Limnological study of the reservoir Sedlice near Zeliv. XI. Periphyton stratification during the first year-long period (June 1957-July 1958). *Fac. Techn. Fuel & Water* 4: 143-261
- SLADECKOVA, A., 1960b. Application of the glass slide method to the periphyton study in the Slapy Reservoir. *Fac. Techn. Fuel & Water* 4: 403-433
- SLADECKOVA, A., 1962. Limnological study of the reservoir Sedlice near Zeliv. XX. Periphyton stratification during the second-year-long period (August 1958-June 1959). *Techn. Water* 6: 221-291
- SLADECKOVA, A., 1985. Hydrobiological problems in air-conditioning systems. *Verh. internat. Ver. Limnol.* 22: 2342-2346
- SLADECKOVA, A. & V. SLADECEK, 1960. Das Frühlingsplankton einiger Talsperren in Ost-Böhmen. *Fac. Techn. Fuel & Water* 4: 487-510
- SLADECKOVA, A. & V. SLADECEK, 1963. Periphyton as indicator of the reservoir water quality. I. True periphyton. *Techn. Water* 7: 507-561
- SMIRNOW, N. S., 1961. Food cycles in sphagnum bogs. *Hydrobiologia* 17: 175-182
- SMITH, M. W., 1969. Changes in environment and biota of a natural lake after fertilization. *J. Fish. Res. Bd. Can.* 26: 3101-3132
- SMYLY, W. J. P., 1968. Some observations on the effect of sampling technique under different conditions on numbers of some freshwater planktonic Entomostraca and Rotifera caught by a water-bottle. *J. Nat. Hist.* 2: 569-575
- SMYLY, W. J. P., 1976. Some effects of enclosures on the zooplankton in a small lake. *Freshw. Biol.* 6: 241-251
- SMYLY, W. J. P., 1978. Further observations on limnetic zooplankton in a small lake and two enclosures containing fish. *Freshw. Biol.* 8: 491-485
- SNELL, T. W., 1979. Intraspecific competition and population structure in Rotifers. *Ecology* 60: 494-502
- SNELL, T. W., 1980. Blue-green algae and selection in rotifer populations. *Oecologica* 46: 343-346

- SNELL, T. W., 1987. Sex, population dynamics and resting egg production in rotifers. *Hydrobiologia* 144: 105-111
- SNELL, T. W., 1989. Systematics, reproductive isolation and species boundaries in monogonont rotifers. *Hydrobiologia* 186/187: 299-310
- SNELL, T. W., C. J. BIEBERICH & R. FÜRST, 1983. The effect of green and blue-green algal diets on the reproductive rate of the rotifer *Brachionus plicatilis*. *Aquacult.* 31: 21-30
- SNELL, T. W., B. E. BURKE & S. D. MESSUR, 1983. Size and distribution of resting eggs in a natural population of the rotifer *Brachionus plicatilis*. *Gulf Res. Reports* 7: 285-287
- SNELL, T. W. & K. CARRILLO, 1984. Body size variation among strains of the rotifer *Brachionus plicatilis*. *Aquacult.* 37: 359-367
- SNELL, T. W. & C. A. HAWKINSON, 1983. Behavioral reproductive isolation among populations of the rotifer *Brachionus plicatilis*. *Evolution* 37: 1294-1305
- SNELL, T. W. & C. E. KING, 1977a. Amphoteric reproduction in *Asplanchna girodi*. *Arch. Hydrobiol. Beih.* 8: 182-183
- SNELL, T. W. & C. E. KING, 1977b. Lifespan and fecundity patterns in rotifers: the cost of reproduction. *Evolution* 31: 882-890
- SNELL, T. W., B. D. MOFFAT, C. JANSEN & G. PERSOONE, 1991. Acute toxicity tests using Rotifers. III. Effects of temperature, strain, and exposure time on the sensitivity of *Brachionus plicatilis*. *Environm. Toxic. Water Qual.* 6: 63-75
- SNELL, T. W. & B. C. WINKLER, 1984. Isozyme analysis of Rotifer Proteins. *Biochem. Syst. & Ecol.* 12: 199-202
- SOKOLOVA, V. A., 1986. Water quality of the Vilyuy River in relation to its zooplankton. *Hydrobiol. Jnl.* 22(1): 273-275
- SOLLBERGER, P. J. & L. J. PAULSON, 1992. Littoral and limnetic zooplankton communities in Lake Mead, Nevada-Arizona, USA. *Hydrobiologia* 237: 175-184
- SOLTERO, R. A., A. F. GASPERINO & W. G. GRAHAM, 1975. Cultural eutrophication of Long Lake, Washington. *Verh. internat. Ver. Limnol.* 19: 1778-1789
- SOROKIN, Y. I. & E. B. PAVELJEVA, 1972. On the quantitative characteristics of the pelagic ecosystem of Dalnee Lakes (Kamchatka). *Hydrobiologia* 40: 519-552
- SOTO, D., I. VILA & B. VILLALOBOS, 1984. Temporal and spatial distribution of rotifera in a Chilean reservoir: a possible effect of impoundment hydrodynamics. *Hydrobiologia* 114: 67-74
- SPANGENBERG, H.-J., 1973. Faunistisch-ökologische Untersuchungen an Gewässern von Gipshöhlen und in Grundwasser des Südharzes und Kyffhäuserns. *Internat. Rev. ges. Hydrobiol.* 58: 501-542
- SPITTNER, P., 1969. Zucht- und Fütterungsversuche an einigen Rädertierarten der Hiddenseer Gewässer. *Limnologica* 7: 207-211
- SPODNIEWSKA, I., E. GRYGIEREK & A. HILLBRICHT-ILKOWSKA, 1966. Some annual successional changes in plankton of temporal water bodies. *Verh. internat. Ver. Limnol.* 16: 585-591
- SPODNIEWSKA, I., A. HILLBRICHT-ILKOWSKA & T. WEGLENSKA, 1973. Long-term changes in the plankton of eutrophic Mikolajskie Lake as an effect of accelerated eutrophication. *Bull. Acad. Pol. Sci. Série Sci. biol.* C1. II, 21: 215-221
- SREENIVASAN, A., 1964. Limnological studies and fish yield in three upland lakes of Madras State, India. *Limnol. Oceanogr.* 9: 564-575
- SREENIVASAN, A., 1968. The limnology and fish productions in two ponds in Chinglepat (Madras). *Hydrobiologia* 32: 131-144
- SRIVASTAVA, O. N., 1981. Planktonic change after restoration of a perennial pond at Chotanagpur, India. *Verh. internat. Ver. Limnol.* 21: 1196-1199
- STANKOVIC, S., 1960. The Balkan Lake Ohrid and its living world. *Ed. Junk, The Hague, NL.* 358 pp.
- STARKWEATHER, P. L., 1988. Reproductive and functional responses of the rotifer *Brachionus plicatilis* to changing food density. *Verh. internat. Ver. Limnol.* 23: 2001-2005
- STARKWEATHER, P. L., 1990. Zooplankton community structure of high elevation lakes: biogeographic and predator-prey interactions. *Verh. internat. Ver. Limnol.* 24: 513-517
- STARKWEATHER, P. L. & K. G. BOGDAN, 1980. Detrital feeding in natural zooplankton communities: discrimination between live and dead algal food. *Hydrobiologia* 73: 83-85
- STARKWEATHER, P. L. & J. J. GILBERT, 1977. Feeding in the rotifer *Brachionus calyciflorus*. II. Effect of food density on feeding rates using *Euglena gracilis* and *Rhodotorula glutinis*. *Oecologia* 28: 133-139

- STARKWEATHER, P. L. & E. J. WALSH, 1989. Influence of cyanobacterial diet on *Asplanchna* predation risk in *Brachionus calyciflorus*. *Hydrobiologia* 186/187: 35-38
- STARMACH, K., 1961. Hydrobiologische Untersuchungen an der Talsperre bei Gazalkowice. *Verh. internat. Ver. Limnol.* 14: 643-646
- STELLA, E., L. FERRERO & F. G. MARGARITORA, 1978. Alterations of the plankton in a much polluted lake in Central Italy (Latium), the Volcanic Lake Nemi. *Verh. internat. Ver. Limnol.* 20: 1049-1054
- STELLA, E. & F. MARGARITORA, 1966. A contribution to the study of Lago Monterosi (Roma, Italy): present biological conditions of the lake. *Verh. internat. Ver. Limnol.* 16: 83-88
- STELLA, E. & F. MARGARITORA, 1972. Planktonic biocenosis of Giulianello Lake (Latium, Italy) during the last four years. *Verh. internat. Ver. Limnol.* 18: 482-488
- STEMBERGER, R. S., 1974. Temporal and spatial distributions of planktonic rotifers in Milwaukee Harbor and adjacent Lake Michigan. *Proc. 17th Conf. Great Lakes Res.*: 120-134
- STEMBERGER, R. S., 1976. New Rotifers from the Laurentian Great Lakes Region. *J. Fish Res. Bd. Can.* 33: 2814-2818
- STEMBERGER, R. S., 1979. A Guide to Rotifers of the Laurentian Great Lakes. *Rep. Univ. Michigan Biol. Station-Pellston, Michigan* 49769
- STEMBERGER, R. S., 1985. Prey selection by the copepod *Diacyclops thomasi*. *Oecologia* 65: 492-497
- STEMBERGER, R. S., 1986. The effects of food deprivation, prey density and volume on clearance and ingestion rates of *Diacyclops thomasi*. *J. Plankton Res.* 8: 243-251
- STEMBERGER, R. S., 1987. The potential for population growth of *Ascomorpha ecaudis*. *Hydrobiologia* 147: 297-301
- STEMBERGER, R. S., 1988. Reproductive costs and hydrodynamic benefits of chemically induced defenses in *Keratella testudo*. *Limnol. Oceanogr.* 33: 593-606
- STEMBERGER, R. S., 1990a. Food limitation, spination and reproduction in *Brachionus calyciflorus*. *Limnol. Oceanogr.* 35: 33-44
- STEMBERGER, R. S., 1990b. An inventory of rotifer species diversity of northern Michigan inland lakes. *Arch. Hydrobiol.* 118: 283-302
- STEMBERGER, R. S., 1990c. *Keratella armudara* (Rotifera: Brachionidae) a new rotifer from a Michigan bog lake. *Can. J. Zool.* 68: 2306-2309
- STEMBERGER, R. S. & M. S. EVANS, 1984. Rotifer seasonal succession and copepod predation in Lake Michigan. *J. Great Lakes Res.* 10: 417-428
- STEMBERGER, R. S. & J. E. GANNON, 1977. Multivariate analysis of rotifer distribution in Lake Huron. *Arch. Hydrobiol. Beih.* 8: 38-42
- STEMBERGER, R. S. & J. J. GILBERT, 1984. Spine development in *Keratella cochlearis*: induction by cyclopoid copepods and *Asplanchna*. *Freshw. Biol.* 14: 639-647
- STEMBERGER, R. S. & J. J. GILBERT, 1985. Body size, food concentration and population growth in planktonic rotifers. *Ecology* 66: 1151-1159
- STEMBERGER, R. S. & J. J. GILBERT, 1987. Multiple-species induction of morphological defenses in the rotifer *Keratella testudo*. *Ecology* 68: 370-378
- STENSON, J. A. E., 1982. Fish impact on rotifer community structure. *Hydrobiologia* 87: 57-64
- STENSON, J. A. E., 1983. Changes in the relative abundance of *Polyarthra vulgaris* and *P. dolichoptera*, following the elimination of fish. *Hydrobiologia* 104: 269-273
- STEPANOVA, L. A., 1972. Rations of *Mesocyclops leuckarti* (Claus) and *Leptodora kindtii* (Focke) populations in Lake Ilmen. *Hydrobiol. Jnl.* 8 (3): 70-72
- STEPHENSON, G. L., P. HAMILTON, N. K. KAUSHIK, J. B. ROBINSON & K. R. SOLOMON, 1984. Spatial distribution of plankton in enclosures of three sizes. *Can. J. Fish. Aquat. Sci.* 41: 1048-1054
- STERNER, R. W., 1990. Resource competition and the autecology of pennate diatoms. *Verh. internat. Ver. Limnol.* 24: 518-523
- STEWART, L. J. & D. G. GEORGE, 1987. Environmental factors influencing the vertical migration of planktonic rotifers in a hypereutrophic tarn. *Hydrobiologia* 147: 203-208
- STICH, H. B., 1991. Phosphorous and carbon values of zooplankton species in Lake Constance. *Verh. internat. Ver. Limnol.* 24: 837-846
- STONEBURGER, D. L. & L. A. SMOCK, 1980. Plankton communities of an acidic polymictic, brown-water lake. *Hydrobiologia* 69: 131-137
- STOUT, V. M., 1969. Lakes in the mountain region of New Zealand. *Verh. internat. Ver. Limnol.* 17: 404-413

- STOUT, V. M., 1972. Plankton composition in relation to nutrient inflow in a small New Zealand lake. *Verh. internat. Ver. Limnol.* 18: 605-612
- STOUT, V. M., 1975. A preliminary account of some lakes at different altitudes in southwest New Zealand. *Verh. internat. Ver. Limnol.* 19: 1452-1460
- STOUT, V. M., 1978. Effects of different silt loads of hydroelectric development on four large lakes. *Verh. internat. Ver. Limnol.* 20: 1182-1185
- STOUT, V. M., 1981. Some year to year fluctuations in a natural and in an artificial lake, South Island, New Zealand. *Verh. internat. Ver. Limnol.* 21: 699-702
- STOUT, V. M., 1984. Seasonal plankton cycles in two adjacent lakes in the South Island, New Zealand. *Verh. internat. Ver. Limnol.* 22: 1266-1230
- STOUT, V. M., 1991. Year to year fluctuations in a small mountain lake. *Verh. internat. Ver. Limnol.* 24: 1145-1148
- STRASKRABA, M., 1965. Contributions to the productivity of the littoral regions of pools and ponds. I. Quantitative study of the littoral zooplankton of the rich vegetation of the backwater Labicko. *Hydrobiologia* 26: 421-443
- STRAYER, D., 1985. The benthic micrometazoans of Mirror Lake, New Hampshire. *Arch. Hydrobiol./Suppl.* 72: 287-426
- SUDZUKI, M., 1960. Limnological survey of the Lake Nojiri on the Tableland in Middle Japan. II. Description of some new species of Rotifers. *Bull. Biogeogr. Soc. Japan* 21: 19-26
- SUDZUKI, M., 1964a. On the microfauna of the Antarctic Region. I. Moss-dwelling water communities at Langhovde. *Jap. Antarctic Res. Exp. Rep.* 19. 41 pp.
- SUDZUKI, M., 1964b. Ueber zwei neue *Encentrum*-Arten (Rotatoria) aus Moospolstern des Südpolar-landes. *Limnologica* 2: 349-353
- SUDZUKI, M., 1967. Rotifers from South Australia. I. A taxonomical note on *Keratella quadrata* from Tasmania. *Jap. Soc. syst. Zool.* 3: 17-19
- SUDZUKI, M., 1971a. Notes on the lava fauna of Mt. Fuji, in UENO, S. I.: *Results of the Co-operate Scientific Survey of Mt. Fuji*: 759-788. Fuji Kyuko Co. Ltd. Tokyo, Japan.
- SUDZUKI, M., 1971b. An analysis of colonoization in freshwater organisms I. Colonization at 17 stations along the 5 lakes of the Mt. Fuji. *Zool. Mag. Tokyo* 80: 191-201
- SUDZUKI, M., 1975. List of Rotifera and Gastrotricha from Garden Ponds of Tokyo 1974-1975 and some notes on *Rhinoglena*, *Fadeewella*, *Neogossea* etc. *Proc. Jap. Soc. syst. Zool.* 11: 5-12
- SUDZUKI, M., 1976a. Microscopical marine animals scarcely known from Japan. I. Micro- and meiofauna around Kasado Island in the Seto inland sea of Japan. *Proc. Jap. Soc. syst. Zool.* 12: 5-12
- SUDZUKI, M., 1976b. Recent portraits of Wild Biota in Japan. II. The inland sea of Japan around Kasado Islands, Yamaguchi Prefecture. *Obun Ronso* 7: 11-32
- SUDZUKI, M., 1977a. Some puzzling problems in the taxonomy of *Brachionus* and *Keratella*. *Arch. Hydrobiologia Beih.* 8: 230-231
- SUDZUKI, M., 1977b. Recent portraits of Wild Biota in Japan. III. A Man-made sandy beach constructed in the Bay of Tokyo. *Obun Ronso* 8: 221-239
- SUDZUKI, M., 1977c. Some approaches to the estimation of the biomass for microfauna communities. I. Analysis of the factors influencing the occurrence of microbiota inhabiting a beach forest, in *Biomass of animals of terrestrial systems in Japan*. Annual report of 1977: 159-198. Ed. Environmental agency, Tokyo
- SUDZUKI, M., 1978a. Recent portrait of Wild Biota in Japan. IV. Animalcules in the area of Mt. Fuji including 5 lakes, grasslands and forests at the foot of the mountain. *Obun ronso* 9: 235-277
- SUDZUKI, M., 1978b. Some approaches to the estimation of the biomass for microfauna communities. II. Difference in the occurrence of microbiota inhabiting litters, mosses, especially soils from four terrestrial ecosystems, in *Biomass of animals of terrestrial ecosystems in Japan*. Annual report of 1978: 181-215. Ed. Environmental agency, Tokyo
- SUDZUKI, M., 1979a. Some approaches to the estimation of the biomass for microfauna communities. III. Individual density and presumable biomass of the microbiota inhabiting terrestrial ecosystems in a subtropical Island, named Iriomote-Jima, in *Biomass of animals of terrestrial ecosystems in Japan*. Re-port of the result of 1978: 213-248. Ed. Environmental agency, Tokyo
- SUDZUKI, M., 1979b. On the microfauna of the antarctic region. III. Microbiota of the terrestrial interstices. *Memoirs Nat. Inst. Pol. Res.* Special issue n° 11: 104-126
- SUDZUKI, M., 1979c. Some aspects of the haline interstitial biota from Ryukyu Shoto, subtropical chain islands, southwest Japan. *Sesoko Mar. Sci. Lab. Techn. Rep.* 6: 37-50

- SUDZUKI, M., 1981a. Faunistic and ecological studies of the sewage biota of Japan (metropolis Tokyo and its adjacent prefecture). *Verh. internat. Ver. Limnol.* 21: 1094-1100
- SUDZUKI, M., 1981b. Periphytic biota (Aufwuchs) in sewage and park waters, in SUDZUKI, M. (ed.): *Some approaches to saprobiofical problems*: 69-107. Sanseido Co Ltd., Tokyo
- SUDZUKI, M., 1982. Recent portrait of wild biota in Japan. VII. Microscopical animals in Fukushima Prefecture. *Obun ronso* 12: 401-442
- SUDZUKI, M., 1985. Tasmanian Rotifers in Mid-August (winter). *Zool. Sci.* 2: 1005
- SUDZUKI, M., 1988a. Planktonic rotifers from Taiwan, R. O. C. *Zool. Sci.* 5: 1355
- SUDZUKI, M., 1988b. Comments on the antarctic rotifera. *Hydrobiologia* 165: 89-96
- SUDZUKI, M., 1989. Rotifera from the Oriental Region and their characteristics. *Special issue celebrating the Centennial Anniversary of the Foundation of "Nihon Daigaku" University Tokyo* 3: 325-366
- SUDZUKI, M., 1990. Summer rotifera from southwest islands of Japan. *Zool. Sci.* 7: 1185
- SUDZUKI, M., 1991a. Free-living microscopic animals of limestone caves Akiyoshi-dai, Yamaguchi, Japan. *Bull. Akiyoshi-dai Nat. Hist.* 26: 21-28
- SUDZUKI, M., 1991b. The rotifera from Singapore and Taiwan. *Proc. Jap. Soc. syst. Zool.* 43: 1-34
- SUDZUKI, M., 1991c. Rotifera from southwestern islands of Japan. III. *Zool. Sci.* 8: 1199
- SUDZUKI, M., 1992. New Rotifera from Southwestern Islands of Japan. *Proc. Japan. Soc. syst. Zool.* 46: 17-28
- SUDZUKI, M. & K. FUKUTA, 1989. Rotifera from Koyama-Ike, in Tottori, the largest coastal pond in Japan. *Zool. Sci.* 6: 1219
- SUDZUKI, M. & K. FUKUTA, 1992. Faunistic changes in the rotifera in the coastal prefectures. I. Tottori & Shimane-Ken. *Zool. Sci.* 9: 1297
- SUDZUKI, M. & Y. HOSOYAMA, 1991. Microscopic animals from Gyokusen-dô Cave and its water quality. *J. speleol. Soc. Japan* 16: 38-44
- SUDZUKI, M., Y. HOSOYAMA & Y. DEGUCHI, 1992. Occurrence of the rotifera and water quality. *Proc. Japan. Soc. syst. Zool.* 46: 71-99
- SUDZUKI, M., T. MATSUMOTO & K. HARITA, 1989. Occurrence of Rotifera in the field under natural and intentionally changed conditions. II. Lake Numasawa. *Hydrobiologia* 186/187: 247-254
- SUDZUKI, M. & B. V. TIMMS, 1977. A new species of *Brachionus* (Rotifera) from the Myall lakes, New South Wales. *Proc. Linn. Soc. NSW* 101: 162-166
- SUDZUKI, M. & B. V. TIMMS, 1980. Planktonic Rotifers of farm dams near Gloucester, NSW. *Bull. Austral. Soc. Limnol.* 7: 1-7
- SUDZUKI, M., K. WATANABE, K. SUZUKI & K. NARITA, 1983. Occurrence of Rotifera in the field under natural and intentionally changed conditions. *Hydrobiologia* 104: 341-347
- SUTHERLAND, I., 1982. A collection of zooplankton from Tuktoyaktuk Harbour, Northwest Territories. *Can. J. Zool.* 60: 477-480
- SYMOENS, J. J., L. DERONDE, I. VAN ELSLANDE, D. VAN DE GUCHT & H. BORGOO, 1983. Some limnological characteristics of a forest pond at Lembeek (Prov. Brabant, Belgium). *Abstract 22nd Congress SIL*: 336
- SZLANER, L., 1969. Response of lake fauna to light. *Pol. Arch. Hydrobiol.* 16: 233-244
- TAIT, R. D., R. J. SHIEL & W. KOSTE, 1984. Structure and dynamics of zooplankton communities, Alligator Rivers Region, N. T., Australia. *Hydrobiologia* 113: 1-13
- TAKAMURA, K., Y. SUGAYA, N. TAKAMURA, T. HANAZOTO, M. YASUNO & T. IWAKUMA, 1989. Primary production of phytoplankton and standing crops of zooplankton and zoobenthos in hypertrophic Lake Teganuma. *Hydrobiologia* 173: 173-184
- TAMAS, G., 1975. Controlled production of rotifer zooplankton in fingerlings ponds. *Hydrobiol. Jnl.* 11 (6): 53-55
- TARNOGRADSKY, D. A., 1961a. Microflora and microfauna of peats in the Caucasus. 5. *Sphagnetum* of Maharsky gouge (Karatchaivo-Circassian Republic). *Trav. Stat. biol. Cauc. nord* 22: 1-32
- TARNOGRADSKY, D. A., 1961b. Panzerrotatorien des Kaukasus - Genus *Lecane* Nitzsch. *Trav. Stat. biol. Cauc. nord* 22: 33-73
- TASSIGNY, M., R. MAILLARD & R. POURRIOT, 1970. Etude hydrobiologique d'un étang de Sologne: l'Etang de Pommereau. *Bull. Assoc. Nat. Orl. Loire moyenne*, N. S. n° 51: 3-36
- TATICCHI, M. I., 1968. Vicende stagionali delle comita littorance del Lago Trasimeno (1963-1965). *Riv. Idrobiol.* 7: 195-302
- TAYLOR, B. E. & D. L. MAHONEY, 1988. Extinction and recolonization: processes regulating zooplankton dynamics in a cooling reservoir. *Verh. internat. Ver. Limnol.* 23: 1536-1541

- TELESH, I. V., 1986. Comparative effectiveness of methods of counting plankton rotifers. *Hydrobiol. Jnl.* 22 (4): 101-104
- TENT, L., 1981. Der Aufwachs im Hamburger Hafen. Struktur einer Biocoenose in einem Belastungszentrum des Elbe-Aestuars. *Arch. Hydrobiol./Suppl.* 61: 1-58
- TEREK, J., 1990a. An annual cycle of zooplankton of dead branch Latorica near Leles (DFS). *Biol. Bratislava* 45: 81-94
- TEREK, J., 1990b. New and rare species of rotifer (Rotatoria) in Czechoslovakia. *Biol. Bratislava* 45: 149-152
- TEREK, J., 1990c. Net zooplankton of hydromelioration canals and its relation to selected ecological factors. *Biol. Bratislava* 45: 801-814
- THANE-FENCHEL, A., 1965. *Proales paguri*, n. sp., a rotifer living on the gills of the hermit crab *Pagurus bernhardus* (L.). *Ophelia* 3: 93-97
- THANE-FENCHEL, A., 1968. Distribution and ecology of non-planktonic brackish-water Rotifers from Scandinavian waters. *Ophelia* 5: 273-297
- THEILACKER, G. H. & M. F. Mc MASTER, 1971. Mass culture of the rotifer *Brachionus plicatilis* and its evaluation as a food for larval anchovies. *Mar. Biol.* 10: 183-188
- THOMAS, C. W., 1965. On populations in Antarctic meltwater pools. *Pacific Sci.* 19: 515-521
- THOMAS, C. W., 1972. Two species of Antarctic rotifers. *Antarct. Journ. U. S.* 7: 186-187
- THOMASSON, K., 1961. Zur Planktonkunde Spitsbergens 2. *Hydrobiologia* 18: 192-198
- THOMASSON, K., 1962. Planktological Notes from western North America. *Arkiv. Bot. Ser. 2, Bd. 4:* 437-463
- THOMASSON, K., 1963a. Algological notes. 2. An interesting *Staurastrum* population. *Revue algol.* 1: 93-99
- THOMASSON, K., 1963b. Araucarian lakes. *Acta Phytogeogr. Sueica* 47: 5-139
- THOMASSON, K., 1964a. Plankton and environment of North Patagonian Lakes. *Ann. Soc. Tart. ad res nat. invest. const. NS* 4: 9-28
- THOMASSON, K., 1964b. Algae from Lakes in Northern Colorado. *Svensk. Bot. Tidskr.* 58: 73-80
- THOMASSON, K., 1967. Phytoplankton from some lakes on Mt. Wilhelm, East New Guinea. *Blumea* 15: 285-296
- THOMASSON, K., 1970. A *Mallomonas* population. *Svensk. Bot. Tidskr.* 64: 303-311
- THOMASSON, K., 1971. Amazonian Algae. *Mém. Inst. r. Sci. nat. Belg., série 2, fasc. 86:* 57 pp.
- THOMASSON, K., 1973. Notes on the plankton of some Sidney reservoirs with description of two interesting desmids. *Contr. NSW Nat. Herb.* 4: 384-394
- THOMASSON, K., 1979. Heleoplankton from a pool in South Trøndelag Province, Central Norway. *Gunneria* 32: 7-21
- THRELKELD, S. T., 1983. Empty loricas and the dynamics of *Kellicottia longispina* in a subalpine, oligotrophic lake. *Hydrobiologia* 104: 367-372
- THRELKELD, S. T. & E. CHOINSKI, 1985. Plankton in a rapidly flushed impoundment: spatial distribution, population dynamics and community structure. *Mem. Ist. it. Idrobiol.* 43: 105-118
- TIEFENBACHER, L., 1972. Beiträge zur Biologie und Oekologie sessiler Rotatorien unter besonderer Berücksichtigung des Gehäusebaus und der Regenerationsfähigkeit. *Arch. Hydrobiol.* 71: 31-78
- TIMMS, B. V., 1981. Animal communities in three Victorian lakes of different salinity. *Hydrobiologia* 81/82: 181-193
- TIMMS, R. M. & B. MOSS, 1984. Prevention of growth of potentially dense phytoplankton populations by zooplankton grazing, in the presence of zooplanktonivorous fish, in a shallow wetland ecosystem. *Limnol. Oceanogr.* 29: 472-486
- TOKAT, M., 1976. Rotatoria of Lake Hazar (Gölcük) and their distribution. *Hydrobiol. Istanbul* 18: 1-13
- TONOLLI, V., 1962. L'attuale situazione del popolamento planctonico del Lago Maggiore. *Mem. Ist. it. Idrobiol.* 15: 81-134
- TORALES, G. J., W. H. HACK & B. TURN, 1972. Criaderos de Culicidos en Bromeliaceas del NW de Corrientes. *Acta zool. Lilloana* 29: 293-308
- TRAHMS, K.-J., 1972. Die Entwicklung der Plankton-Biocoenosen in Restgewässern des Rheinischen Braunkohlgebietes. *Internat. Rev. ges. Hydrobiol.* 57: 695-758
- TUNDISI, J. G., T. MATSUMARA-TUNDISI, M. C. CALIJURI & E. M. L. NOVO, 1991. Comparative limnology of five reservoirs in the middle Tietê River, S. Paulo State. *Verh. internat. Ver. Limnol.* 24: 1489-1496
- TURNER, P. N., 1986. Some rotifers from Republic of Korea. *Hydrobiologia* 137: 3-7

- TURNER, P. N., 1987a. Some rotifers from Alaska, USA, with a note to Researchers. *Microscopy* 35: 541-548
- TURNER, P. N., 1987b. Some rotifers from Kathmandu, Nepal. *Microscopy* 35: 549-550
- TURNER, P. N., 1987c. A new rotifer from a coastal lake in Southwestern Brazil: *Hexarthra longicornicula* n. sp. *Hydrobiologia* 135: 169-174
- TURNER, P. N., 1987d. *Keratella* rotifera found in Brazil, and a survey of *Keratella* rotifers from the Neotropis. *Amazoniana* 10: 223-236
- TURNER, P. N., 1988. Some rotifers encountered in Colorado, USA. *Microscopy* 36: 174-180
- TURNER, 1990a. Some rotifers from coastal lakes of Brazil, with description of a new rotifer, *Lepadella (Xenolepadella) curvicaudata*, n. sp. *Hydrobiologia* 208: 141-152
- TURNER, P. N., 1990b. Some interstitial rotifera from a Florida, USA, Beach. *Trans. am. Microscop. Soc.* 109: 417-421
- TURNER, P. N., 1990c. The rotifer genus *Platyias* Harring (1913) in the Neotropis. *Acta limnol. Brasil.* 3: 741-756
- TURNER, P. N. & C. DA SILVA, 1992. Littoral rotifers from the State of Mato Grosso, Brazil. *Environment* 27: 227-241
- TWORNBLY, S. & W. M. LEWIS Jr., 1987. Zooplankton abundance and species composition in Laguna la Orsinera, a Venezuelan floodplain lake. *Arch. Hydrobiol./Suppl.* 79: 87-107
- TZSCHASCHEL, G., 1979. Marine rotatoria aus dem Interstitial der Nordseeinsel Sylt. *Akad. Wiss. Literatur. Math.-Naturw. Klasse Reihe "Mikrofauna des Meeresbodens"* 71: 3-64
- TZSCHASCHEL, G., 1980. Verteilung, Abundanzdynamik und Biologie mariner interstitieller Rotatoria. *Mikrofauna des Meeresbodens* 81: 3-56
- TZSCHASCHEL, G., 1983. Seasonal abundance of psammon rotifers. *Hydrobiologia* 104: 275-278
- UENO, M., 1966. Freshwater zooplankton of Southeast Asia. *SE Asian Studies* 3: 94-109
- UENO, M., 1967. Zooplankton of Lake Titicaca on the Bolivian side. *Hydrobiologia* 29: 547-568
- UENO, M., R. HOSHINO, T. MITZUNO & M. TANAKA, 1961. A limnological study of the small lakes at the foot of Mt. Shiratakagama, west of the city of Yamagata. *Jap. J. Limnol.* 22: 31-43
- URABE, J., 1990. Stable horizontal variation in the zooplankton community structure of a reservoir maintained by predation and competition. *Limnol. Oceanogr.* 35: 1703-1717
- URABE, J., 1992. Midsummer succession of rotifer plankton in a shallow eutrophic pond. *J. Plankt. Res.* 14: 851-866
- USTAOGLU, R., 1986. Zooplankton (Metazoa) of the Karagöl (Jamanlar, Izmir-Turkey). *Biol. Gallo-Hell.* 12: 273-281
- USTAOGLU, R., in press. Zooplankton (Metazoa) of Lake Marmara (Turkey). *Biol. Gallo-Hell.*
- USTAOGLU, R. & S. BALIK, 1987. Rotifer fauna of Akgöl (Selçuk-Izmir) VIII. *Ulusal Biyoloji Kongresi Cilt II*: 614-626
- USTAOGLU, R. & S. BALIK, 1990a. Zooplankton of Lake Kus (Sandırma). X. *Ulusal Biyoloji Kongresi 18-20 Temmuz 1990, Erzutum*: 11-19
- USTAOGLU, R. & S. BALIK, 1990b. Zooplankton of Lake Gebekirse (Izmir-Turkey). *Rapp. Comm. Int. Mer Médit.* 32: 74
- VALLENTYNE, J. R., 1971. A description of Hutchinsons Lake, Ontario. *Limnol. Oceanogr.* 16: 473-475
- VALOVAYA, M. A., 1991. A new parasitic rotifer, *Albertia ovagranulata* sp. n. (Dicranophoridae) from the intestine of the Oligochaete *Enchytraeus albidus* from the White Sea sublittoral. *Hydrobiol. Jnl.* 27 (3): 58-61
- VANCIL, J. E., 1983. A method for the laboratory culture of the planktonic rotifer *Keratella cochlearis* (Gosse). *Hydrobiologia* 107: 47-50
- VAN DEN BOSCH, F. & J. RINGELBERG, 1985. Seasonal succession and population dynamics of *Keratella cochlearis* (Ehrb.) and *Kellicottia longispina* (Kellicott) in Lake Maarsseveen I (Netherlands). *Arch. Hydrobiol.* 103: 273-290
- VANDERBEKEN, C., 1977. Studie van de zooplanktonpopulaties in de "Nationale Watersportbaan" te Gent in de periode 1976-1977. *Lic. thesis, RUG*
- VAN DONK, E., R. D. GULATI & M. P. GRIMM, 1990. Restoration by manipulation in a small hypertrophic lake: first year results. *Hydrobiologia* 191: 285-295
- VAN ELSLANDE, I., 1975. Bijdrage tot de kennis van de secundaire produktie van een bosvijver te Lembeek. *Lic. thesis, VUB, Brussels*
- VANNI, M. J., 1986. Competition in plankton communities: suppression of small species by *Daphnia pulex*. *Limnol. Oceanogr.* 31: 1039-1056

- VAN TONGEREN, O. F. R., L. VAN LIERE, R. D. GULATI, G. POSTEMA & P. J. BOESEWINKEL-DE BRUYN, 1992. Multivariate analysis of the plankton communities of the Loosdrecht lakes: relationship with the chemical and physical environment. *Hydrobiologia* 233: 105-117
- VAN VOSSEL, A., 1973. Preliminair onderzoek van geschikte kweekcondities voor een zoetwater-rotifeer: *Epiphantes senta*. *Lic. thesis*, RUG
- VARDAPETYAN, S. M., 1972. Food relations of predatory crustaceans in lake zooplankton. *Sov. J. Ecol.* 3: 222-227
- VARGA, L., 1962. Rotatorien vom Hoverla-Berg. *Allat. Közlem.* 49: 141-150
- VARGA, L., 1963. Weitere Untersuchungen über die aquatile Mikrofauna der Boradla-Höhle bei Aggtelek (Ungarn). *Acta zool. Hung.* 9: 439-458
- VARNHAGEN, E., E. R. BYRON & Ch. R. GOLDMAN, 1988. *Epischura* density as a factor controlling the establishment of *Bosmina* populations in Lake Tahoe. *Verh. internat. Ver. Limnol.* 23: 2082-2086
- VASCOUCELOS, V., 1990. Seasonal fluctuations in the zooplankton community of Azibo reservoir (Portugal). *Hydrobiologia* 196: 183-191
- VASISHT, H. S. & R. JINDAL, 1980. Rheological survey of a pucca stream of Patiala (Punjab, India). *Limnologica* 12: 77-83
- VASQUEZ, E., 1984a. Estudio de las comunidades de Rotiferos del Orinoco medio, bajo Caroni y algunas lagunas de inundacion (Venezuela). *Mem. Soc. Cienc. Nat. La Salle* 121: 95-108
- VASQUEZ, E., 1984b. El zooplankton de la seccion baja de un rio de aguas negras (Rio Caroni) y de un embalse hidroelectrico (Macagua I), Venezuela. *Mem. Soc. Ciencias Nat. La Salle* 121: 109-129
- VASQUEZ, E. & W. KOSTE, 1988. Form variation of the rotifer *Brachionus variabilis* (Hempel, 1896) from the Orinoco River (Venezuela). *Annls. limnol.* 24: 127-129
- VASQUEZ, E., W. KOSTE & M. L. MEDINA, 1991. Variaciones morfologicas del rotifero *Keratella americana* (Carlin, 1943) de una laguna de inundacion del rio Orinoco, Venezuela. *Rev. Hydrobiol. trop.* 24: 83-90
- VASQUEZ, E. & M. L. MEDINA, 1991. Predation by dwarf forms of *Asplanchna sieboldi* (Rotatoria) from a floodplain lake of the Orinoco River (Venezuela). *Annls. Limnol.* 27: 111-117
- VASQUEZ, E. & J. REY, 1989. A longitudinal study of zooplankton along the Lower Orinoco River and its Delta (Venezuela). *Annls. Limnol.* 25: 107-120
- VASQUEZ, E. & J. REY, 1992. Composition, abundance and biomass of zooplankton in Orinoco floodplain lakes, Venezuela. *Annls. Limnol.* 28: 3-18
- VASS, K. K. & D. P. ZUTSHI, 1983. Energy flow, trophic evaluation and ecosystem management of a Kashmir Himalayan Lake (Dal Lake). *Arch. hydrobiol.* 97: 39-59
- VASS, K. K., A. WANGANEO, H. S. RAINA, D. P. ZUTSHI & R. WANGANEO, 1989. Summer limnology and fisheries of high mountain lakes of Kashmir Himalayas. *Arch. Hydrobiol.* 114: 603-619
- VASSILIEVA, G. L. & L. A. KUTIKOVA, 1969. Rotifères benthiques du genre *Notholca* dans le lac Baikal. *Zool. Zhu.* 48: 791-802
- VELASCO, J. L., 1990. Lista faunistica & bibliographia de los Rotiferos (Rotatoria) de la Peninsula Iberica e Islas Baleares & Canarias. *AEL*, Publ. 8. 195 pp.
- VELIKORETSJAKA, J. J. & L. F. FORSH, 1972. Lakes of different landscapes of the Kola Peninsula. *Verh. internat. Ver. Limnol.* 18: 517-521
- VENGLENIK, D. L., V. A. SOKOLOVA & M. M. TYAPTIRAYANOV, 1985. Zooplankton of lakes of Northeast Yakutia. *Hydrobiol. Jnl.* 21 (1): 68-73
- VERHEYE, H., 1980. Experimentele studie van de vertikale migratie bij zoetwaterplankton (Rotatoria - Cladocera - Copepoda). *Lic. thesis* RUG
- VERRAES, W., 1968. Hydrobiologisch onderzoek in Zeeschelde en Rupel ter hoogte van de Rupelmonding en in een visvijver te Bornem. *Natuurwet. Tijdschrift* 50: 172-173, 9 figs.
- VERRETH, J., 1990. The accuracy of population density estimates of a horizontally distributed zooplankton community in Dutch fish ponds. *Hydrobiologia* 203: 53-61
- VIETINGHOFF, U., N. ERDMANN, H. ARNDT, V. KELL & M. L. HUBERT, 1984. Integrated samples provide accurate means for parameters characterizing aquatic ecosystems. *Internat. Rev. ges. Hydrobiol.* 69: 121-131
- VIETINGHOFF, U., M. L. HUBERT, R. HEERKLOSS, H. ARNDT & W. SCHNESE, 1984. A mathematical submodel for the planktonic rotatoria in the ecosystem of the Barther Bodden (Southern Baltic Sea). *Internat. Rev. ges. Hydrobiol.* 69: 159-172

- VILA CLARA, G. & V. SLADECEK, 1989. Mexican rotifers as indicators of water quality, with description of *Collotheca riverai*, n. sp. *Arch. Hydrobiol.* 115: 257-263
- VILLEGAS, C. T., 1990. The effects on growth and survival of feeding water fleas (*Moina macrocopa* Straus) and rotifers (*Brachionus plicatilis*) to milkfish fry (*Chanos chanos* Forskall). *Isr. J. Aquacult.* 42: 10
- VOIGT, M., 1957. Rotatoria. Die Rädertere Mitteleuropas. Berlin, Nikolaussee. 2 Teile: 508 pp., 115 plates.
- VOLGIN, M. V., M. S. KUKOU, L. L. SIPKO & M. V. TIMOFYEVA, 1982. Hydrobiology of the Cooling ponds of the Barabinskaya Power Plant. *Hydrobiol. Jnl.* 18 (5): 46-51
- VORSTMAN, A. G., 1960. Hydrobiologisch onderzoek van het Meyendelgebied. 12e verslag.
- VOS, J., 1975. Limnologische studie van de hypertrofiering van het zuidelijk gedeelte van het Donkmeer (O. Vlaanderen) tengevolge van het inpompen van grondwater. *Lic. thesis UIA*, Antwerpen
- VOS-KELK, P. & C. DAVIDS, 1977. A short note on *Lecane* species (Rotifera) found in *Utricula minor* vegetation. *Hydrobiol. Bull.* 11: 53-55
- V'YUSHKOVA, V. P., 1985. Principal zooplankton species in Volgograd and Saratov Reservoirs. *Hydrobiol. Jnl.* 21 (5): 15-19
- V'YUSHKOVA, V. P. & V. A. LAKHNOVA, 1974. Rotifers of the Volgograd Reservoir. *Hydrobiol. Jnl.* 10 (1): 66-69
- WAEYAERT, R., 1973a. Verwerking van een staalname uit de Samber in Frankrijk. *Maandwerk Lab. Morf. Syst. RUG*
- WAEYAERT, R., 1973b. Studie van enige eutroficate- en pollutieverschijnselen in het Donkmeer (O. Vl.). *Lic. thesis RUG*
- WALLACE, R. L., 1975. Larval behavior of the sessile rotifer *Ptygura beauchampi* (Edmondson). *Verh. internat. Ver. Limnol.* 19: 2811-2815
- WALLACE, R. L., 1977. Distribution of sessile rotifers in an acid bog pond. *Arch. Hydrobiol.* 79: 478-505
- WALLACE, R. L., 1980. Ecology of sessile rotifers. *Hydrobiologia* 73: 181-193
- WALLACE, R. L. & W. T. EDMONDSON, 1986. Mechanism and adaptive significance of substrate selection by a sessile rotifer. *Ecology* 67: 314-323
- WALLACE, R. L. & P. L. STARKWEATHER, 1983. Clearance rates of sessile rotifers: *in situ* determinations. *Hydrobiologia* 104: 379-383
- WALLACE, R. L. & P. L. STARKWEATHER, 1985. Clearance rates of sessile rotifers: *in vitro* determinations. *Hydrobiologia* 121: 139-144
- WALSH, E. J., 1989. Oviposition behavior of the littoral rotifer *Euchlanis dilatata*. *Hydrobiologia* 186/187: 157-161
- WALTER, G., 1961. Die aktive und passive Einwanderung tierischer und pflanzlicher Organismen in einem neuangelegten Durchfluszgraben. *Verh. internat. Ver. Limnol.* 14: 508-512
- WALTER, G. & R. SCHARF, 1961. Das biologische Gütebild der unteren Oder und der Lausitzer Neisse. *Internat. Rev. ges. Hydrobiol.* 46: 130-161
- WALZ, N., 1983a. Individual culture and experimental population dynamics of *Keratella cochlearis* (Rotatoria). *Hydrobiologia* 107: 35-45
- WALZ, N., 1983b. Continuous culture of the pelagic rotifers *Keratella cochlearis* and *Brachionus angularis*. *Arch. Hydrobiol.* 98: 70-72
- WALZ, N., 1986. Wie werden Rotatorienpopulationen reguliert? Beziehungen zwischen Rotatorien, Crustaceen und Phytoplankton. Akademie für Naturschutz und Landschaftspflege. Limnol. Station Seeon. *Laufener Seminarbeiträge* 2/86: 80-90
- WALZ, N., H.-J. ELSTER & M. MEZGER, 1987. The development of the rotifer community structure in Lake Constance during its eutrophication. *Arch. Hydrobiol./Suppl.* 74: 452-487
- WARD, J. V., 1975. Downstream fate of zooplankton from a hypolimnia release mountain reservoir. *Verh. internat. Ver. Limnol.* 19: 1798-1804
- WATTIEZ, C., 1978. Agrégation et migration verticale du zooplancton dans de petits étangs peu profonds. *Hydrobiologia* 61: 49-67
- WAWRİK, F., 1966. Zur Kenntnis alpiner Hochgebirgs-Kleingewässer. *Verh. internat. Ver. Limnol.* 16: 543-553
- WAWRİK, F., 1968. Zur Frage: Führt der Donaustrom autochtones Plankton, II. *Arch. Hydrobiol./Suppl.* 34: 339-363
- WAWRİK, F., 1969. Braunwasserteich-Studien im niederösterreichischen Waldviertel. *Verh. internat. Ver. Limnol.* 17: 570-574

- WAWRIK, F., 1972. Vergleichende Braunwasser-Teichstudien im niederösterreichischen Waldviertel. *Hydrobiologia* 39: 17-82
- WEIMIN, C., 1988. Some aspects of zooplankton species, biomass and distribution to environmental factors in Dianchi and Erhai Lakes with an evaluation of water trophic status. *Verh. internat. Ver. Limnol.* 23: 566
- WENINGER, G., 1971. Das Auftreten kleiner Metazoen bei Abbauprozessen in vergleichender Sicht. *Sitzungsber. Oesterr. Akad. Wiss. Abt. 1*, 179: 129-158
- WEU, Y. H., 1992. Contribution of bacterioplankton, phytoplankton, zooplankton and detritus to organic seston carbon load in a Changjiang floodplain lake (China). *Arch. Hydrobiol.* 126: 213-238
- WHITMAN, R. L., A. V. GOCHEC & P. L. RUCKMAN, 1988. Biological assemblages of Miller Woods ponds, Indiana Dunes National Lakeshore, Gary, Indiana. *Verh. internat. Ver. Limnol.* 23: 1041-1048
- WHYTE, J. N. C. & W. D. NAGATA, 1990. Carbohydrate and fatty acid composition of the Rotifer *Brachionus plicatilis*, fed monospecific diets of yeast or zooplankton. *Aquaculture* 89: 263-272
- WIBAUT-ISEBREE MOENS, N. L., 1964. Onderzoek van grindgaten langs de Maas. *Publ. natuur-hist. Gen. Limb.* 14: 1-44
- WIKTOR, J., 1960. Outline of biological conditions of the Szecezecin Lagoons waters. *Polsk. Arch. Hydrobiol.* 7: 7-27
- WIKTOR, K., 1968. Seasonal modifications in fertility of some Rotifer species. *Polsk. Arch. Hydrobiol.* 15: 9-22
- WILLEN, T., 1975. Biological long-term investigations of Swedish lakes. *Verh. internat. Ver. Limnol.* 19: 1117-1124
- WILLIAMS, W. D., 1991. Chinese and Mongolian saline lakes: a limnological overview. *Hydrobiologia* 210: 39-66
- WILLIAMSON, C. E., 1984. Laboratory and field experiments on the feeding and ecology of the cyclopoid copepod, *Mesocyclops edax*. *Freshw. Biol.* 14: 575-585
- WILLIAMSON, C. E., 1987. Predator-prey interactions between omnivorous diaptomid copepods and rotifers. The role of prey morphology and behavior. *Limnol. Oceanogr.* 32: 167-177
- WILLIAMSON, C. E. & N. M. BUTLER, 1986. Predation on rotifers by the suspension-feeding calanoid copepod *Diaptomus pallidus*. *Limnol. Oceanogr.* 31: 393-402
- WILLIAMSON, C. E. & J. J. GILBERT, 1980. Variation among zooplankton predators: the potential of *Asplanchna*, *Mesocyclops* and *Cyclops* to attack, capture and eat various rotifer prey, in KERFOOT, W. Ch.: *Evolution and ecology of zooplankton communities*: 509-517. Ed. University Press of New England, Hanover. 793 pp.
- WILLIAMSON, C. E., M. E. STOECKEL & L. J. SCHONECK, 1989. Predation risk and the structure of freshwater zooplankton communities. *Oecologia* 79: 76-82
- WILMS, A. L., R. D. GULATI & G. POSTEMA, 1991. First attempt to measure the clearance rate of *Anuraeopsis fissa*. *Verh. internat. Ver. Limnol.* 24: 742-744
- WISSMAR, R. C. & R. G. WETZEL, 1978. Analysis of the North American lake ecosystems. VI. Consumer community and production. *Verh. internat. Ver. Limnol.* 20: 587-597
- WISZNIEWSKI, J., 1954. Matériaux relatifs à la nomenclature et à la bibliographie des Rotifères. *Polski Arch. Hydrobiol.* 2/15: 7-249
- WRIGHT, H. G. S., 1961. A new species and a new subspecies of the Rotatorian genus *Collotheca*. *J. Quak. micr. Club*, Ser. 4, vol. 5, nr. 13: 361-365
- WROBEL, S., 1972. Some remarks to the production of basic communities in ponds with inorganic fertilization. *Verh. internat. Ver. Limnol.* 18: 221-227
- WULFERT, K., 1960a. Die Rädertiere saurer Gewässer der Dübener Heide. I. Die Rotatorien des Zadlitzmoors und des Wildenhainer Bruchs. *Arch. Hydrobiol.* 56: 261-298
- WULFERT, K., 1960b. Die Rädertiere saurer Gewässer der Dübener Heide. II. Die Rotatorien des Krebsscherentümpels bei Winkelmühle. *Arch. Hydrobiol.* 56: 311-333
- WULFERT, K., 1961. Die Rädertiere saurer Gewässer der Dübener Heide. III. Die Rotatorien des Presseler und des Winkelmühler Teiches. *Arch. Hydrobiol.* 58: 72-102
- WULFERT, K., 1966. Rotatorien aus dem Stausee und der Trinkwasserbereitung der Stadt Baroda (Indien). *Limnologica* 1: 53-93
- WULFERT, K., 1968. Rädertiere aus China, I. *Limnologica* 6: 405-416
- YAMAMOTO, K., 1960a. Plankton rotatoria in Japanese inland waters. *Hydrobiologia* 16: 364-411
- YAMAMOTO, K., 1960b. The Rotifers found in the plankton samples of Lake Biwa-ko. *Jap. J. Limnol.* 21: 327-334

- YAMAMOTO, K., 1967. First report of the regular limnological survey of Lake Biwa. III. Zooplankton. *Mem. Fac. Sci., Kyoto Univ., Ser. Biol.* 1: 62-77
- YAMAMOTO, K., 1968. Second report of the regular limnological survey of Lake Biwa (1967). II. Zooplankton. *Mem. Fac. Sci., Kyoto Univ., Ser. Biol.* 2: 92-106
- YAMAZI, I., 1960. Plankton investigations in inlet waters along the coast of Japan. IV. The plankton of Nanao Bay. *Seto Mar. Biol. Lab. Contr.* 191: 309-319
- YAN, N. D. & W. GEILING, 1985. Elevated planktonic rotifer biomass in acidified metal-contaminated lakes near Sudbury, Ontario. *Hydrobiologia* 120: 199-205
- YEREKEYEV, U. Ye. & S. K. KAZAKHBAYEV, 1972. Zooplankton of Aral Sea spawning grounds (Mujak, Adjhibay & Drzytphas). *Hydrobiol. Jnl.* 8 (3): 67-70
- YUFERA, M., 1982. Morphometric characterization of a small-sized strain of *Brachionus plicatilis* in culture. *Aquaculture* 27: 55-61
- YUFERA, M., L. M. LUBIAU & E. PASCUAL, 1984. Estudio del zooplancton de las Salinas de Cadiz. *Limnetica* 1: 62-69
- ZAGARESE, H. E. & M. C. MARINONE, 1992. Induction and inhibition of spine development in the rotifer *Keratella tropica*: evidence from field observations and laboratory experiments. *Freshw. Biol.* 28: 289-300
- ZANKAI, N. P., 1968. Ueber die Rädertieren - (Rotatoria) fauna des Plattensees, nach Literaturangaben von 1897 bis 1960. *Ann. biol. Tihany* 35: 247-272
- ZANKAI, N. P., 1984. Predation of *Cyclops vicinus* (Ent., Crustacea) on small zooplankton animals in Lake Balaton (Hungary). *Arch. Hydrobiol.* 99: 360-378
- ZANKAI, N. P., 1989. Horizontal distribution of rotifer plankton along a trophic gradient in Lake Balaton: changes of community structure and abundance during the past 20 years. *Arch. Hydrobiol.* 115: 111-123
- ZANKAI, N. P. & G. KERTESZ, 1967. Horizontal plankton investigations in Lake Balaton. VI. A study of the open water Rotatoria of Lake Balaton, based on collections in 1965. *Ann. biol. Tihany* 34: 255-275
- ZANKAI, N. P. & J. E. PONYI, 1970. The quantitative proportions Rotifera plankton in Lake Balaton, in 1967. *Ann. biol. Tihany* 37: 291-308
- ZANKAI, N. P. & J. E. PONYI, 1971. The horizontal distribution of rotifera plankton in Lake Balaton. *Ann. biol. Tihany* 38: 285-304
- ZANKAI, N. P. & J. E. PONYI, 1972. Quantitative relationships of the rotatoria plankton in Lake Balaton during 1965-1966. *Ann. biol. Tihany* 39: 189-240
- ZANKAI, N. P. & J. E. PONYI, 1973. The biomass of rotatoria in Lake Balaton. *Ann. Biol. Tihany* 40: 285-292
- ZARFDJIAN, M. H., M. VRANOVSKY & P. S. ECONOMIDIS, 1990. Les invertébrés planctoniques du Lac Volvi (Macédoine, Grèce). *Internat. Rev. ges. Hydrobiol.* 75: 403-412
- ZAYTSEV, Yu. P., G. P. GARKAVAYA, D. A. NESTROVA & L. N. POLISHCHUK, 1989. The Danube as the principal source of eutrophication of the Black Sea. *Hydrobiol. Jnl.* 25 (4): 18-20
- ZEEB, Ia. Ia., 1972. Die limnologischen Untersuchungen der Dnjeper-Stauseen (Hauptergebnisse und Perspektive). *Verh. internat. Ver. Limnol.* 18: 848-853
- ZHADIN, V. I., N. A. AKATOVA, L. A. KUTIKOVA & N. G. OZERETSKOVSKAYA, 1972. Fresh-water psammon on the shoreline of the Courland Spit. *Hydrobiol. Jnl.* 8 (6): 50-63
- ZHAO, Y. & C. E. KING, 1989. Ecological genetics of the rotifer *Brachionus plicatilis* in Soda Lake, Nevada, USA. *Hydrobiologia* 185: 175-181
- ZIMMERMANN, C., 1974. Die pelagischen Rotatorien des Sempachersees, mit spezieller Berücksichtigung der Brachioniden und der Ernährungsfrage. *Schweiz. Z. Hydrol.* 36: 205-300
- ZINEVICI, V. & L. TEODORESCU, 1990. L'évolution de la structure taxonomique du zooplankton dans les écosystèmes de type lacustre du delta du Danube, sous l'action du facteur anthropique (pendant les années 1975-1987). *Rev. roum. biol.-Biol. anim.* 35: 69-81
- ZINEVICI, V. & L. TEODORESCU, 1991. L'évolution de la structure gravimétrique du zooplankton dans les écosystèmes de type lacustre du delta du Danube, sous l'impact du processus d'eutrophisation (1975-1987). *Rev. roum. biol.-Biol. anim.* 35: 155-167
- ZIVKOVIC, A., 1968. Das Zooplankton der Jugoslawischen Donaustrecke. *Arch. Hydrobiol./ Suppl.* 34: 155-167
- ZIVKOVIC, A., 1987. Fauna Rotatoria in the yugoslav part of the Danube river and its flood-regions at Apatin, with description of three new species. *Proc. Fauna SR Serbia* 4: 1-110
- ZOUFAL, W., 1989. Developmental times of *Synchaeta oblonga* eggs from the Danube, Austria.

- ZUNIGA, L. R., V. CAMPOS, H. PINOCHET & B. PRADO, 1991. A limnological reconnaissance of Lake Tebenquiche, Salar de Atacama, Chile. *Hydrobiologia* 210: 19-24
- ZUREK, R., 1980. The effect of suspended materials of the zooplankton. *Acta hydrobiol.* 22: 449-471
- ZUTSHI, D. P., V. KAUL & K. K. VASS, 1972. Limnology of high altitude Kashmir lakes. *Verh. internat. Ver. Limnol.* 18: 599-604
- ZUTSHI, D. P., B. A. SUBLA, M. A. KHAN & A. WANGANEW, 1980. Comparative limnology of Jammu and Kashmir Himalayas. *Hydrobiologia* 72: 101-112